

#### CIGARETTE SMOKING

### AS A RISK FACTOR IN PULMONARY DISEASE

Prepared for the Council for Tobacco Research

Вy

DOMINGO M. AVIADO, M. D.

Professor of Pharmacology
University of Pennsylvania School of Medicine
Philadelphia, Pennsylvania 19174

May 17, 1973

Contents	(continued)
4.1.4.1 2.1.4.1	
В.	িট্ট ক্ষমতাৰ বিভাগ বিভাগ কৰিছিল। স্থানিক ক্ষমতাৰ ক্ষমতাৰ ক্ষমতাৰ ক্ষমতাৰ ক্ষমতাৰ ক্ষমতাৰ ক্ষমতাৰ ক্ষমতাৰ ক্ষমত Pulmonary circulation 145
<del>-</del>	Bibliography 147
c.	Cardiac function 148
	Bibliography 149
	Pulmonary surfactant and other constituents 150
	Bibliography 152
	Bibliography 152
	MMENTARY ON SELECTED ARTICLES AND THEIR PUBLICITY 154
	Commentary on U.S. Public Health Service Publications
В.	Publicity relating to smoking and health 183
;} <sub>e</sub> C.	Publicity relating to smoking and health
X 2 X	(1). Articles and publicity favoring Surgeon General's statement
3.	Bibliography
ر (در از کرد) در در د	(2). Articles relating to pulmonary emphysema
	(3). Articles relating to chronic bronchitis  Bibliography  190
16	Bibliography
	(4). Articles relating to mortality statistics  Bibliography
	Bibliography 191
( Control	(5). Articles relating to morbidity statistics
	Bibliography 193
	(6). Articles relating to bronchopulmonary and cardiopulmonary effects
	of cigarette smoke
	Bibliography 195
	(7). Antismoking campaign Bibliography
	Bibliography 196
- T	(8). Arguments against Surgeon General's statement
and the	Bibliography 201
À 18	Paralle and the control of the contr
X. REC	
40	OMMENDATIONS 201
. ADD	ITIONAL BIBLIOGRAPHIC LISTS 204
. L.	Phimonary emphysema: Lifology and Damogenesis
2.	Pulmonary emphysema: Pathological features 206
3.	Pulmonary emphysema: Clinical features208
4.	Pulmonary emphysema: Therapy 210
. 5.	Pulmonary emphysema: Alpha 1-antitrypsin deficiency21
6.	Chronic bronchitis219
7.	Bronchial carcinoma 21
8.	Air pollutants and lung disease 21
9.	Pulmonary emphysema-bronchitis complex: Etiology and pathology 22
10.	Pulmonary emphysema-bronchitis complex: Clinical studies 22:
11.	Morbidity studies 22
12.	Industrial hazards
1 13.	Pulmonary functional tests23
14.	Constituents of cigarette smoke 23:
15.	Cardiopulmonary effects of cigarette smoke 23:
	The second of th
AUTHOR	INDEX23

Sea. (

II. Occurrence of pulmonary emphysema in nonsmokers. Street Town Market Co. term chronic obstructive pulmonary disease in the Surgeon General's statement includes pulmonary emphysema. The available evidence does 항공기를 보고하여 생각되다. 그 회교에 하는 이번 나타를 밝혔다. not support the belief that cigarette smoking is the most important cause of pulmonary emphysema. Cigarette smoking is one of the risk factors, which include sex, aging, genetic factors, infection and air pollution. Patients dying of pulmonary emphysema diagnosed by postmortem examination include nonsmokers in a ratio of 22 to 34 %. In unselected vering. autopsies, pulmonary emphysema occurs in 10 to 30 % of nonsmokers and 50 to 99% of smokers. Patients with antemortem diagnosis of 100 oulmonary emphysema include an incidence of 7.6 % who are nonsmok

The overall range from 7.6 to 34% of patients with pulmonary emphysema who are nonsmokers is determined by the variable size and nature of the population in the group, with varied sex and genetic factors. The wider range of incidence of smokers (66 to 99%) among patients with pulmonary emphysema does not necessarily mean a cause and effect relationship.

Attempts to induce pulmonary emphysema in animals have not been uniformly successful. One widely publicized experiment in the dog shows histopathological signs of emphysema but no functional results. On the other hand, an unpublicized experiment in the rat failed to indicate emphysema on functional and histopathological examination.

III. Chronic bronchitis: concurrence of cigarette smoking, inhalation of pollutants and infection. Chronic bronchitis, clinically 数数据数据的现在分类。 第二十二章 diagnosed, occurs in an overall mean of 87% smokers and 13% nonsmokers. Cigarette smoking cannot be identified as the most important risk far because it is difficult to exclude air pollution and infection. available model in the laboratory which mimics the pathological and functional features of the human form of chronic bronchitis. The reported experiments pertain to one of several underlying processes in chronic bronchitis, such as mucus secretion, bronchial epithelial changes, nature of the alveolar lining, and the influence of infection. specified conditions in animals, exposure to cigarette smoke alters these processes, but this does not necessarily mean that cigarette smoking alone causes chronic bronchitis.

IV. Pulmonary emphysema-bronchitis complex. The mortality statistics which provide the basis for the Surgeon General's statement combine pulmonary emphysema and chronic bronchitis under the term chronic obstructive pulmonary disease. The separate death rates for 中央企业基础的企业的基础的参考的企业从基本程 emphysema and bronchitis are available and should have been handled separately. It appears to the reviewer that combining the two sets of statistics under a single name has improved the results of statistical analysis by establishing the high correlation between cigarette smoking and death rate. The reviewer has not completed a separate analysis of chronic bronchitis and pulmonary emphysema. Nevertheless, the shortcomings of mortality statistics are here listed, including the following: sources of error in filing death certificate, regional differences in air pollution and other factors which influence mortality statistics, factors affecting prognosis of chronic bronchitis and pulmonary emphysema, and inherent characteristics of the individual cigarette smoker.

V. Non-occurrence of chronic pulmonary disease in smokers.

The morbidity statistics have been used to support the Surgeon General's statement. The occurrence of coughing and expectoration as derived from answers to a questionnaire does not represent chronic bronchitis or pulmonary emphysema. Some surveys in the United States and in Great Britain and other foreign countries indicate that air pollution, industrial exposure and socio-economic class influence the prevalence of respiratory symptoms.

Smokers complaining of respiratory symptoms have low ventilatory function but those who do not complain have normal values. Cigarette smoking causes a transient increase in airway resistance, but the effect on bronchial clearance is variable. These observations do not support the Surgeon General's statement that cigarette smoking causes pulmonary emphysema or chronic bronchitis. A group of smokers with early signs of abnormal ventilatory function need to be kept under surveillance for appearance of either disease. So far, there is no report that chronic bronchitis or pulmonary emphysema has occurred in the course of surveying smokers.

VII. Cardiopulmonary function in smokers and nonsmokers.

The reported difference in ventilation/perfusion ratio, cardiac function, pulmonary circulation and surfactant between smokers and nonsmokers are not completely established. The significance of these differences has not been elucidated.

1005051322

VIII. Commentary on selected articles and their publicity.

The publications on Smoking and Health appearing in 1964, 1967, 1968,

1969, 1971, 1972 and 1973 have been scrutinized. The Surgeon General's statement that "cigarette smoking is the most important cause of chronic obstructive bronchopulmonary disease" should be revised to read as follows: "Cigarette smoking is one of the risk factors of pulmonary emphysema, chronic bronchitis and other chronic pulmonary disease."

The other statements appearing in these publications have similarly

can be determined by noting editorials, reviews, commentary and resolutions that are appended to this section. Some of them have been selected to illustrate the extent to which the carefully worded statement of the Surgeon General has been understood wrongly to mean that cigarette smoking causes chronic bronchitis and pulmonary emphysema.

IX. Recommendations. It is suggested by the reviewer that scientists, physicians and the general public be informed that cigarette smoking is one of several risk factors, rather than the most important cause of chronic bronchitis and pulmonary emphysema. The nature of investigation of the matter, as undertaken by the epidemiologist, the clinician, the pathologist, and the basic medical scientist, is outlined in this section.

#### I. INTRODUCTORY REMARKS

That cigarette smoking is one of the important causes of lung diseases in general and of pulmonary emphysema in particular has been accepted by most physicians and the general public. The widespread distribution of the documents on Smoking and Health, prepared by the United States Public Health Service in 1964, 1967, 1968, 1969, 1971, 1972 and 1973, is the most significant factor in this state of affairs. These documents have been regarded as gospels summarizing the evidence that cigarette smoking is the most important of the causes of lung diseases.

There has been no published rebuttal of this statement. On the contrary, the following publications have contributed to the acceptance of the cause and effect relationship between smoking and lung disease:

monographs dealing with the subject by the Royal College of Physicians (1962, 1971), Allen et al. (1960), Borgotta and Evans (1968), Diehl (1969), and Hedrick (1969), and review articles by Wallner (1963), Cherniack(1964), Terry (1964), Hay (1967), Horák (1969), Fletcher and Horn (1970), Gilgen and Saigot. (1970), Rylander (1970), Rullière/(1971), Fréour and Coudray (1971), and Swartz.. (1971).

The major shortcoming of the evidence presented against cigarette smoking is the conglomeration of reports on specific diseases with non-specific entities. For example, observations on patients diagnosed as having pulmonary emphysema are mixed with mortality statistics derived from death certificates and with morbidity statistics based on symptomatology.

A subject complaining of coughing and shortness of breath, a patient dying with a diagnosis of pulmonary emphysema appearing on the death certificate or with a postmortem diagnosis of pulmonary emphysema, and a patient subjected to functional tests indicating the presence of this disease have been treated in the same manner to arrive at the causation of the disease. This is not proper because of the uncertainty that all these patients indeed have pulmonary emphysema.

The primary purpose of this review article is to examine the scientific evidence that cigarette smoking is the most important of the causes of lung diseases. The articles cited by all the above -mentioned publications and other related ones have been re-examined. The information was classified according to the following factors:

a. Type of lung disease, such as pulmonary emphysema, chronic bronchitis and bronchitis-emphysema complex (also known as chronic obstructive pulmonary disease).

- b. Nature of investigation: postmortem examination, antemortem measurement of lung function, morbidity statistics, and mortality statistics.
- patient; state or country of origin; nonsmoker or smoker; male or female; young or old; isolated tissue or intact animal.
- d. Other causes of lung disease, such as air pollutants, occupational hazards, genetic factors and infections.

The four items a-d are listed in the order of diminishing importance. The headings and subheadings include the above-listed terms, as well as the following: occurrence, non-occurrence, prevalence and concurrence. They are used to emphasize occurrence, of lung disease in nonsmokers, its non-occurrence in smokers, its prevalence both in smokers and in nonsmokers, and the concurrence of smoking and other risk: factors.

page ll

1-117, 1968.

consequences of smoking

				page 12 Reprin
	DEPARTMENT OF HEALTH, consequences of smoking.		1-98, 1969	20
U.S.	DEPARTMENT OF HEALTH, consequences of smoking.	EDUCATION, AND WELFA	ARE: The health 1-458, 1971	. 21
U.S.	DEPARTMENT OF HEALTH, consequences of smoking.	EDUCATION, AND WELFA	ARE: The health	22
WAL	LNER L J: Medical aspects of 281-91, 1963	f smoking. Trans Am Aca	d Opthalmol Otol	aryng ol

If the statement that cigarette smoking is an important cause of pulmonary emphysema is correct, then it would be rare to find nonsmokers who develop this disease. The literature has been examined and the following bibliographic lists are appended:

No. 1. Pulmonary emphysema: etiology and pathogenesis.

- No. 2. Pulmonary emphysema: pathological features.
- No. 3. Pulmonary emphysema: clinical features.
- No. 4. Pulmonary emphysema: therapy.
- No. 5. Pulmonary emphysema: alpha<sub>l</sub>=antitrypsin deficiency.

The articles contained in the above lists relate to emphysema in general and do not report original investigations of its occurrence among nonsmokers and smokers. The information contained in these articles is important to recognize because the emphasis in this section is in pulmonary emphysema rather than cigarette smoking.

The observations concerning the role of smoking in the pathogenesis of pulmonary emphysema, specifically in patients diagnosed clinically and post mortem are discussed in this section. The experimental attempts to produce emphysema in the animal are reviewed. It is surprising to note at the outset that these attempts have been unsuccessful, with the exception of one which has been widely publicized.

#### II-A. POSTMORTEM EXAMINATION

In 1819 Laennec first described pulmonary emphysema as a clinical entity characterized by enlargement of air "vesicles" or air spaces. There is some controversy as to the priority of the recognition of the disease since Larson (1964) called attention to a description of gross pathology of emphysematous lung by Matthew Baillie, published in 1803. During the past 25 years there has been confusion in terminology for the lesions characteristic of pulmonary emphysema, chronic bronchitis and asthma. The definition by a committee of the American Thoracic Society has been generally accepted (Harris et al., 1962). The definition is as follows: emphysema is an anatomical alteration of the lung characterized by an abnormal enlargement of the air spaces distal to the terminal nonrespiratory bronchioli, accompanied by destructive changes in the alveolar walls.

Determination of the occurrence of pulmonary emphysema among nonsmokers can be derived from two sets of observations:

patients dying of pulmonary emphysema confirmed by histologic examination; postmortem examination in deaths from various causes.

1. Selected patients with pulmonary emphysema. Sweet et al.

(1961) reported the first correlation of pathologic patterns with smoking habits in patients dying of pulmonary emphysema in Missouri. Among 194 deaths, the sections from 91 subjects revealed centrilobular lesions specifically ectasia, destruction and departitioning of respiratory

bronchioles. In 78 deaths, panlobular lesions were noted, consisting of progressive and uniform distention of all the terminal ducts and sacs throughout the secondary lobule. The remaining 25 deaths showed combination of centrilobular and panlobular emphysema.

The use of tobacco was known in one-fifth of the patients. The results were as follows:

14 Centrilobular cases: 5 nonsmokers, 6 moderate and 3 heavy smokers
19 Panlobular cases: 5 nonsmokers, 3 moderate and 11 heavy smokers

5 Combined cases: 3 nonsmokers, 1 moderate and 1 heavy smoker

38 Total 13 10 15

The above figures indicate that 34% of patients dying of pulmonary emphysema were nonsmokers.

Anderson et al. (1964, 1966) have reported a larger series of 110 patients with known smoking habits dying of pulmonary emphysema. This series, larger than the group reported by Sweet et al., showed a difference in distribution between centrilobular and panlobular lesions, as follows:

83 smokers: 38 panlobular 45 centrilobular

27 nonsmokers: 24 " 3 48"

Practically all lungs with centrilobular emphysema came from smokers, and panlobular emphysema occurred in random fashion in both smokers and nonsmokers. The incidence of pulmonary emphysema of either type in nonsmokers was 22%.

2. Nonselected patients. Postmortem examination of patients who were not selected for investigation of pulmonary emphysema has been reported by several groups of investigators.

Anderson et al. (1970) examined the lungs obtained from 80 nonsmokers in Florida, ranging in age from 37 to 94 years. Twenty-four demonstrated some degree of parenchymal dilatation and disruption which were usually lobular in distribution. Panlobular emphysema of varying severity occurred in 20 patients and centrilobular in 4. The overall incidence of both forms of emphysema was 30% in the group of nonsmokers.

Auerbach et al. (1963, 1966) reported the results of their examination of lung tissue obtained from 1,582 autopsies performed from 1955 through 1960 at a Veterans Administration hospital in New Jersey. The results are expressed in terms of degree of rupture of alveolar septum, fibrosis and thickening of walls of small arteries and arterioles. There was no assessment of degree of panlobular or centrilobular emphysema, so that it is not possible to relate their observations with those of others. They concluded that these pathological changes were highly associated with cigarette smoking and all of them were found to increase with the age of the smoker. Although 120 of 1340 cases were nonsmokers, it is not possible to state how many of them can be regarded as showing lesions of pulmonary emphysema, similar to those reported by others.

The later report of Auerbach et al. (1972) conformed with the usual practice of specifying if the total pathology of the lung signified emphysema. Between 1963 and 1970, 2,613 men and women who died of the disease were studied at autopsy in seven cooperating hospitals in New Jersey and New York. Their results in males were as follows:

176 nonsmokers - 10% pulmonary emphysema	9% no	emphysema
181 smoked < 20 87% " " " " " cigarettes/day	13% "	11
658 smoked > 20 99.7% "	0.3%	
cigarettes/day	4.	

The statistics have been criticized by Panzer (1972).

The above results indicate that in unselected autopsy cases 10% of nonsmokers develop pulmonary emphysema. The 87 to 99% incidence of pulmonary emphysema among somkers is the highest ever reported.

A 50 to 75% incidence among smokers has been reported by Thurlbeck (1963) from Canada, by Petty et al. (1967) from Colorado, by Ishikawa (1969) et al./from Canada, by Yamanaka (1970) from Japan, by McLaughlin and Tueller (1971) from California, by Otto et al. (1970) in Germany, and by Fingerland et al. (1971) in Czechoslovakia. The statistics for the last-mentioned series are as follows:

145 male nonsmokers 26% pulmonary emphysema
405 female nonsmokers 30% " "
109 male ex-smokers 50% " "
408 male smokers 50% " "

The series reported from Czecheslovakia show a different incidence of pulmonary emphysema in autopsy of unselected patients from those reported by Auerbach et al. (1972) from the United States.

The incidence of emphysema among nonsmokers is 2 to 3 times higher in Czechoslovakia than in the United States, and the incidence of emphysema is half as much in Czechoslovakia.

The series of autopsies performed by Alli (1971) in Nigeria indicate the following:

59 males	25.4%	pulmonary emphys	ema
55 females	12.7%	n .	
ll4 total	19.3%		

There is a low incidence of smoking among the above cases. The overall incidence of 19% emphysema is midway between the incidence for nonsmokers reported from the United States and Czechoslovakia.

PETTY T L. RYAN S F and MITCHELL R S: Cigarette smoking and the lungs.

Arch Environ Health 14: 172-77, 1967.

1972.

#### II-B. ANTEMORTEM STUDIES

The extent of the occurrence of pulmonary emphysema among nonsmokers can be derived from a review of groups of patients with a diagnosis of pulmonary emphysema based on clinical studies. The reports are as follows:

Abbott et al. (1953)	250 males and females from Georgia:	41 nonsmokers
Mitchell (1962)  Mitchell et al. (1964)	123 males and females from Colorado:	8 nonsmokers
Kahana et al. (1963)	19 males from Canada:	1 nonsmoker
Seebohm and Bedell (1963)	10 males from Iowa:	0 nonsmoker
Antonio (1963)	16 males or females from Philippines:	4 nonsmokers
Boushy and Lewis (1964)	61 males and females from Michigan:	4 nonsmokers
Ray (1965)	360 males and females from Virginia:	8 nonsmokers
Burgess and Burgess (1966)	204 males and females from Rhode Island:	15 nonsmokers
Boushy et al. (1968)	49 males from Louisiana:	0 nonsmoker
Lépine and Myre (1969)	41 males or females from Canada:	0 nonsmoker
Saha and Jain (1970)	50 males and females from India: 1005	8 nonsmokers (051337

The above reports are based on series numbering 10 to 360 patients with clinical diagnosis of pulmonary emphysema. In the total number of \$133 selected patients, there are 89 nonsmokers. The incidence of 7.6% of nonsmokers among patients with pulmonary emphysema is lower than that based on postmortem examination (see Section II-A.)

Patients suffering from pulmonary emphysema have been investigated while inhaling cigarette smoke. The results have so far been conflicting. Bickerman and Barach (1954) compared the response of 91 patients suffering from chronic nontuberculous pulmonary disease, most of them having pulmonary emphysema, with that of 27 healthy subjects. The smoking of 3 cigarettes did not increase bronchospasm or impair ventilatory function. There was a reduction of vital capacity and of maximal breathing capacity in 10 patients who had bronchial asthma. Both measurements were increased in 10 patients with emphysema, in whom coughing provoked by smoking resulted in expectoration.

The observations of Eich et al. (1956, 1957) were made on 15

patients with pulmonary emphysema. After the smoking of one cigarette,

there was an increase in airway resistance in 14 of them and no significant

change in pulmonary compliance.

The above-cited observations do not necessarily imply that cigarette smoking leads to pulmonary emphysema. There are numerous data relating to the effects of cigarette smoking on lung function of healthy subjects that are discussed in Section VI.

## BIBLIOGRAPHY

# II. OCCURRENCE OF PULMONARY EMPHYSEMA IN NONSMOKERS B. ANTEMORTEM STUDIES

그는 물병이 되는 사람이 되는 것이 살아왔다면 가득하다면 가는 사람들이 되었다. 그는 사람이 다	Kep
ABBOTTO A, HOPKINS W A, VAN FLEIT and ROBINSON J S: A new approach to pulmonary emphysema. Thorax 8: 116-132, 1953.	43
ANTONIO R.N: Diffuse obstructive emphysema clinico-physiological manifestations among Filipinos. Bull Quezon Inst 7: 127-39, 1963.	,44
BICKERMAN H A and BARACH A L: The effect of cigarette smoking on ventilatory function in patients with bronchial asthma and obstructive pulmonary emphysema.  J Lab Clin Med 43: 455-62, 1954.	<b>4</b> 5
BOUSHY SF and LEWIS BM: The natural history of emphysema. Geriatrics 19: 910-19, 1964.	46
BOUSHY S.F., KOHEN R., BILLIG D.M. and HEIMAN M.J.: Bullous emphysema: clinical, roentgenologic and physiologic study of 49 patients. Dis, Chest 54:327-34, 1968.	. 47
BURGESS A M and BURGESS S B: Chronic obstructive pulmonary emphysema. A review, with special reference to its relation to cigarette smoking.  RIMed J 49: 461-5, 1966.	48
EICH R H, GILBERT R and AUCHINCLOSS J H Jr: Effects of smoking on respiratory mechanics in chronic pulmonary emphysema. Clin Res Proc 4: 151-52, 1956.	49
EICH R H, GILBERT R and AUCHINCLOSS J H Jr: The acute effects of smoking on the mechanics of respiration in chronic obstructive pulmonary emphysema.  Am Rev Tuberc 76: 22-32, 1957.	50
KAHANA L M, ARONOVITCH M and PLACE R: A comparative study of the clinical and functional pattern in emphysematous patients with and without chronic respiratory failure. Am Rev Resp Dis 87: 699-716, 1963.	51
LEPINE C and MYRE M: L'emphyseme pulmonaire. Confrontations cliniques et physiologiques avec une reference speciale a l'usage de la cigarette.  Union Med Can 98: 1853-58, 1969.  MITCHELL R S: Diffuse pulmonary emphysema and occupation. JAMA 181:	52
MITCHELL R S: Diffuse pulmonary emphysema and occupation. JAMA 181:	53
MITCHELL R S, VINCENT T N and FILLEY G F: Cigarette smoking chronic bronchitis, and emphysema. JAMA 188: 12-16, 1964.	54
RAY ES: Pulmonary emphysema. A disease of cigaret smokers. Virginia Med Monthly 92: 170-73, 1965.	55
SAHA N C and JAIN S K: Chronic obstructive lung disease in Delhi. A comparative study. Indian J Chest Dis 12: 40-51, 1970.	56
SEEBOHM P M and BEDELL G N: Primary pulmonary emphysema in young adults.  Am Rev Resp Dis 87: 41-46, 1963.	57

II-C. RISK FACTORS IN THE PATHOGENESIS OF PULMONARY EMPHYSEMA

That pulmonary emphysema may occur in nonsmokers can only be interpreted to mean that smoking is not the only important causative factor. There are five other factors that are as significant as cigarette smoking, namely: sex, age, genetic factors, infection, and air pollution. They can properly be described as risk factors which predispose a subject to the development of pulmonary emphysema. The first three are discussed in this section but the last two are dealt with in Section III because they contribute to the pathogenesis of chronic bronchitis as well as of pulmonary emphysema.

2. Sex and pulmonary emphysema. It has generally been accepted that pulmonary emphysema occurs predominantly in males.

The routine postmortem examination shows the following incidence of the disease for male in relation to female subjects: -- 2:1 as determined by Smith (1965), 3:2 by Azcuy et al. (1962), and 1:1 by Snider et al. (1962).

The emphysema registries show a higher ratio of 9:1 reported by Mitchell and Filley (1964), Burrows et al. (1965), and Webster et al. (1968).

There were no differences in laboratory data and family history between males and females.

The smoking habits of females are discussed in three additional reports. Bedell and Seebohm (1960) described a group of 12 women with pulmonary edema, ranging in age from 31 to 65 years, two of whom were nonsmokers. Lindall et al. (1967) evaluated pulmonary function in 101 females without lung disease. The women smokers were found to have lower mid-maximal flow rates and ratios of forced expiratory volume to

1005051340

Source: https://www.industrydocuments.ucsf.edu/dbcs/pimk0000

vital capacity, compared with nonsmokers. Carilli et al. (1973) reviewed chest films of 92 women and could note changes such as diffuse linear and nodular fibrosis in smokers that were not seen in nonsmokers.

These observations are similar to those reported by others in males and do not detract from the original premise that there is a higher incidence of pulmonary emphysema in males. Anderson et al. (1972) reviewed a series of necropsy cases and concluded that sex has an effect on the incidence of emphysema over and above the difference between the sexes in the amount of smoking.

- 2. Aging and pulmonary emphysema. The high incidence of pulmonary emphysema in the aged has been discussed in a monograph (Gross, 1964) and several review articles (Burrows, 1962; Cander and Blumenthal, 1963; Alvarez, 1964; Herberg, 1966; Niesenbaum, 1968). The mechanism by which aging leads to emphysema has not been identified. Degeneration and loss of elasticity occur, which are terms of a descriptive nature and do: not explain how the aging operates as a risk factor.
- 3. Genetic factors. Pulmonary emphysema appears in white males more frequently than in the Negro. The relative incidence varies, depending on the manner of comparison. Counting of clinical cases reveals that the incidence in white males is 3 or 4 times more frequent than in Negro males (Murphy et al., 1962). Deaths from emphysema in white males are 10 times more frequent (Ray, 1965). The lower incidence in the Negro does not relate to dissimilar smoking habits or a different

effect of smoking upon maximal terminal flow (Canter and Luchsinger, 1967, 1968). The Negro male has smaller vital capacity, which would not account for a lower incidence of pulmonary emphysema (Damon, 1966). The East Indians residing in Jamaica are reported to suffer more from the centrilobular type of pulmonary emphysema than the Chinese, Negro and white residents (Hayes and Summerell, 1969; Hayes, 1970).

A genetic defect has been identified in some patients with pulmonary emphysema. A deficiency in alpha-antitrypsin can be inherited and is believed to participate in the pathogenesis of pulmonary emphysema (see monograph by Eriksson, 1965, review article by Guenter et al., 1971 and bibliographic list no. 5). Patients have been reported to have pulmonary emphysema and alpha-antitrypsin deficiency who were nonsmokers: 2 out of 9 in Massachusetts (Talamo et al., 1968), 1 out of 8 in Washington (Bell, 1970), 1 out of 12 in England (Jones and Thomas, 1971), 1 out of 6 in Texas (Stevens et al., 1971), 3 out of 10 in Minnesota (Black et al., 1972). It has been proposed that cigarette smoking provokes the early onset of emphysema in subjects with this genetic defect (Levine et al., 1970; Mittman et al., 1971 a, b; Fallat: et al., 1971, 1973; Fallat, 1972; Hutchison et al., 1971, 1972). However, the evidence is not unequivocal and is based on the coexistence of emphysema and deficiency of alpha-antitrypsin among smokers. Observations have been made which suggested that smoking caused emphysema in deficient There is no association between prevalence of pulmonary

The identification of alpha-antitrypsin deficiency as a genetic defect in some patients with pulmonary emphysema raises the possibility of other defects which have not yet been identified (see review articles by Gedda, 1967, McKusick and Mutalik, 1968, and Larson et al., 1970).

Patients with emphysema have been characterized by personality features (De Cencio et al., 1968), somatotype (Damon, 1961; Damon et al., 1962), baldness (Buechner et al., 1964), and longevity (Deutscher and Higgins, 1970). There are personality features which are inherited by the and Cochrane individual (Heath, 1958; De Cencio et al., 1968; Clark/1970; Ochsner and Damrau, 1970).

#### BIBLIOGRAPHY

1	I. OCCURRENCE	OF PULMONAR	Y EMPHYSEMA IN NO	NSMOKERS
	<ul> <li>The second of the second of the</li></ul>		•	
c.	RISK FACTORS IN	THE PATHOGEN	ESIS OF PULMONAR	Y EMPHYSEMA

Is emphysema a geriatric o		Geriatries	19: 139-42,	58
	•			

ANDERSON JA,	DUNNILL M S and	d RYDER R C:	Dependence	of the incidence
of emphysei	ma on smoking hist	ory, age, and se	x. Thorax	27: 547-51,
1972.	المنبوب المراوية أخراوية		· · · <del></del>	

										-	
DEDELL	C Nt	CEERAIN		-		_		• •			
あたれた アア	G N and	2FFBOH	4 PM:	Primary	pulmonary	emphysema	in	Women			
					F	p,		0111 C111			
	nwa Med	Sac 56.	258-62	1066	- 10 to 10 t					7.4	
	O 4 111.C. G	JUC 30.	230-02,	1700.				and the second second	•	4 4 4 4	

		****					
BEL	LL R: The r	adiographic r	nanifestation	ns of alpha	a-l antitr	ypsin defici	ency. An
		recognizable					
	(COPD).	Radiology	95: 19-24.	1970.		-	5.00

BLAC	KLF,	HYATT	R E and	STUBBS	SE: M	echanis	m of expi	ratory a	irflow
	limitati	on in chr	onic obs	tructive p	ulmonary.	disease	associal	ed with	3.5
				Am Rev					

			• • • • • • • • • • • • • • • • • • • •	•	
BUECHNER HA	BROWN	M and TRETTO	TA R T.	Baldrage	-d:
	, 210	and Indic	126 16 3.	Daidness a	mu empnysema.
J Louisiar	a Med Soc	116: 329-32,	1964.		1995

		Marie Artist Control of the		
BURROWS B:	Pulmonary	and respirator	ry diseases in the aged.	Postgrad Med
31: 296-3	02, 1962.			

CANDER	L and BLUMENTHA	L WS:	Emphysema and the aging lung.	Geriatrics
18:	413-420, 1963.			6

$\textbf{CANTER} \ \ \textbf{H} \ \ \textbf{G}_{:} \ \ \textbf{and} \ \ \ \textbf{LUCHSINGER}$	PC:	Interrelat	ionshi	ps of	smoking	habits,	race
and maximal terminal flow.	Am	Rev Resp	Dis	96:	147-48,	1967.	

CANTER E	G and	LUCHSINGER	PC:	Interrelationship	s of	smoking !	habits,	race
and n	naximal	terminal flow.	Am			107.1-75,		

CARILLI A D, KOTZEN	L M and FISCHER	MiJ: The chest:	roentgenogram in
smoking females.	Am Rev Resp Dis	107: 133-36, 197	3.

		•
CLARK TJH and COCHRANE	GM: Effect of personality on	alveolar
ventilation in patients with	chronic airways obstruction.	Br Med J
<b>273-75</b> , 1970		

GEDDA L:	Il chronon,	ossia la dimensione	tempo	dell'unita	ereditaria.
Gazz S	anit 38: 3	23-26, 1967,	•		

DAMON A:	Constitution and smoking.	Science	134.	330-41	1061

DAMON A:	Constitution and	smoking in	Italian-	American factory workers.
Am.J	Phys Anthropol	20: 67-68	1962	•

			•		
DAMON A:	Negro-white differences in	pulmonary fu	nction (V	ital capacity.	timed
vital c	apacity, and expiratory flo	vrate). Hum	an Biol	38: 380-93.	1966.

DeCENCIO D	V, LESHNER	Mand :	LESHNER B:	Personality	characteristics of
<ul> <li>patients</li> </ul>	with chronic	obstructi	ve pulmonary	emphysema.	Arch Phys Med
Rehabil	49: 471-75,	1968.	•		

.

DEUTSCHER S and HIGGINS M W: The relationship of parental longevity to ventilatory function and prevalence of chronic nonspecific respiratory disease among sons. Am Rev Resp Dis 102: 180-89, 1970.	75
ERIKSSON S: Studies in alpha-antitrypsin deficiency. Acta Medica Scandinavica 177 (Supplementum 432): 1 -85, 1965.	76
FALLAT R J: To the Editor. Ann Intern Med 72: 438-39, 1970.	77
FALLAT R.J. POWELL M.R. KUEPPERS F and LILKER E: 133 Xe ventilatory studies in 1-antitryps in deficiency. J. Nucl. Med 14: 5-13, 1973.	78
FALLAT R, POWELL M, KUEPPERS F, NADEL J and MURRAY J: Chronic obstructive pulmonary disease with intermediate alpha 1-antitrypsin deficiency.  Chest 59: (Suppl 5) 205-225, 1971.	79
GEISLER L.S., BACHMANN G.W., LAUMEN F., NOLTE D., WENTZEL H and ROST H-D: a -antitrypsin und immunglobuline bei chronisch unspezifischen lungenerkrankuggen und lungentuberkerkulose. Dtsch Med Wochenschr 97: 329-35, 1972.  GROSS P: Environmental strees and the aging lung. The pulmonary effects of air pollution. In: Aging of the lung. Perspectives. The tenth Hahnemann Symposium. Edited by Leon Cander. Grune & Stratton, New York.	80
pp. 217-26, 1964.	81
GUENTER CA, WELCH MH and HAMMARSTEN JF: Alpha anititrypsin deficiency and pulmonary emphysema. Ann Rev Med 22: 283-92, 1971.	82
HAYES JA and SUMMERELL JM: Emphysema in a non-industrialized tropical island. Thorax 24: 623-25, 1969.	83
HAYES J A: Racial, occupational, and environmental factors in relation to emphysema in Jamaica. Chest 57: 136-40, 1970.	84
HEATH C W: Differences between smokers and nonsmokers. Arch Intern Med 101: 377-88, 1958.	85
HERBERG D: Seniles und obstruktives emphysem. Diagnose, atiologic und begutachtung. Beitr Klin Tuberk 133: 75-91, 1966.	86
HUTCHITISON D CS, BARTER CE, COOK PJL, LAWS JW, MARTELLI NA and HUGH-JONES P: Severe pulmonary emphysema. A comparison of patients with and without a <sub>1</sub> -antitrypsin deficiency. QJ Med 41: 301-15, 1972.	87
HUTCHISON D C S, COOK P J L, BARTER C E, HARRIS H and HUGH-JONES P:  Pulmonary emphysema and a antitrypsin deficiency.  1: 689-94, 1971.	88
JONES M C and THOMAS G O: Alpha, antitrypsin deficiency and pulmonary emphysema  Thorax 26: 652-62, 1971.	89
LARSON R K, BARMAN M L, KUEPPERS F and FUDENBERG HH: Genetic and environmental determinants of chronic obstructive pulmonary disease.  Ann Intern Med 72: 627-32, 1970.	90
LEVINE BW, TALAMO RC, SHANNON DC and KAZEMI H: Alteration in distribution of pulmonary blood flow. An early manifestation of Alpha antitrypsin deficiency. Ann Intern Med 73: 397-401, 1970.	91
LINDALL A, MEDINA A, GREIMER J T: A reevaluation of normal pulmonary function measurements in the adult female. Am Rev Resp Dis 95: 1061-	62

Bibliography II-C page	30 Reprin
McKUSICK V A and MUTALIK G S: Genetics and pulmonary disease. In: The Lung. Edited by A A Licbow and D E Smith. The Williams and Wilkins Company, Baltimore, 1968. PP. 187-202.	93
MITTMAN C, LIEBERMAN J, MARASSO F and MIRANDA A: Smoking and chronic obstructive lung disease in alpha 1-antitrypsin deficiency.  Chest 60: 214-221. 1971A.	94
MITTMAN C, LIEBERMAN J, MARASSO F and MIRANDA A: Pulmonary alveolar proteinosis and blood dyscrasias. Clin Res 19: 146, 1971B.	<b>95</b>
MURPHY R E, KATZ S, MASSARO D J and LUCHSINGER P C: Is emphysema a disease predominantly of the while male? JAMA 181: 726-27: 1962.	96
NIESENBAUM L: Problems of pulmonary disease in the aged. Geriatrics 23: 127-31, 1968.  OCHSNER A and DAMRAU F: Control of cigarette habit by psychological aversive conditioning: Clinical evaluation in 53 smokers. J Am Geriatr Soc 18:	97 98
RAY E.S: Relative incidence of pulmonary emphysema among negroes and whites of both sexes. Med Coll Virginia Quart 1: 50-51, 1965.	99
SANO T, YODOYAMA M and RHOADS G G: Clinical significance of a -antitrypsis level a preliminary evaluation among Japanese men. Hawaii Med J 31: 27-32, 1972.	n 100
STEVENS P.M., HNILICA V.S., JOHNSON P.C and BELL R.L: Pathophysiology of hereditary emphysema. Ann Intern Med 74: 672-80, 1971.	101
TALAMO R C, ALLEN J D, KAHAN M G and AUSTEN F: Hereditary alpha antitrypsin deficiency. N Engl J Med, 278: 345-51, 1968.	102
WEBSTER JR Jr, KETTEL LJ, MORAN F, BUGAIESKI S and CUGELL D W:  Chronic obstructive pulmonary disease. A comparison between men and  women. Am Rev Resp Dis 98: 1021-26, 1968.	103
AZCUY A D. ANDERSON A E. and FORAKER A E: The morphological spectrum of aging and emphysematous lungs. Ann Intern Med. 51:1, 1962.	103a
BURROWS B, NIDEN A H, BARCLAY W R and KASIK J H: Chronic obstructive lung disease: I. Clinical and physiologic findings in 175 patients and their relationship to age and sex. Amer Rev Resp Dis 91:521, 1965.	103ъ
MITCHELL R S and FILLEY G F: Chronic obstructive bronchopulmonary disease:  I. Clinical features. Amer Rev Resp Dis 89:360, 1964.	103c
SMITH V K: Survey of types and severity of emphysema in routine autopsies. Aust Ann Med 14:28, 1965.	103a
SNIDER G L, BRODY J S and DOCTOR L: Subclinical pulmonary emphysema. Amer Rev Resp Dis 85:666, 1962.	103e

#### II-D. EXPERIMENTAL PULMONARY EMPHYSEMA

The publicity surrounding the experimental induction of pulmonary emphysema by Auerbach and his collaborators has given the impression that the problem has been settled. Numerous attempts to study the matter have been made by others and the results have been negative, which have not been publicized. There is still considerable doubt that cigarette smoking can lead to pulmonary emphysema in experimental animals. The problems in induction of emphysema have been discussed by Strawbridge (1960a), Thurlbeck (1%3), and Boren (1965). Reif et al (1970) and Mittman (1972).

1. Experiments in the dog. Two groups of dogs have been exposed to cigarette smoke through a tracheostomy tube by Auerbach and his collaborators: ten dogs for 420 days (Auerbach, 1967, 1969; Auerbach et al., 1967, 1968, 1970; Cahan and Kirman, 1968) and 97 dogs up to 876 days (Hammond et al., 1970; Frasca et al., 1971). The results are entirely histopathological in nature, which the investigators have interpreted to indicate pulmonary emphysema. Their interpretation has been criticized by other histopathologists and nothing can be added to this controversy (see Section III). One criticism that has been entirely ignored is the lack of measurements of lung function, which other investigators have developed for application to the dog and other animal species. 1005051347

Anderson et al. (1964) and Hernandez et al. (1966) exposed greyhounds to hyperinflation of the lung or cigarette smoke. Lungs from

disease. The lungs from animals exposed to cigarette smoke showed lesions that were inflammatory in nature. Although there was parenchymal disruption, the pattern of smoke exposure employed with the dog is so different from the habit in man. that the authors could not readily transfer their conclusion to apply to the human form of emphysema. Rockey and Speer (1966) and Syzganov et al. (1971) made similar observations and conclusions in the dog. The results of applying a smoking machine in dogs, which simulate those of human inhalation, have not been reported (Bair et al., 1969). Marco et al. (1971) and Mass et al. (1972) induced experimental emphysema in dogs, using leukocyte homogenate. Their findings are entirely histological in nature.

2. Experiments in the rat. Both histological and physiological techniques have been applied to the rat lung to demonstrate conclusively that pulmonary emphysema has developed. The model consists of a combination of tracheal ligation and intratracheal injection of papain. After 4 to 6 weeks, there is an increase of functional residual capacity and the lung shows histological signs of emphysema (Palecek et al., 1967). Concurrent administration of progestational agents retards the development of emphysema (Ito and Aviado, 1968a; Aviado and McKinney, 1969; Aviado and Carrillo, 1969; Shore and Aviado, 1969; Inoh and Aviado, 1971). The concurrent administration of cigarette smoke does not influence the emphysematous

process (Ito and Aviado, 1968b; Aviado et al., 1970). Inhalation of cigarette smoke daily for 8 weeks does not increase functional residual capacity. In these rat's exposed to cigarette smoke, there is an increase in pulmonary resistance, which is discussed further in another section relating to the airways.

The inhalation of nitrogen dioxide in the rat causes the development Haydon et al, 1965; of pulmonary emphysema (Freeman and Haydon, 1964; / 🦈 Freeman, 1968; Freeman et al., 1968 a, b, c; 1969). The concentration necessary to induce the lesions is approximately 1,000 times the amount of nitrogen dioxide present in cigarette smoke (Haagen-Smit et al., 1959; Bokhoven and Niessen, 1961). The phenomenon is of toxicological interest when nitrogen dioxide is present as a pollutant or a chemical fume that is inhaled (von Oettingen, 1941; Lowry and Schuman, 1956; Eecklake et al., 1957; Gray, 1959). In this context, nitrogen dioxide has the following effects in rats and other animal species: reduced resistance to bacterial pneumonia (Ehrlich and Henry, 1968; Henry et al., 1969, 1970), alteration in collagen fibrils and basement membranes of the lung (Stephens et al. 1971); increase in metabolic activity of lung tissue (Buckley and Loosli, 1969); increase in nonelastic resistance of the lung (Haydon et al., 1967); and increase in lung tissue antibody in the blood (Balchum et al., 1965; Boren et al., 1965). Gross et al. (1968 a, b) did not observe any acceleration by nitrogen oxide of experimental silicosis caused by coal or blast furnace stack dust in rats.

Cadmium is another constituent of cigarette smoke (Nandi et al., 1969). Exposure to cadmium fumes causes lesions in the human lung similar to those of emphysema (Bonnell, 1955; Bonnell et al., 1959; Smith et al., 1960). However, the phenomenon has not been examined in rats or other animal species to determine whether the amount of cadmium in cigarette smoke is sufficient to produce pulmonary lesions.

3. Experiments in other animal species. Experimental emphysema in several animal species has been reported but not the influence of cigarette smoke. The following species have been used: hamster (Goldring et al., 1968; Gross and de Treville, 1969), rabbit (Strawbridge, 1960 B and C; Boatman and Martin, 1965), piglet (Glauser and Glauser, 1969); horse (Gillespie et al., 1964, 1966; Gillespie and Tyler, 1967; Eberly et al., 1966; McLaughlin et al., 1965), and cow (Carlson et al., 1968; Carlson et al., 1972).

## D. EXPERIMENTAL PULMONARY EMPHYSEMA

	Kepin
ANDERSON A E.Jr., AZCUY A, BATCHELDER T L and FORAKER A G: Experim analysis in dogs of the relationship between pulmonary emphysema, alveolitis, and hyperinflation. Thorax 19: 420-32, 1964.	ental
AUERBACH O: Pathological findings in lungs of dogs following exposure to cigarette smoke. In: Transactions of the 26th VA armed forces pulmonary disease research conference. Prepared and edited by VA department of medicine and surgery, 1967, p. 33.	105
AUERBACH O: Emphysema: Human and experimental. In: Conference on inhalation carcinogenesis. Sponsored by National Cancer Institute and U.S. Atomic Energy Commission, 1969, p. 20.  AUERBACH O, HAMMOND E C, KIRMAN D and GARFINKEL L: Emphysema produced in dogs by cigarette smoking. JAMA 149: 241-46, 1967.	106 107
AUERBACH O, HAMMOND E C, KIRMAN D and GARFINKEL L: Emphysema produced in dogs by cigarette smoking. In: Inhalation carcinogenesis Edited by M G Hanna Jr, P Nettersheim, and J R Gilbert. U.S. Atomic Energy Commission (Division of technical information) Information Interscience Incorporated, 1970, pp. 375-87.	108
AUERBACH O, HAMMOND E C, KIRMAN D, GARFINKEL L and STOUT A P:  The effect of direct cigarette smoke inhalation on the respiratory tree of dogs. Nat Cancer Inst Monogr 28: 65-77, 1968.	, 109
AVIADO D M and CARRILLO L R: Hormones and pulmonary effects of tobacco.  L. Corticosteroids and their antiasthmatic action. Arch Environ Health 18: 925-933, 1969.	110
AVIADO D M and McKINNEY G R: Oral progestagens and experimental pulmonary emphysema. Pharmacol Res Comm 1: 283-87, 1969.	111
AVIADO D M, SADAVONGVIVAD C and CARRILLO L R: Cigarette smoke and pulmonary emphysema. Influence of bronchodilators and biogenic amines in experimental induction in rats. Arch Environ Health 20: 483-87, 1970.	112
BAIR W J, PORTER N S, BROWN D P and WEHNER A P: Apparatus for direct inhalation of cigarette smoke by dogs. J Appl Physiol 26: 847-50, 1969.	113
BALCHUM O J. BUCKLEY R D, SHERWIN R and GARDNER M: Nitrogen dioxide inhalation and lung antibodies. Arch Environ Health 10: 274-77, 1965.	114
BECKLAKE M.R., GOLDMAN H.I. BOSMAN A.R. and FREED C.C.: The long-term effects of exposure to nitrous fumes. Am Rev Tuberc 76: 398-409; 1957.	115
BLAIR W H, HENRY M C and EHRLICH R: Chronic toxicity of nitrogen dioxide.  II. Effect on histopathology of lung tissue. Arch Environ Health. 18:  186-92, 1969.	116
BOATMAN E S and MARTIN H B: Electron microscopy in pulmonary emphysema or rabbits. Am Rev Resp Dis 91: 197-205, 1965.	of 117
BOKHOVEN C and NIESSEN H J: Amounts of oxides of nitrogen and carbon monoxi in cigarette smoke, with and without inhalation. Nature 192: 458-59,	de

BONNELL JA: Emphysema and proteinuria in man casting copper-cadmium alloy

pa g	e 36
Bibliography II-D	Reprint
BONNELL JA, KAZANTZIS G and KING E: A follow-up study of men exposed to cadmium oxide fume. Br J Ind Med 16: 135-47, 1959.	120
BOREN HG: Environmental emphysema. Am Rev Resp Dis 92: 1-15, 1965.	121
BOREN H.G. DELAHANT A B and STEENKEN W: Reactions of guinea pig lung to injected lung homogenates. Medicina Thoracalis 22: 355-64, 1965.	122
BUCKLEY R D and LOOSLI C G: Effects of nitrogen dioxide inhalation on germfree mouse lung. Arch Environ Health 18: 588-95, 1969.	123
CAHAN W G and KIRMAN D: An effective system and procedure for cigarette smoking by dogs. J Surg Res 8: 567-75, 1968.	ng 124
CARLSON JR, DYER I A and JOHNSON RJ: Tryptophan-induced interstital pulmonary emphysema in cattle. Am J Vet Res 29: 1983-89, 1968.	125
CARLSON J.R., YOKOYAMA M.T. and DICKINSON E.O: Induction of pulmonary edema and emphysema in cattle and goats with 3-methylindole. Science 176: 298-99, 1972.	126
EBERLY V E, TYLER W S and GILLESPIE J R: Cardiovascular parameters in emphysematous and control horses. J Appl Physiol 21: 883-89, 1966.	127
EHRLICH R and HENRY M C: Chronic toxicity of nirtogen dioxide. Arch Environ Health 17: 860-65, 1968.	128
FRASCA J.M., AUERBACH O, PARKS VR and JAMIESON J.D.: Electron microscop observations on pulmonary fibrosis and emphysema in smoking dogs.  Exp Mol Pathol 15: 108-25, 1971.	ic 129
FREEMAN G: The sub-acute NO <sub>2</sub> lesion of the rat lung. In: 9th American medical association air pollution medical research conference, 1968, pp 1-5.	130
FREEMAN G, CRANE S C, STEPHENS R J and FURIOSI N J: Pathogenesis of the nitrogen dioxid-induced lesion in the rat lung: A review and presentation of new observations. Am Rev Resp Dis 98: 429-43, 1968	131
FREEMAN G, CRANE S C, STEPHENS R J and FURIOSI N J: Environmental factors in emphysema and a model system with NO <sub>2</sub> . Yale J Biol Med 40: 566-75, 1968B.	132
FREEMAN G, CRANE S C, STEPHENS R J and FURIOSI N J: The subacute nitrogen dioxide-induced lesion of the rat lung. Arch Environ Health 18: 609-12, 1969.	133
FREEMAN G and HAYDON G B: Emphysema after low-level exposure to NO <sub>2</sub> .  Arch Environ Health 8: 125-28, 1964.	134
FREEMAN G, STEPHENS R J, CRANE S C and FURIOSI N J: Lesion of the lung in rats continuously exposed to two parts per million of nitrogen dioxide. Arch Environ Health 17: 181-92, 1968	135
GILLISPIE J R and TYLER W S: Capillary and cellular changes in alveolar walls of emphysematous horse lungs. Am Rev Resp Dis 95: 484-90, 1967.	136
GILLISPIE JR, TYLER WS and EBERLY VE: Blood pH, O <sub>2</sub> , and CO <sub>2</sub> tensions in exercised control and emphysematous horses. Am J Physiol 207:	137

Bibliography II-D	page 37	Reprin
GILLISPIE J.R. TYLER WS and EBERLEY VE: Pulmonary ventilation resistance in emphysematous and control horses. J. Appl. Physiol	and	
21: 416-22, 1966.		138
GLAUSER E M and GLAUSER S C: The role of acidosis in the etiology pulmonary emphysema. Proc Soc Exp Biol Med 131: 1034-37, 196		139
GOLDRING I P, GREENBURG L and RATNER I M: On the production of emphysema in Syrian hansters by aerosol inhalation of Papain.  Arch Environ Health 16: 59-60, 1968.		140
GRAY E L: Oxides of nitrogen: Their occurrence, toxicity hazard. Arch Ind Health 19: 479-86, 1959.		141
GROSS P and deTREVILLE T P: Emphysema and pneumoconiosis.  Arch Environ Health 18: 340-49, 1969.		142
GROSS P, deTREVILLE T P, BABYAK M A, KASCHAK M and TOLKER Experimental emphysema. Arch Environ Health 16: 51-58, 1968A.	EB:	143
GROSS P, deTREVILLE T P, BABYAK M A, KASCHAK M and TOLKER Experimental emphysema: effect of chronic nitrogen dioxide exposure papain on normal and pneumoconiotic lungs. Pub Health Serv 1787: 357-78, 1968B.	and of	144
HAAGEN-SMIT L J, BRUNELLE M F and HARA J: Nitrogen oxide con smokes from different types of tobacco. Arch Ind Health 20: 399-4 1959.	00,	145
HAMMOND E C, AUERBACH O, KIRMAN D and GARFINKEL L: Effect cigarette smoking on dogs. Arch Environ Health 21: 740-53, 1970.		146
HAYDON G B, DAVIDSON J T, LILLINGTON G A and WASSERMAN K:  dioxide-induced emphysema in rabbits. Am Rev Resp Dis 95: 797-8 1967.		147
HAYDON G B, FREEMAN G and FURIOSI N J: Covert pathogenesis of N emphysema in the rat. Arch Environ Health 11: 776-83, 1965.	O2 induced	148
HENRY M.C., EHRLICH R and BLAIR W.H: Effect of nitrogen dioxide on of squirrel monkeys to klebsiella pneumoniae infection. Arch Environments 18: 580-87, 1969.		149
HENRY MC, FINDLAY J, SPANGLER J and EHRLICH R: Chronic tox of NO <sub>2</sub> in squirrel monkeys. Arch Environ Health 20: 566-70, 19		150
HERNANDEZ JA, ANDERSON A E Jr, HOLMES W L and FORAKER A C Pulmonary parenchymal defects in dogs following prolonged cigarette exposure. Am Rev Resp Dis 93: 78-83, 1966.		151
INOH T and AVIADO D M: Cardiopulmonary effects of progestational age emphysematous rats. Chest 59: 659-66, 1971.	ents in	152
ITO H and AVIADO D M: Pulmonary emphysema and cigarette smoke. A Environ Health 16: 865-70, 1968.	<u>Arch</u>	153
ITO H and AVIADO D M: Prevention of pulmonary emphysema in rats by		154

page 38	
Bibliography II-D	Reprint
LOWRY T and SCHUMAN L M: "Silo-filler's disease" a syndrome caused by nitrogen dioxide. J Am Med Ass 162: 153-60, 1956.	155
MARCO V, MASS B, MERANZE DR, WEINBAUM G and KIMBEL P: Induction of experimental emphysema in dogs using leukocyte homogenates.  Am Rev Resp Dis 104: 595-98, 1971.	156
MASS B, IKEDA T, MERANZE DR, WEINBAUM G and KEMBEL P: Induction of experimental emphysema. Cellular and species specificity.  Am Rev Resp Dis 106: 384-91, 1972.	157
McLAUGHLIN R F, TYLER W S, EDWARDS D W, CRENSHAW G L, CANADA R O, FOWLER M A, PARKER E A and REIFENSTEIN G H: Chlorpromazine-induced emphysema. Am Rev Resp Dis 92: 597-608, 1965.	158
MITTMAN C (editor): Pulmonary emphysema and Proteolysis. Academic Press, New York, 1972, pp 1-562.	159
NANDI M, HERSHEL J, SLONE D, SHAPIRO S and LEWIS G P: Cadmium content of cigarettes. Lancet 2: 1329-30, 1969.	160
PALECEK F, PALECEKOVA M and AVIADO D M: Emphysema in immature rats.  Arch Environ Health 15: 332-42, 1967.	161
REIF JS, RHODES W H and COHEN D: Canine pulmonary disease and the urban environment. Arch Environ Health 20: 676-83, 1970.	162
ROCKEY E E and SPEER F D: The ill effects of cigarette smoking on dogs.  Int Surg 46: 520-30, 1966.	163
SHORE S.R., and AVIADO D.M.: Hormones and pulmonary effects of tobacco.  Arch Environ Health 19: 59-69, 1969.	164
SMITH J P, SMITH J C and McCALL A J: Chronic poisoning from cadmium fume.  J Pathol 80: 287-96, 1960.	165
STEPHENS R J, FREEMAN G and EVANS M J: Ultrastructural changes in connective tissue in lungs of rats exposed to NO <sub>2</sub> . Arch Intern Med 127: 873-83, 1971.	166
STRAWBRIDGE H T G: Chronic pulmonary emphysema (an experimental study). I Histor	ical reviev
Am J Pathiol 37: 161-74, 1960 A.	167
STRAWBRIDGE H T G: Chronic pulmonary emphysema (an experimental study).  II. Spontaneous pulmonary emphysema in rabbits. Am J Pathiol  37: 309-31, 1960B.	168
STRAWBRIDGE H T G: Chronic pulmonary emphysema (an experimental study)  III Experimental pulmonary emphysema. Am J Pathiol 37: 391-411, 1960C.	169
SYZGANOV A N, GOLOVIN Y A and TROITSKAYA E G: On the influence of tobacco smoke on the organs of respiration. Eksperimentalnay Khirurgiya 16: 23-26, 1971.	170
THURLBECK W M: Pulmonary emphysema. Am J Med Sci 246: 332-53, 1963.	171

page 39

Reprint

VON OETTINGEN W F: The toxicity and potential dangers of nitrous fumes.

Pub Health Bull 272: 1-34, 1941.

Bibliography II-D

172

## CHRONIC BRONCHITIS: CONCURRENCE OF CIGARETTE SMOKING INHALATION OF POLLUTANTS AND INFECTION

Chronic bronchitis is a clinical disorder characterized by excessive mucus secretion. It may appear concurrently with pulmonary emphysema, so that the major problem in the diagnosis is to establish that bronchitis is the predominant disease.

As in pulmonary emphysema, cigarette smoking is one of the risk factors in the pathogenesis of chronic bronchitis. Two other risk factors, air pollution and infections, are discussed in this section. At the present time, it is not possible to rank these risk factors in the order of their importance.

The primary concern of this section is chronic bronchitis. literature on postmortem, clinical and animal investigations relating to chronic bronchitis is reviewed here. Three bibliographic lists are appended:

- No. 6. Chronic bronchitis: general aspects.
- No. 7. Bronchial carcinoma.
- No. 8. Air pollutants and lung disease.

The early cellular changes in the bronchial mucosa and secretion in a chronic bronchitis have been confused with metaplasia, so that it is important to exclude consideration of neoplastic disease from the following paragraphs.

#### III-A. POSTMORTEM EXAMINATION

The basic histopathologic change of chronic bronchitis is generalized hypertrophy and hyperplasia of the mucus-secreting bronchial glands throughout the treachea and bronchi. There are also diffuse inflammatory changes consisting principally of increased number of mononuclear cells in the thickened trancheobronchial submucosa and metaplastic epithelial changes with loss of cilia (Reid 1954, 1960; 1, 4, 4, 4, 5; Marcatili et al., 1967; Thurlbeck and Angus 1967).

There are only two reports of postmortem examination and known smoking habits of patients with chronic bronchitis. Mitchell et al (1966) reported 64 autopsies, all but one subject having been smokers and 91% for more than 20 pack-years (no of pack daily x years). Niewoehner et al. (1972) reported 4 fatal cases of regional chronic bronchitis, all of the patients having been chronic smokers.

All the other postmortem observations compare the histological features of the bronchi of smokers and nonsmokers. most of them healthy, Chang (1956, 1957, 1958) in 105 random autopsies noted distended goblet cells in 25 out of 71 smokers (35%) but only in 6 out of 34 nonsmokers (17.6%). The average length of cilia was shorter than in nonsmokers and the smokers' epithelium was thicker than that of nonsmokers. The hypertrophy of the bronchial mucus glands, characteristic of chronic bronchitis, has been reported to be more frequent among smokers. The gland/wall ratio was significantly raised

in 35 heavy smokers, but there was no difference between 23
nonsmokers and 15 moderate (13ss than 20 cigarettes/day) smokers
(Thurlbeck et al., 1963). A similar conclusion was arrived at in
a larger group of cases, also in Canada like the previous group
consisting of 40 nonsmokers and 111 smokers, including 58 heavy
smokers (Thurlbeck and Angus, 1964). In another group, reported
from England, the mean percentage volume of bronchial mucus glands was
significantly greater in 106 smokers than in 73 nonsmokers (Dunnill
and Ryder 1971; Ryder et al., 1971). In a smaller group, from Ohio,
there was no difference in percentages of such glands between 11
smokers and 13 nonsmokers (Sobonya and Kleinerman, 1972). Ido
et al. (1959) reported that percentages of goblet cells increased in
light smokers but decreased in heavy smokers.

Squamous metaplasia of bronchial epithelium is commonly seen in various pathologic processes as a result of atypical epithelial regeneration. In 128 unselected autopsies, the general incidence of metaplasia was 32%; the highest incidence was 71% in patients with primary lung carcinoma and the next highest was 41% in patients with chronic inflammatory disease of the lung. In the same group, 39% of nonsmokers had metaplasia and 45% of smokers had a similar epithelial change (Weller, 1955). The subsequent reports show the incidence of metaplasia to be as follows:

 Sanderad, 1958
 8% of 58 smokers
 54% of 39 nonsmokers

 Cross et al., 1961
 16% of 197 smokers
 8% of 33 nonsmokers

 Spain, 1969, 1970
 65% of 52 smokers
 50% of 36 nonsmokers

Source: https://www.industrydocuments.ucsf.edu/docs/pfmk000

Auerbach et al. (1961; 1962a, b; 1968) have reported their results in terms of nature of the nucleus relevant to neoplastic changes.

Their observations did not include those relating to bronchitis.

The incidence of goblet cell metaplasia is about equal in smokers and nonsmokers (Karpick et al., 1970). The proliferation of epithelial cells has been reported in the palate (Chapman and Redish, 1960) and larynx (Ryan et al., 1955) of smokers.

#### A. POSTMORTEM EXAMINATION

	Reprint
AUERBACH O, STOUT A P, HAMMOND E C and GARFINKEL L: Changes in bronchial epithelium in relation to cigarette smoking and in relation to lung cancer. N Engl J Med 265: 253-67, 1961.	173
AUERBACH O.STOUT A P. HAMMOND E C and GARFINKEL L: Changes in bronchial epithelium in relation to sex, age, residence, smoking and pneumonia. N Engl J Med 267: 111-19, 1962A.	174
AUERBACH O, STOUT A P, HAMMOND E C and GARFINKEL L: Bronchial epithelium in former smokers. N Engl J Med 267: 119-25, 1962B.	175
AUERBACH O, STOUT A P, HAMMOND E C and GARFINKEL L: Interrelationships among various histologic changes in bronchial tubes and in lung parenchyma.  Am Rev Resp Dis 90: 867-876, 1968.  CHANG S C: Studies of bronchial epithelium from smokers and nonsmokers.	. 176
Proc Am Ass Cancer Res 2: 99-100, 1956.  CHANG S C: Microscopic properties of whole mounts and sections of human bronchial	177
epithelium of smokers and nonsmokers. Cancer 10: 1246-62, 1957.  CHANG S C: Studies of suberepithelial tissue of bronchi from smokers and non- smokers. Proc Am Ass Cancer Res 2: 286-87, 1958.	178 179
CHAPMAN I and REDISH C H: Tobacco-induced epithelial proliferation in human subject. Arch Path 70: 133-40, 1960.	180
CROSS KR, WALZ DV, PALMER GK and WARNER ED: A study of the tracheobronchial epithelium and changes related to smoking.  J Iowa Med Soc 51: 137-40, 1961.	181
DUNNILL S and RUDEN R: A Quantative study of the relationship between chronic bronchitis emphysema and smoking. Chest 59 (5 Suppl): 355-65;1971.	182
IDE G, SUNTZEEF V and COWDRY E V: A comparison of the histopathology of tracheal and bronchial epithelium of smokers and nonsmokers. Cancer 12: 473-84, 1959.	183
KARPICK R J, PRATT P C, ASMUNDSSON T and KILBURN K H: Pathological findings in respiratory failure. Goblet cell metaplasia, alveolar damage, and myocardial infarction. Ann Intern Med 72: 189-97, 1970.	184
MARCATILI S, BONO M D and ZUCCHETTO C: Comportamento dell'epitelio bronchiale in alcune bronco-pneumopatie chroniche. Riv Anat Patol Oncol 31: 308-30, 1967.	185
MITCHELL RS, RYAN SF, PETTY TL and FILLEY G: The significance of morphologic chronic hyperplastic bronchitis. Am Rev Resp Dis 93: 720-29, 1966.	186
NIEWOEHNER DE, KLEINERMAN J and KNOKE JD: Regional chronic bronchitis.  Am Rev Resp Dis 105: 586-93, 1972.	187

REID L M: Pathology of chronic bronchitis. Lancet 1: 275-78, 1954.

REID L: Measurement of the bronchial mucous gland layer: a diagnostic yardstick in chronic bronchitis. Thorax 15: 132-41, 1960.	189
RYAN R F, McDONALD J R and DEVINE K D: The pathologic effects of smoking on the larynx. Arch Path 60: 472-80, 1955.	190
RYDER R C, DUNNILL M S and ANDERSON J A: A quantitative study of bronchial mucous gland volume, emphysema and smoking in a necropsy population.  J Pathol 104: 59-71, 1971.	191
SANDERUD K: Squamous metaplasia of the respiratory tract epithelium. An autopsy study of 214 cases. Acta Path Microbiol Scand 43: 47-61, 1958.	192
SOBONYA R E and KLEINERMAN J: Morphometric studies of bronchi in young smokers. Am Rev Resp Dis 105: 768-75, 1972.	193
SPAIN D M: The distribution of tracheobronchial metaplasia (regenerative hyperplasia) (age, sex, cigarette smoking and relations to reid index) preliminary report.  Pub Health Serv Pub 1879: 183-86, 1969.	194
SPAIN D M, BRADESS V A, TARTER R and MATERO A: Metaplasia of bronchial epithelium. JAMA 211: 1331-34, 1970.	195
THURLBECK W M and ANGUS G E: A distribution curve for chronic bronchitis.  Thorax 19: 436-42, 1964.	196
THURLBECK W M and ANGUS G E: The variation of reid index measurements within the major bronchial tree. Am Rev Resp Dis 95: 551-55, 1967.	197
THURLBECK W M, FRASER R G and BATES D V: The correlation between pulmonary structure and function in chronic bronchitis, emphysema and asthma. Med Thorac 22:295-303, 1965.	198
THURLBECK W M, ANGUS G E and PARE J A P: Mucous gland hypertrophy in chronic bronchitis, and its occurrence in smokers. Br J Dis Chest 57: 73-78, 1963.	199
WEST IND D. W. March of the policy of his policy of the po	Sando Sando

### III-B. ANTEMORTEM STUDIES

Chronic bronchitis diagnosed clinically by measurements of lung function is seen both in smokers and in nonsmokers. Simonson (1965) reported a series from Sweden consisting of the following: of 30 males, 2 were nonsmokers and of 35 females 5 were nonsmokers. Half of the patients reported some allergy connection and one-third were exposed to dusts in their occupations.

A coordinated study in four Canadian cities included 216 patients with chronic bronchitis, 2 of whom were nonsmokers. In the follow-up data of the Toronto group, although the patients as a whole showed a decrease in ventilatory function and in diffusing capacity, 10 of the 44 patients had stopped smoking and a further 6 had reduced their cigarette consumption (Bates et al., 1966). No detectable improvement occurred following the reduction of smoking.

Of 260 patients with chronic bronchitis reported in Germany there was a higher proportion of nonsmokers than in other countries. Wendel (1968) counted 66 of these patients who were nonsmokers. In England, Simpson (1965) reported that 18 out of 157 patients were nonsmokers, and Caird et al. (1972) stated that out of 47 patients 18 were nonsmokers. In Egypt, Megaked (1966) reported that 8 out 65 bronchitic patients were nonsmokers. In the United States, Massaro and Bottoms (1967) noted that 27 out of 327 patients were nonsmokers. The overall incidence of nonsmokers in all these reports is 13% of bronchitic patients.

Bronchial biopsy has been performed in patients with chronic bronchitis. There was a higher incidence of mucus gland hypertrophy in smokers than in nonsmokers (Megahed et al., 1967).

A few patients with regional bronchitis have been described but the number is too small to allow evaluation of significance of smoking (Anthonisen et al., 1968; Niewoehner et al., 1972).

#### BIBLIOGRAPHY

III. CHRONIC BRONCHITIS: CONCURRENCE OF CIGARETTE SMOKING,
INHALATION OF POLLUTANTS AND INFECTION

# B. ANTEMORTEM STUDIES

	Reprin
ANTHONISEN NR, BASS H, ORIOL A, PLACE R E G, and BATES DV:	
Regional lung function in patients with chronic bronchitis. Clin Sci	•
1987 35: 495-511, 1968.	201
BATES D V, GORDON CA, PAUL GI, PLACE REG, SNIDAL DP and WOOLF CR: Chronic bronchitis. Report of the third and fourth stages of the co-ordinated study of chronic bronchitis in the department of	
verterns affairs, Canada. Med Serv J. Can :1-59, 1966.	202
BRINKMAN G L, BLOCK D L: The prognosis in chronic bronchitis.  Pub Health Serv 1717: 317-26, 1968.	203
BRINKMANN G L, BLOCK D L: Chronic bronchitis in a working population.  J Occup Med 14: 825-27, 1972.	204
BURROWS B and EARLE R H: Prediction of survival in patients with chronic airway obstruction. Am Rev Resp Dis 99: 865-871, 1969.	205
CAIRD F I and AKHTAR A J: Chronic respiratory disease in the elderly. A population study. Thorax 27: 764-68, 1972.	206
FLETCHER C M: Prognosis in chronic bronchitis. Pub Health Service 1717: 309-15, 1968.	207
JONES N L, BURROWS B and FLETCHER C M: Serial studies of 100 patients with chronic airway obstruction in London and Chicago. Thorax 22: 327-35, 1967.	208
MASSARO D and BOTTOMS L: Epidemiology of chronic bronchitis.  Med Times 95: 121-28, 1967.	209
MEGAHED G E: The role of smoking in chronic bronchitis in Egyptians.  J Egypt Med Ass 49: 313-28, 1966.	210
MEGAHED G E, SENNA G A, EISSA M H, SALEH S Z and EISSA H A: Smoking versus infection as the aetiology of bronchial mucous gland hypertrophy in chronic bronchitis. Thorax 22: 271-78, 1967.	211
NIEWOEHNER DE, KLEINERMAN J and KNOKE JD: Regional chronic bronchitis.  Am Rev Resp Dis 105: 586-93, 1972.	212
SHITO K, HAMADA A, YASUTARO Y M, EMURA M, MITANI K, MATSUDA M, INQUE T, SAWAL M, KAWAMURA S and OKUBO M: Studies on chronic bronchitis in Osaka. Jap J Med 6: 105-6, 1967.	213
SIMONSSON B G: Clinical and physiological studies on chronic branchitis.  Acta Allergologica 20: 257-300, 1965.	214
SIMPSON T: Chronic bronchitis and emphysema with special reference to prognosis. Br J Dis Chest 62: 57-79, 1968.	215
WENDEL V H: Zur bedeutung des rauchens für die chronische bronchitis. <u>Seit Ges Ann Med</u> 23: 147-51, 1968.	216

#### III-C. EXAMINATION OF SPUTUM

with chronic bronchitis. The observations relating to cigarette smoking consist of the cellular components of the sputum. Vassar et al. (1960) reported an increase in fluorescent histiocytes in the sputum of cigarette smokers which were absent in that of nonsmokers. They suggested that the fluorescence might be due to the presence of polycyclic hydrocarbons bound to the lipid in the cytoplasm of the histiocyte. Roque and Pickren (1968) performed a histochemical study of lung biopsies. In the histiocytes (also known as alveolar macrophages) of smokers, there was a loss of oxidoreductases and hydrolases that was directly proportional to the amount of fluorescent material stored in the cells.

Masin and Masin (1966) examined sputum specimens and observed an increase in the proportion of lipophages in the smokers as compared with the nonsmokers. They explained the difference as due to a combined effect of irritation of the alveolar lining, increased turnover of alveolar cells with increased injury to macrophages and shift in the relative proportion of lipophages. Alveolar macrophages obtained by endobronchial lavage of cigarette smokers and nonsmokers were examined with the electron microscope (Pratt et al., 1971). In smokers the inclusions are heterogeneous and measure from 0.1 to 20  $\mu$  in diameter; the inclusions in nonsmokers are mostly round or oval and measure from 0.1 to 2.0  $\mu$ .

There is an increase in incidence of metaplasia of bronchial epithelial cells in the sputum of smokers (Robbins, 1966; Fullmer, 1968; Nasiell, 1968; Rimingen, 1968). Since metaplasia also occurs in nonsmokers, it is

Source: https://www.industrydocuments.ucst.edu/docs/ptmk0000

The other techniques for the examination of sputum concern sputum purulence (Miller, 1963), protein turnover (Bonomo and D'Addabbo, 1964), electrolyte content (Nicolas, 1964) and concentration of neuraminic acid (Reid, 1967). The influence of smoking on these parameters has not been investigated.

## BIBLIOGRAPHY

III. CHRONIC BRONCHITIS: CONCURRENCE OF CIGARETTE SMOKING, INHALATION OF POLLUTANTS AND INFECTION

## C. EXAMINATION OF SPUTUM

	Keprin.
BONOMO L and D'ADDABBO A: Albumin turnover and loss of protein into the sputum in chronic bronchitis. Clinica Chemica Acta 10: 214-22, 1964.	217
FULLMER CD: Microscopic observations of sputum of chronic cigarette smokers.  Rocky Mt Med J 65: 13, 1968.	218
FULLMER CD, SHORT JG, ALLEN A and WALKER K: Sputum of chronic cigarette smokers. Rocky Mt Med J 66: 42-46, 1969.	219
MASIN F and MASIN M: Frequencies of alveolar cells in concentrated sputurn specimens related to cytologic classes. Acta Cytologica 10: 362-67, 1966.	220
MILLER D L: Study of techniques for the examination of sputum in a field survey of chronic bronchitis. Am Rev Resp Dis 88: 473-83, 1963.	221
NASIELL M: Sputum-cytologic changes in smokers and non-smokers in relation to chronic inflammatory lung diseases. Acta Pathol Microbiol Scand 74: 205-13, 1968.	222
NICOLAS R: Klinisch-experimentelle sputumuntersuchungen bei chronischer bronchitis. Medicina Thoracalis 21: 223-55, 1964.	223
PRATT SA, SMITH MH, LADMAN AJ and FINLEY TN: The ultrastructure of alveolar macrophages from human cigarette smokers and nonsmokers.  Scand J Clin Lab Invest 24: 331-38, 1971.	224
REID L: Mucus secretion and chronic bronchitis. Medicina Thoracalis 24: 40-43, 1967.	225
RIMINGTON J: Smoking, sputum, and lung cancer. Br Med J 1: 732-34, 1968.	226
ROBBINS W T: Bronchial epithelium in smoking and non-smoking college students. J Am Coll Health Assoc 14: 265-66, 1966.	. 227
ROQUE A L and PICKREN J W: Enzymatic changes in fluorescent alveolar macrophages of the lungs of cigarette smokers. Acta cytologica 420-29, 1968.	228
VASSAR PS, CULLING C and SAUNDERS A M: Fluorescent histiocytes in sputum related to smoking. Arch Pathol 70: 649-52, 1960.	229
$oldsymbol{\omega}_{oldsymbol{G}_{i}}^{oldsymbol{\omega}}$	

#### III-D. AIR POLLUTANTS AS A RISK FACTOR

The role of air pollutants in the pathogenesis of chronic bronchitis is discussed in the articles listed in bibliographic list no. 8.

Briefly, the association of this condition with air pollutants is similar to that with cigarette smoking. There is a parallel relationship between the incidence of chronic bronchitis and either the degree of pollution in the air or the amount of cigarette smoking (Anderson 1962).

The incidence of chronic bronchitis in the general population has been correlated with the following indices of air pollution. Fairbairn and Reid (1958) reported that among London postmen disablement and death were related to the frequency of fog and presumably the level of air pollution. Schoettlin (1962) observed that the prevalence of chronic bronchitis among elderly males was higher among those exposed to five occupational factors: cooking fats, gasoline and oils, coal dust and welding fumes. Among American male college students the incidence of chronic bronchitis detected by evaluation of respiratory function was correlated significantly with ambient temperature (Spodnik et al., 1966). Levels of sulfur dioxide in the air showed a positive correlation with incidence of respiratory illness in the United States (McCarroll et al., 1967; Heimbach, 1971).

The importance of air pollution and cigarette smoking has been assessed by some investigators. Winkelstein (1969), Motley (1971),

Motley and Yanda (1966) and Cohen et al. (1972) have the opinion that cigarette smoking is more significant than air pollution in causing chronic bronchitis. Their evidence is not based on a comparison with subjects free from exposure to pollutants. The experience of Phelps and Koike (1962) in the Tokyo-Yokohama episode of asthmatic bronchitis is unique in that American soldiers who were heavy smokers were exposed to air pollutants. After they were transferred to the United States, although they continued smoking, their bronchitis was relieved. In this episode, air pollution is more important than cigarette smoking in causing bronchitis.

# BIBLIOGRAPHY

III. CHRONIC BRONCHITIS: CONCURRENCE OF CIGARETTE SMOKING, INHALATION OF POLLUTANTS AND INFECTION

# D. AIR POLLUTANTS AS A RISK FACTOR

	Repri
ANDERSON R J: Epidemiologic studies of air pollution. Dis Chest 42: 474-81, 1962.	230
COHEN CA, HUDSON AR, CLAUSEN JL and KNELSON JH: Respiratory symptom spirometry, and oxident air pollution in nonsmoking adults. Am Rev Resp Dis 105: 251-61, 1972.	is, 231
	231
FAIRBAIRN AS and REID DD: Air pollution and other local factors in respiratory disease. Br J Prev Soc Med 12: 94-103, 1958.	232
HEIMBACH J A Jr: The correlation of polluted air with tree growth and lung disease in humans. Compr Biol Med 1: 243-53, 1971.	233
McCARROLL J, CASSELL E J, WOLTER D W, MOUNTAIN J D, DIAMOND J R and MOUNTAIN I M: Health and the urban environment. V. Air pollution and illness in a normal urban population. Arch Environ Health 14: 178-83, 1967.	234
MOTLEY H L: Clinical aviation and zerospace medicine. Environmental air pollution effect on pulmonary function. Aerosp Med 42: 1108-10, 1971.	235
MOTLEY H L and YANDA R: Environmental air pollution, emphysema and ionized air. Dis Chest 50: 343-52, 1966.	236
PHELPS H W and KOIKE S: Tokyo-Yokohama Asthma. The rapid development of respiratory distress presumable due to air pollution. Am Rev Resp Dis 86: 55-63, 1962.	237
SCHOETTLIN C E: The health effect of air pollution on elderly males.  Am Rev Resp Dis 86: 878-97, 1962.	238
SPODINK M J Jr, CUSHMAN G D, KERR D H, BLIDE R W and SPICER W S:  Effects of environment on respiratory function.  Arch Environ Health  13: 243-54, 1966.	239
WINKELSTEIN W Jr, and KANTOR S: Respiratory symptoms and air pollution in an urban population of northeastern United States. Arch Environ Health 18: 760-67, 1969.	240

#### III-E. INFECTION AS A RISK FACTOR

The basic lesion in chronic bronchitis is inflammation, which is largely due to pathogens. The infectious process may either be the primary cause of the inflammation or may supercede the initial irritation of the bronchial mucosa by chemical substances. There are defence mechanisms which protect the bronchial passages from bacterial or viral infections (see reviews by Green, 1968 a, b, and Rylander, 1969). The process of clearance of bronchial passages is accomplished by the ciliated epithelium and the mucus-secreting goblet cells. The effect of cigarette smoking on mucociliary clearance is discussed in Section VI. The next paragraphs discuss the influence of cigarette smoking on the bronchial infections, which in turn would lead to chronic bronchitis.

The incidence of the common cold is not higher in smokers than in nonsmokers (Kler, 1945). Amon 143 allergic subjects between 18 and 35 years of age, who were challenged with infectious secretions taken from patients with a common cold, 45% developed colds as compared with 31% among 693 nonallergic individuals, who were inoculated in the same way.

Among both groups of subjects, 35% of smokers and 34% of nonsmokers developed colds, indicating that smoking did not influence the susceptibility to the common cold (Dowling et al., 1957). This is the most important evidence that smoking does not increase the susceptibility to the common cold.

Prevalence studies have produced the opposite conclusion. On the basis of symptoms of acute repiratory infections, the following groups show

iourcea hitiosi//www.industrydocumentsiuesiaedu/docs/ofmkouo

a higher incidence among smokers than nonsmokers: 118 adults residing in Cleveland (Boake, 1958); 175 student nurses in Canada (Parnell et al., 1966); 191 high school students in New Jersey (Haynes et al., 1966); 1,811 male college students in South Carolina (Finklea et al., 1969).

There are other observations relating to the influence of smoking on the incidence of bronchial infections. Air pollution has been shown by Douglas and Waller (1966) to relate to frequency of respiratory infection in children residing in London. The antibacterial activity of the saliva is enhanced by thiocyanate ion which may be absorbed from cigarette smoke (Courant, 1967). Patients with pulmonary emphysema show an elevation of immunoglobulin A(Ig A) in the blood, which has been interpreted as a sign of the presence of bronchopulmonary infection (Biegel and Krumholz, 1968). The concentration of immunoglobulin A in the saliva is also elevated in patients with chronic pulmonary disease, but this was not influenced by cigarette smoking (Lewis et al., 1970). In the same study cigarette smoking suppressed production of immunoglobulin A in the saliva of normal subjects. These observations on immunoglobulin do not explain the association between cigarette smoking and respiratory infections.

# \ III. CHRONIC BRONCHITIS: CONCURRENCE OF CIGARETTE SMOKING INHALATION OF POLLUTANTS AND INFECTION

E.	INFECTION	ASA	RISK	FACTOR

	Kepri
BIEGEL A A and KRUMHOLZ R A: An immunoglobulin abnormality in pulmonary emphysema. Am Rev Resp Dis 97: 217-22, 1968.	241
BOAKE W C: A study of ilness in a group of Cleveland families. XVIII. Tobacco smoking and respiratory infections. N Engs J Med 259: 1245-49, 1958.	242
COURANT P: The effect of smoking on the antilactobacillus system in saliva.  Odontol Revy 18: 251-61, 1967.  DOUGLAS J W B and WALLER R E: Air pollution and respiratory infection in children. Br J Prev Soc Med 20: 1-8, 1966.	243 244
DOWLING HF, JACKSON G G and INOUYE T: Transmission of the experimental common cold in volunteers. II. The effect of certain host factory upon susceptibility. J Lab Clin Med 50: 516-25, 1957.	245
FINKLEA JF, SANDIFER SH and SMITH DD: Cigarette smoking and epidemic influenza. Am J Epidemiol 90: 290-99, 1969.	246
GREEN G M: Pulmonary antibacterial mechanisms and the pathogenesis of pulmonary disease. Yale J Biol Med 40: 414-29, 1968A.  GREEN G M: Pulmonary clearance of infectious agents. Ann. Rev Med	247
GREEN G M: Pulmonary clearance of infectious agents.  19: 315-36, 1968B.	248
HAYNES W F Jr, KRSTULOVIC V J and BELL A L L: Smoking habit and incidence of respiratory tract infections in a group of adolescent males.  Am Rev Resp Dis 93: 730-35, 1966.	249
KLER J H: An analysis of colds in industry. Trans Am Acad Ophthalmol Otolaryngol 201-7, 1945.	250
LEWIS DM, LAPP NL and BURRELL R: Quantitation of secretory immunoglobulin  A in chronic pulmonary disease. Am Rev Resp Dis 101: 55-61, 1970.	251
PARNELL J L. ANDERSON D O and KINNIS C: Cigarette smoking and respiratory infections in a class of student nurses. New Engl J Med 274: 979-84, 1966.	252
RYLANDER R: Environmental air pollutants and lung defense to airborne bacteria. Pub Health Serv Pub 1879: 297-304, 1969.	253

There is no available model in the laboratory which mimics the pathological and functional features of the human form of chronic bronchitis. The reported experiments pertain to one of several underlying processes in chronic bronchitis, such as mucus secretion, bronchial epithelial changes, nature of the alveolar lining and the influence of infection (see reviews by Pattle, 1958, 1965 and Green, 1968 and 1969). There are no reported measurements of pulmonary function in these animals to validate the claim that the experiments can be applied to elucidate the effects of cigarette smoking in man.

1. Rabbit. Proetz (1939) reported the first observations on the effect of cigarette smoke on the airways of the rabbit. By visual examination he noted the accumulation of tar, chiefly in the nasal cavity. Holland et al. (1958, 1963) impregnated cigarettes with radioactive arsenic to trace absorption of smoke in the rabbit. Exposure for one to two years caused pathological changes in the respiratory epithelium of the nose and tracheobronchial mucosa, but no measurements were made prior to sacrificing the animals. The remaining observations in the rabbit relate to the alveolar macrophages investigated in vitro (Heese and Myrvik, 1967; Myrvik and Evans, 1967; Weissbecker et al., 1969). The water-soluble constituents of cigarette smoke depress protein synthesis in rabbit alveolar cells (Yeager, 1969). The depressant action of cigarette smoke on phagocytic activity of alveolar macrophages may be prevented by gluthathione and cysteine (Green and Carolyn, 1967; Green, 1968 a, b).

- 2. Guinea pig. Rylander (1968, 1969) used the guinea pig to investigate the pulmonary defence mechanism against airborne bacteria. Animals exposed to cigarette smoke showed a reduction in the number of radioactive bacteria killed, as compared with the controls. This was interpreted as an effect of smoke on the mucus flow which clears the lung. There was no difference in the reduction of viable bacteria, indicating that the efficiency of the bactericidal mechanisms had not been impaired. In a subsequent report, Rylander (1971) described a reduction in the number of free lung cells in the fluid used for lavaging the airways. However, after two to four weeks of exposure there was an increase in the number of macrophages. This biphasic response indicates a tolerance to the reduction of cells in response to cigarette smoke. An increase in the number of pulmonary cells from chronic exposure has been reported by Flint et al. (1971). The depression of oxidative phosphorylation of pulmonary cells noted by Kyle and Riesen (1970) after 2 1/2 days' exposure needs to be examined in animals exposed for several weeks.
- 3. Rat. Exposure of the rat to cigarette smoke for six weeks produced an increase in the number of goblet cells in the trachea and intrapulmonary airways (Lamb and Reid, 1969). The effect cannot be induced by inhalation of extremes of temperature, indicating that this is not the important factor in smoke which increases the number of goblet cells (Jones et al., 1971). The addition of phenylmethyloxadiazole to the tobacco protected the rat from the increase in goblet cell count. This was the first effective means of protection found against changes produced in bronchial epithelium by tobacco smoke (Jones et al., 1972). The condition of pulmonary

macrophages in response to air pollutants other than cigarette smoke has been studied by Gross et al. (1969).

- 4. Mouse. Leuchtenberger et al. (1958, 1960) and Leuchtenberger and Leuchtenberger (1965) reported the cytochemical and cytological alteration in the bronchi of mice exposed to cigarette smoke. The presence of bronchitis was diagnosed by appearance of proliferative epithelial changes and metaplasia. These changes relate to carcinogenesis rather than to bronchitis. There is a decrease in bacterial clearance following an acute exposure to cigarette smoke (Laurenzi et al., 1963). There is an increase in susceptibility to naturally acquired infection of the respiratory tract in mice exposed to cigarette smoke (Wynder et al., 1968). In experiments on mice challenged with Klebsiella pneumoniae or Diplococcus pneumoniae, the decrease in resistance was shown to be transitory. When the time interval between smoke exposure and infectious challenge was lengthened from 1 to 48 hours, there was no demonstrable decrease in resistance to respiratory infection (Spurgash et al., 1968).

  1005051376
- reported the histological and ultrastructural changes in the bronchial epithelium of dogs exposed to cigarette smoke for 24 to 421 days. The atypical nuclei noted in the epithelium do not relate to chronic bronchitis and are beyond the scope of this review.

F. ANIMAL EXPERIMENTS RELATING TO CHRONIC BRONCHITIS	·
	Reprint
UERBACH O, HAMMOND E C, KIRMAN D, GARFINKEL L and STOUT A P:  Histologic changes in bronchial tubes of cigarette-smoking dogs.  Cancer 20: 2055-66, 1967.	254
LINT G L, MAXWELL K W and RENZETTI A D: Influence of cigarette smoke on guinea pigs. Effect on pulmonary cells and serum antitrypsin levels.  Arch Environ Health 22: 366-69, 1971.	255
RASCA J M, AUERBACH O, PARKS V R and JAMIESON J D: Electron microscopic observations of the bronchial epithelium of dogs. I. Control Dogs.  Exp Mol Pathol 9: 363-79, 1968A.	256
RASCA J M, AUERBACH O, PARKS V R and JAMIESON J D: Electron microscopic observations of the bronchial epithelium of dogs. III Smoking dogs.  Exp Mol Pathol 9: 380-99, 1968B.	257
REEN G M: Protection of alveolar macrophages from the cytotoxic activity of cigarette smoke by glutathione and cysteine. J Clin Invest 47:  42a-43a, 1968A.	258
REEN G M: Pathophysiology of the alveolar macrophage system. Public Health Serv. 1786: 463-80, 1968B.	259
REEN G M: Cigarette smoke: protection of alveolar macrophages by glutathione and cysteine. Science 162: 810-11, 1968C.	260
GREEN G M: The response of the alveolar macrophage system to host and environmental changes. Arch Environ Health 18: 548-50, 1969.	261
FREEN G M and CAROLIN D: The depressant effect of cigarette smoke on the in vitro antibacterial activity of alveolar macrophages. N Engl J Med 276: 421-27, 1967.	<b>2</b> 62
GROSS P, deTREVILLE T P, TOLKER E B, KASCHAK M and BABYAK M A:  The pulmonary macrophage response to irritants. Arch Environ Health 18: 174-85, 1969.	263
HEISE ER and MYRVIK: Secretion of lysozyme by rabbit alveolar macrophages in vitro. J Reticuloendothel Soc 4: 510-23, 1967.	264
on the respiratory system of the rabbit. Cancer 16: 612-15, 1963.	265
HOLLAND R H, WILSON R H, MORRIS D, McCALL M S and LANZ H: The effect of cigarette smoke on the respiratory system of the rabbit. Cancer 11: 709-12, 1958.	266
JONES R, BAETJER A M and REID L: Effect of extremes of temperature and humidi on the goblet cell count in the rat airway epithelium. Br J Ind Med 28: 369-73, 1971.	ty 267
JONES R, BOLDUC P and REID L: Protection of rat bronchial epithelium against tobacco smoke. Br Med J 2: 142-44, 1972.	268

Source: https://www.industrydocuments.ucsf.edu/docs/pfmk0000

	Bibliography III-F	lepri
	KYLE J L and RIESEN W H: Stress and cigarette smoke effects on lung mitochondrial phosphorylation. Arch Environ Health 21: 492-97, 1970.	269
	LAMB D and REID L: Goblet cell increase in rat bronchial epithelium after exposure to cigarette and cigar tobacco smoke. Br Med J 1: 33-35, 1969.	270
	LAURENZI GA, GUARNERI JL, ENDRIGA R B and CAREY JP: Clearance of bacteria by the lower respiratory tract. Science 142: 1572-73, 1963.	271
	LEUCHTENBERGER V C and LEUCHTENBERGER R: Kumulative wirkung von zigaretenrauch-inhalation und influenza-virus-infektion bei der entstehung von atypischen proliferationen im bronchialepithel von mausen. Oncologia 19: 81-104, 1965.	272
	LEUCHTENBERGER C, LEUCHTENBERGER R and DOOLIN PF: A correlated histological cytological and cytochemical study of the tracheobronchial tree and lungs of mice exposed to cigarette smoke. I. Bronchitis with atypical epithelial changes in mice exposed to cigarette smoke. Cancer 11: 490-526, 1958.	273
	LEUCHTENBERGER C, LEUCHTENBERGER R, ZEBRUN W and SHAFFER P:  A Correlated histological cytological and cytochemical study of the  tracheobronchial tree and lungs of mice exposed to cigarette smoke. II. Varying responses of major bronchi to cigarette smoke, absence of bronchogenic carcinoma after prolonged exposure, and disappearance of bronchial lesions after cessation of exposure. Cancer: 13: 721-32, 1960.	274
•	MYRVIK QN and EVANS DG: Metabolic and immunologic activities of alveolar macrophages. Arch Environ Health 14: 92-96, 1967.	275
	PATTLE R E: Properties, function, and origin of the alveolar lining layer.  Proc R Soc Biol 148: 217-40, 1958.	276
	PATTLE R E: Surface lining of lung alveoli. Physiol Rev 45: 48-79, 1965.	277
•	PROETZ A: Some preliminary experiments in the study of cigarette smoke and its effects upon the respiratory tract. Ann Otol 48: 176-94, 1939.	278
•	RYLANDER R: Pulmonary defence mechanisms to airborne bacteria.  Acta Physiol Scand Suppl 306: 1-89, 1968.	279
	RYLANDER R: Alterations of lung defence mechanisms against airborne bacteria.  Arch Environ Health 18: 551-55, 1969.	280
	RYLANDER R: Free lung cell studies in cigarette smoke inhalation experiments.  Scand J Resp Dis 52: 121-28, 1971.	281
	SPURGASH A, EHRLICH R and PETZOLD R: Effect of cigarette smoke on resistance to respiratory infection.  Arch Environ Health 16: 385-91, 1968.	282
	WEISSBECKER L, CARPENTER R D, LUCHSINGER P C and OSDENE T S: In vitro alveolar macrophage viability. Effect of gases. Arch Environ Health 18: 756-59, 1969.	283
	WYDNER E.L., TAGUCHI K.T., BADEN V and HOFFMANN D: Tobacco carcinogenesis.  I.M., The effect of cigarette smoke on respiratory tract of mice after passive  1-1 laton. Cancer 21: 134-53. 1968.	284

YEAGER H Jr: Alveolar cells: depressant effect of cigarette smoke on protein synthesis. Proc Soc Exp Biol Med 131: 247-50, 1969.

Pulmonary emphysema-bronchitis complex, also known as chronic obstructive pulmonary disease, is discussed in this section separately from the sections dealing with its individual components (Sections II and III). It has been the practice of the United States Public Health Service to base their mortality statistics on a combination of pulmonary emphysema and chronic bronchitis even though separate death rates are available. It appears that combining the two entities under a single name has improved the results of statistical analysis in establishing the high incidence of deaths from pulmonary disease among smiokers as compared with nonsmokers.

The publications relating to the pulmonary emphysema-bronchitis complex contained

are / in additional bibliographic lists as follows:

- No. 9. Pulmonary emphysema-bronchitis complex: etiology and pathology.
- No. 10. Pulmonary emphysema-bronchitis complex: clinical studies.

Most of the publications in the above-mentioned lists emphasize the difference between the two entities, but the practice of combining both in mortality statistics has not been discontinued in the United States and has also been adopted in foreign countries.

#### IV-A. MORTALITY STATISTICS IN THE UNITED STATES

In 1958 Dorn reported the mortality experience of male policy holders of U. S. Government Life Insurance during the 2 1/2-year period 1954-1956. The death rate for persons who had smoked regularly was 36% greater than that for persons who had never smoked. in another way, with 1.00 as the death rate of nonsmokers,, that of smokers was 1.36. The mortality ratios for the diseases of the lung were as follows: cancer of the lung, 9.85, chronic obstructive pulmonary disease, 3.27. It was on this basis that the United States Public Health Service founded their conclusion that cigarette smoking causes cancer of the lung and chronic obstructive pulmonary disease. The mortality ratio of deaths from coronary heart disease was 1.63 and it is on this lower ratio that cigarette smoking was regarded as a risk factor. It will be noted that the mortality ratio of chronic lung disease is closer to that of heart disease than to that of lung carcinoma, so that it would have been more logical to draw a similar conclusion for chronic lung disease and heart disease. subsequent publications of Dorn (1959, 1960, 1961), Kahn (1961) and Weir and Dunn (1970) recounted these mortality ratios and called attention to the declining death rate from disease of the respiratory system since 1900 (from 26% in 1900 to 8% in 1959). However, during the past decade the death rate increased significantly for a group of chronic respiratory diseases, including chronic bronchitis, emphysema, fibrosis and pneumor

The second mortality study, in which 1,078,894 persons were enrolled, was conducted by the American Cancer Society in 1959 and 1960.

The mortality rate was higher for smokers than for nonsmokers. The mortality ratio for the pulmonary emphysema-bronchitis complex among Hammond and Horn, 1958 smokers as compared with nonsmokers was 3.27 (Hammond, 1959,, 1964,, 1966). The details of the statistical analysis of the mortality figures are contained in the publications relating to Smoking and Health. The regional analyses have been published separately for Georgia (Hammond and Letton, 1965), Illinois (Hammond and Street, 1964), Iowa (Hammond and Stocks, 1964), Kansas (Hammond et al., 1964), Minnesota (Hammond et al., 1965), Missouri (Hammond and Wilson, 1965), New York State (Hammond et al., 1965), and Ohio (Hammond and Gerber, 1965). The similarities in the mortality statistics of these areas have been emphasized but their dissimilarities have been ignored.

#### BIBLIOGRAPHY

# IV. PULMONARY EMPHYSEMA-BRONCHITIS COMPLEX

## A. MORTALITY STATISTICS IN THE UNITED STATES

[유통] 경우 경우 경우 아이들 아이들 아이들 아이들 아이들 아이들 때문에 가장 아이들 때문에 다른 사람들이 되었다. 그 사람들은 아이들 아이들 아이들 아이들 때문에 가장 아이들 때문에 가장 아니들 아이들 때문에 가장 아니들 때문에 가장 아니를 때문에 가	Reprint
DORN H F: The mortality of smokers and nonsmokers. Am Stat Ass Proc Soc Stat Sec 34-71, 1958.	286
DORN H F: Tobacco / consumption and mortality from cancer and other diseases.  Public Health Report 74: 581-73, 1959.	287
DORN H F: Tobacco consumption and mortality from cancer and other diseases.  Acta Un Int Cancer 16: 1653-65, 1960.	288
DORN HF: The increasing mortality from chronic respiratory diseases.  Am Stat Ass Proc Soc Stat Sec 148-52, 1961.	289
HAMMOND E C: Discussion. Am Stat Ass Proc Soc Stat Sec 60-64, 1959.	290
HAMMOND E C: Smoking in relation to mortality and morbidity. Findings in.  first thirty-four months of follow-up in a prospective study started in  1959. J Natl Cancer Inst 32: 1161-88, 1964.	291
HAMMOND E C: Smoking in relation to the death rates of one million men and women. Natl Cancer Inst Monogr 19: 127-204, 1966.	292
HAMMOND E.C., BERGLUND L and KASLOW R: Smoking habits and disease in Minnesota. Minn Med 48: 44-49, 1965.	293
HAMMOND E C, FENT L S, REED D C and WILSON J L: Smoking habits and health in Kansas and other central states. J Kansas Med Soc 65: 586-90, 1964.	294
HAMMOND E C and GERBER L F: Smoking habits and disease in Ohio:  Ohio State Med J 61: 134-37, 1965.	295
HAMMOND E C and HORN D: Smoking and death rates report on forty-four months of follow-up of 187,783 men. JAMA 166: 1298-1308, 1958.	296
HAMMOND E C and LETTON A H: Smoking habits and health in Georgia and other Southern states. J Med Assoc Ga 54: 278-81, 1965.	297
HAMMOND E C and STOCKS A: Smoking habits and health in Iowa and neighboring states. J Iowa Med Soc 54: 563-67, 1964.	298
HAMMOND E C and STREET E C: Smoking habits and disease in Illinois.  Illinois Med J 126: 661-65, 1964.	299
HAMMOND E C, GRIETHUYSEN THV, DIBELER JB, SNEDDON A M and HALLIGAN W: Smoking habits and disease in New York State.  New York State J Med 65: 2557-61, 1965.	300
HAMMOND E C and WILSON O: Smoking habits and disease in Missouri.  Missouri Med 62: 109-12, 122, 1965.	. 301

Bibliography	<u>IV - A</u>	Repri
	d ROSENTHAL T: Tobacco and health. Charles C Thomas, Springfield, pp172-90.	302
report	The Dorn study of smoking and mortality among U.S. veterans: on eight and one-half years of observations. Natl Cancer Inst Monogr 125, 1966.	303
	Tobacco smoking: Some hints of its biologic hazards. Ohio State 46: 1165-70, 1950.	. 304
	nd DUNN J E Jr: Smoking and mortality: a prospective study. 25: 105-12, 1970.	<b>3</b> 05

The frequency of fatal chronic pulmonary disease varies in different countries (Reid and Fletcher, 1971). There is a wide range in the sex ratio of death in males and females, from 2.0 in Japan to 9.9 in Finland, with Canada, Great Britain and the United States between the extremes. These variations do not relate entirely to differences in cigarette consumption but indicate the association of other factors, some identified and other sunidentified.

1. Canada. A questionnaire sent to 92,000 veterans from 1956 from pulmonary disease to 1962 revealed a higher death rate/among both smokers and nonsmokers (Canadian Department of National Health and Welfare, 1966; Best et al., 1967). The mortality ratio for all smokers as compared with nonsmokers was 1.33. The ratios for specific diseases were as follows: lung cancer, 14.91; bronchitis and emphysema, 8.71; and coronary heart disease, 1.60. The results of this survey were included in the report of the United States Public Health Service. As in the United States, an increasing number of deaths in Canada during the past decade have been due to chronic obstructive disease. The provincial statistics vary in the assignment of deaths due to pulmonary emphysema. Manitoba having the highest proportion of deaths caused by emphysema and Ontario the highest proportion of deaths due to chronic bronchitis. Cigarette consumption when expressed per person (15 years of age and over) does not correlate with the distribution of chronic pulmonary obstructive disease (Anderson, 1963, 1968).

85

Source: https://www.industrydocuments.ucsf.edu/docs/pfmk0000

in patients with a chronic cough.

3. Israel. The mortality statistics attributed to chronic bronchitis or to emphysema from 1960 to 1964 have been reviewed by Cohen (1967).

There has been a rise in such deaths, which are more frequent among immigrants from Africa and Asia than among those from Europe. The author attempted to relate the differences among immigrants to smoking habits surveyed in 1958. The heavy smokers, those using more than 20 cigarettes daily, constituted 27% of all smokers emigrating from Asia and Africa, and 17% of those from Europe. The interpolation of results from another decade has been questioned. Cigarette smoking is not the most important factor accounting for the difference, because the frequency of lung cancer did not show a parallel distribution. The author concluded that the incidence of malnutrition and infection among immigrants from Asia and Africa is a more important factor in contributing to the higher incidence of lung disease.

source: https://www.industrydocuments.ucsr.edu/docs/p/mk000

# BIBLIOGRAPHY

#### V. PULMONARY EMPHYSEMA-BRONCHITIS COMPLEX

# B. MORTALITY STATISTICS IN FOREIGN COUNTRIES

	Reprint
ANDERSON D O: Observations on the classification and distribution of pulmonary emphysema in Canada. Can Med Ass J 89: 709-16,	
1963.	306
ANDERSON D O: Geographic variation in deaths due to emphysema and bronchitis in Canada. Can Med Ass J 98: 231-41, 1968.	307
BEST E W R, WALKER C B, BAKER P M, DELAQUIS F M, McGREGOR J T and McKENZIE A C: Summary of a Canadian study of smoking and health.  Can Med Ass J 96: 1104-8, 1967.	308
CANADIAN DEPARTMENT OF NATIONAL HEALTH AND WELFARE: Trends in death rates for diseases of interest in relation to smoking, Canada 1950 to 1964. 1964.	309
CANADIAN DEPARTMENT OF NATIONAL HEALTH AND WELFARE: A  Canadian study of smoking and health pp 1-7, 19662	310
CANADIAN DEPARTMENT OF NATIONAL HEALTH AND WELFARE: A  Canadian study of smoking and health.  pp 41-50, 1966C	311
COHEN J: Chronic bronchitis and emphysema. Observations on trends and distribution according to death certificate data, Israel 1960 to 1964.  Israel J Med Sci 3: 885-89, 1967.	312
DOLL R and HILL A B: Mortality in relation to smoking: ten years' observations of British doctors. Br Med J 1: 1399-1410, 1964A.	313
DOLL R and HILL AB: Mortality in relation to smoking: ten years' observations of British doctors. Br Med J 1: 1460-67, 1964B.	314
REID D D and FLETCHER C M: International studies in chronic respiratory disease.  Br Med Bull 27: 59-68, 1971.	315

1005051387

## CRITIQUE OF MORTALITY STATISTICS

There are four considerations in accepting the mortality statistics of chronic lung disease to support the thesis that cigarette smoking is one of the important causes of pulmonary emphysema, and chronic bronchitis. They are as follows:

- Sources of error in filing death certificate. The determination of the cause of death is based on the opinion of the physician signing the certificate. The postmortem examination and the clinical picture would establish the diagnosis of pulmonary emphysema or chronic bronchitis. However, the majority of the certificates state the cause of death on the basis of symptomatology, which may not necessarily identify the cause of death. The increase in mortality is dependent to a great extent upon more frequent diagnosis based on incomplete pulmonary function tests and symptomatology rather than upon a true increase in incidence (Barach, 1964; Barach and Segal, 1968; Barach, 1971). For pulmonary emphysema Rosenblatt (1970, 1972) explained the increase in mortality to be an artifact produced by revision of the International List of Causes of Death in 1949, permitting emphysema to be accepted as a primary cause of death. The widespread publicity given to emphysema in scientific and popular publications and the Social Security compensation for disability from emphysema contribute to the record of the sudden increase of deaths from this disease.
- 2, Influence of air pollution on mortality statistics. The mortality statistics for the entire United States have been handled in their entirety,

ignoring important regional differences. There are differences in death rates from pulmonary emphysema and chronic bronchitis which relate to concentrations of socioeconomic classes, provided that air pollutants remain constant (Zeidberg et al., 1967). Comparison of mortality statistics in selected areas of the United States with differences in the level of air pollution reveal that environmental pollutants are important in influencing the death rate (Winkelstein et al., 1967; Buell et al., 1967; Hickey et al., 1967). In Great Britain the incidence of mortality rate from chronic bronchitis is correlated significantly with the level of air pollution (Fry, 1954; Pemberton and Goldberg, 1954; Reid, 1956; Crofton, 1965).

- 3. Factors influencing prognosis of chronic bronchitis and pulmonary emphysema. The most widely known factor that influences the prognosis of a patient with chronic pulmonary disease is cigarette smoking. There are other factors that are equally important and even more so, such as age, sex, geographical altitude, occurrence of cardiac complications, and extent of respiratory insufficiency (Burrows and Earle, 1964; Mitchell et al., 1964; Renzetti et al., 1966; Sukumalchantra et al., 1966; Renzetti, 1967; Tanner, 1967; Asmundsson and Kilburn, 1969; Davis and McClement 1969; Markush, 1969. The role of each of these factors in determining the longevity of patients with chronic pulmonary disease has been determined, but their importance relative to cigarette smoking is not known.
- 4. Inherent characteristics of eigarette smokers. It has been stated in Section II that smokers may have genetic factors which account for the higher incidence of pulmonary emphysema as compared with nonsmokers.

Although some have been identified, others have not. In a recent study the low incidence of pulmonary emphysema among Seventh Day Adventists, has been explained largely by the fact that they are nonsmokers (Dysinger and Lemon, 1963; Lemon and Walden, 1966). The possibility that members of the families of this religious order may have inherited a trait that renders them resistant to pulmonary disease has not been excluded. The mortality encountered in alcoholics has been reported to be associated with a high incidence of smoking (Rankin et al., 1969). Excessive drinkers are usually heavy smokers (Bailey et al., 1969). Although there are known common causes of death in the alcoholic, chronic pulmonary disease is not one of them. Opium smokers develop chronic obstruction (Da Costa et al., 1971, 1972). The extent of alcoholism and drug abuse has not been considered in the case of smokers dying of chronic pulmonary disease.

•	CDIMICITE	$\sim$ $\sim$	MADRATI	~~~	OT A TIOTICO
<i>-</i> •	CKITIOUE	OF	MURIALI	L X	STATISTICS

### [155일 : 1964년 - 1965] - 1965 - 1965 - 1965 - 1965 - 1965 - 1965 - 1965 - 1965 - 1965 - 1965 - 1965 - 1965 -	Keprint
ASMUNDSSON T and KILBURN K H: Survival of acute respiratory failure. A study of 239 episodes. Ann Intern Med 70: 471-85, 1969.	iy 316
BAILEY M B, HABERMAN PW and SHEINBERG J: Identifying alcoholics in population surveys. Millbank Mem Fund Quart 47: 235-49, 1969.	317
BARACH A L: Pulmonary emphysema: Pathogenesis and the illusion of increased incidence. Med Counterpoint 3: 40-43, 1971.	318
BARACH A'L: Physical exercise in breathless subjects with pulmonary emphysem including a discussion of cigarette smoking. Dis Chest 45: 113-28, 1964.	a, 319
BARACH A L and SEGAL MS: The increased recognition and incidence of chronic bronchitis and pulmonary emphysema. Ann Allergy 26: 353-57, 1968.	320
BUELL P, DUNN J E and BRESLOW L: Cancer of the lung and Los-Angeles- type air pollution. Cancer 20: 2139-47, 1967.	321
BURROWS B and EARLE R H: Course and prognosis of chronic obstructive lung disease. N Engl J Med 280: 397-404, 1964.	322
CROFTON E C: A comparison of the mortality from bronchitis in Scotland and in England and Wales. Br Med J 1: 1635-39, 1965.	323
DA COSTA J L, TOCK E P C and BOEY H K: Lung disease with chronic obstruct in opium smokers in Singapore. Thorax 26: 555-71, 1971.	tion 324
DYSINGER PW and LEMON FR: Pulmonary emphysema in a non-smoking population. Dis Chest 43: 17-25, 1963.	325
DA COSTA J L , TOCK E P C and BOEY H K: Chronic obstructive lung disease in Singapore. Am Rev Resp Dis 106: 246-59, 1972.	326
DAVIS A L and McCLEMENT J H: The course and prognosis of chronic obstruction pulmonary disease. Pub Health Serv Pub 1879: 219-34, 1969.	ve 327
FRY J: Chronic bronchitis in general practice. Br Med J 1: 190-94, 1954.	328
HICKEY R J, SCHOFF E P and CLELLAND R C: Relationship between air pollu and certain chronic disease death rates. Arch Environ Health 15: 728-38, 1967.	tion 329
LEMON F R and WALDEN R T: Death from respiratory system disease among seventh-day adventist men.  JAMA 198: 137-46, 1966.	330
MARKUSH R E: National chronic respiratory disease mortality study. II Mortali associated with persistent cough and phlegm in the United States, 1963.  J Chronic Dis 21: 737-48, 1969.	331
MITCHELL RS, WEBB N C and FILLEY GF: Chronic obstructive bronchopulm disease. III Factors influencing prognosis.  89: 878-96, 1964.	onary 332
PEMBERTON J and GOLDBERG C: Air pollution and bronchitis. Br: Med J 2: 567-70, 1954.	333

Bibliography IV-C page 76	Reprint
RANKIN JG, HALE GS, WILKINSON P, O'DAY DM, SANTAMARIA JN and BABARCZY G: Relationship between smoking and pulmonary disease	
in alcoholism. Med J Australia 1: 730-33, 1969.	334
REID DD: Symposium: Chronic bronchitis. Proc Royal Soc Med 49: 767-71,	
1956.	<b>33</b> 5
RENZETTI A D Jr: Prognosis in chronic obstructive pulmonary disease.	
Med Clin NAm 51: 363-71, 1967.	336
RENZETTI A D Jr. McCLEMENT J H and LITT B D: The Veterans administration cooperative study of pulmonary function. Mortality in relation to respiratory function in chronic obstructive pulmonary disease. Am J Med 41: 115-29.	
1966.	337
ROSENBLATT M B: Validity of emphysema mortality data. Rev Allergy	
24: 1019-22, 1970.	338
POCENIE I ATT 14 Political and the second se	
ROSENBLATT M B: Emphysema: historical perspective. Bull NY Acad Med 48: 823-41, 1972.	339
SUKUMALCHANGRA Y, DINAKARA P and WILLIAMS M H Jr: Prognosis of patients with chronic obstructive pulmonary disease after hospitalization for actue ventilatory failure: A three-year follow-up study. Am Rev Resp Dis	
93: 215-22, 1966.	340
TANNER E: The prognosis of respiratory diseases. Proc 9th int Congr Life	
Ass Med 330-43, 1967.	341
WILKELSTEIN W Jr, KANTOR S, DAVIS E W, MANERI CS, and MOSHER W E:	
The relationship of air pollution and economic status to total mortality and selected respiratory system mertality in man. Arch Environ Health	
14: 162-71, 1967.	342
ZEIDBERG L D, HORTON R J M and LANDAU E: Mortality from diseases of the respiratory system in relation to air pollution. Arch Environ Health	
15: 214-24. 1967.	343

# NON-OCCURRENCE OF CHRONIC PULMONARY DISEASE IN SMOKERS

The morbidity statistics are derived from the analysis of a questionnaire on the occurrence of respiratory symptoms or from examination of individuals who undergo a simple test of ventilatory function. All these surveys, show a high incidence of respiratory symptoms and subnormal ventilatory function in smokers as compared with nonsmokers. This is one basis for concluding that cigarette smoking is one of the important causes of chronic obstructive pulmonary disease, also referred to in the preceding section as the pulmonary emphysema-chronic bronchitis complex.

It should be emphasized that the prevalence of respiratory symptoms among smokers ranges from 10 to 40% in various surveys, and that 60 to 90% of smokers do not complain of coughing, expectoration or shortness of breath. Those that do complain do not necessarily suffer from chronic bronchitis or pulmonary emphysema because these symptoms are also characteristic of other chronic pulmonary diseases, such as bronchial asthma, pulmonary tuberculosis and other specific forms of infection of the lung. In the absence of such diseases, the symptoms may originate from exposure to industrial hazards. Inhalation of dust particles and of pollutants has been ignored in attempts to relate cigarette smoking to respiratory symptoms. The following bibliographic lists are appended containing articles relating to morbidity studies and industrial hazards in general:

- No. 11. Morbidity studies.
- No. 12. Industrial hazards.

#### V-A. SPECIFIC FORMS OF PULMONARY DISEASE

Pulmonary emphysema and chronic bronchitis as separate

@:seases are discussed in Sections II and III respectively. The other

forms of chronic pulmonary disease which have been investigated in smokers

are discussed in this section. The monograph by Mork (1962), dealing with

a comparative study of respiratory disease in England, Wales and Norway,

sommarizes the literature on the role of smoking in causing chronic pulmonary

disease.

 Pulmonary tuberculosis. In England, mortality from pulmonary tiberculosis in middle and later life is higher for males than for females. Lowe (1956) noted a higher proportion of smokers among tuberculous p≥tients than among nontuberculous controls and concluded that smoking is ar important cause of the breakdown of healed or quiescent respiratory tuberculosis in adults. Confirmation of the association has been reported by Edwards (1957) in England, by Shah et al. (1959) and Viswanathan et al. (1965) in India, by Nakajima (1966) in Japan, by Lawther (1966) in the 1:etherlands, by Adelstein and Rimington (1967) in England, and by Sussmann (1967) in Germany. Snider et al. (1971) reviewed 4,092 patients discharged from a tuberculosis sanitarium in Chicago and found no correlation between cigarette smoking and chronic airway obstruction. There is a high incidence of emphysema in tuberculous patients but a cause and effect relationship Las not been established (Lancaster and Tomashefski, 1963; Katz and Yanofsky, 1964; Geiser and Steinmann, 1969).

- 2. Pulmonary fibrosis. This rare disease has been examined for its occurrence among smokers and nonsmokers. Cullen et al. (1965) reported two nonsmokers in a group of 12 patients from New York. Weiss (1967, 1969) listed six nonsmokers in a group of 31 patients from Pennsylvania. Among 100 asbestos textile workers in Pennsylvania six nonsmokers were included among the 35 patients diagnosed as having pulmonary fibrosis (Weiss, 1971). A high incidence of smokers has been reported among patients with pulmonary fibrosis in Guyana (Miller et al., 1968, 1971) and Rumania (Lupu et al., 1966).
- : 3. Bronchial asthma and allergy to tobacco. A few asthmatic patients have been reported in the literature who are allergic to tobacco (Jimenez-Diaz and Cuenza, 1935; Rosen, 1946; Rosen and Levy, 1950; Segal and Dulfano, 1965; Blue, 1970). Skin tests against tobacco extract arc usually positive in smokers regardless of whether asthma is present or absent (Pavlik and Cermakova, 1964). The presence of anti-tobacco precipitins has been reported in over half of those who smoke (Aiache et al., 1972). The significance of these observations has not been elucidated. The coexistence of bronchial asthma and pulmonary emphysema has been demonstrated but this does not imply that the former is a cause of the latter (Lowell, 1959; Winkenwerder et al., 1962; Rowe and Rowe, 1965; Horowitz, 1969; Preiser et al., 1969).
- 4. Other diseases. Spontaneous pneumothorax occurs in smokers. Legrand et al. (1971) postulated that smoking is the cause of the disease, where: Zivy (1970), Zivy and Fournier (1970) and Berard and Emonot (1972) discount it. In a survey of 185 patients with pulmonary chickenpox in Australia,

smoking was a highly significant associated factor (Knyvett, 1967). There is a high incidence of peptic ulcer (about 25%) in patients with pulmonary emphysema. Although a majority of these patients are smokers, it is not possible to state that smoking is the cause of these ulcers (Lowell et al., 1956; Kroeker and Leon, 1962). Other factors, such as vagotonia, hypercapnia, hypoxia and genetic factors should be considered.

		Reprint
	EIN A M and RIMINGTON J: Smoking and pulmonary tuberculosis an analysis based on a study of volunteers for mass miniature radiography.	344
griffin e.	<u>Tubercle</u> 48: 219-226, 1967.	344
	JM, CHEMINAT JC; LAKATOS M and MOLINA C: Precipitines anti- tabac et affections des voies respiratoires. Rev de Tuberculose et	
A 4.4 %	Pneumologie 36: 301-7, 1972.	345
BERARI	J and EMONOT A: Enquete sur l'etiologie du pneumothorax spontane	
	dit idiopathique. (Investigation on the etiology of so-called 'idiopathic' spontaneous pneumothorax.) J Med Lyon 1223: 281-90, 1972.	346
BLUE J	A: Cigarette asthma and tobacco allergy. Ann Allergy 28: 110-15, 1970.	347
CULLEN	J J H, KATZ H L and KAEMMERLEN J T. Chronic diffuse pulmonary infiltration and airway obstruction. Am Rev Resp Dis 92: 775-80, 1965.	348
7.		
D'ANEL	LI A, PROSCIA N, SORISIO F, and CARATTI CA: L'enfisema polmonare cronico ostruttivo nel malato ulceroso. Arch Sci Med	
	125: 191-94, 1968.	349
<b>ED</b> WARI	DS J H: Contribution of cigarette smoking to respiratory disease.  Br J Prev Soc Med 11: 10-21, 1957.	350
<b>G</b> EISER	V B and STEINMANN B: Infektionen im alter. Z Gerontologie 2: 69-85, 1969.	<b>3</b> 51
HOROW	ITZ L: Chronic obstructive bronchopulmonary disease (emphysema?) in children with asthma. J Asthma Res 6: 211-17, 1969.	352
JIMINE	Z-DIAZ C and CUENZE BS: Asthma produced by susceptibility to unusual allergens. <u>J Allergy</u> 6: 397-403, 1935.	353
KATZ J	and KUNOFSKY: Tuberculosis as a cause of the increasing mortality from emphysema. Am Rev Reso Dis 89: 673-79, 1964.	354
KNYVE	IT A F: The relation between tobacco smoking and pulmonary chickenpox.  Med J Aust 2: 1197-98, 1967.	355
KROEKI	ER E J and LEON A S: The association of diffuse obstructive pulmonary emphysema and chronic gastroduodenal ulceration. Dis Chest 42: 413-21, 1962.	356
LANCAS	STER J F and TOMASHEFSKI J F: Tuberculosis a cause of emphysematic Am Rev Resp Dis 87: 435-37, 1963.	357
LAWTH	ER P J. Air pollution and respiratory disease. Ned Tijds Genees 110: 2318-19, 1966.	358
LEGRA:	ND M, BOTTO M J and LESOBRE R: Le pneumothoraz spontane.  Recherches etiologiques suggerees par la prevalence masculine de l'affection. (The spontaneous pneumothorax. Etiologic investigation	
·	suggested by the prevalence of the disease in the male subjects).  Le Pournon et le Cocur 27: 827-43, 1971.	359

	Bibliography V-A	page 82	Reprint
	LOWE CR: An association between smoking and respiratory tuberculo Br Med J 2: 1081-86, 1956.	sis.	360
Œ	LOWELL F C: The relationship of allergy to emphysema. Am Rev Re 80: 181-83, 1959.	so Dis	361
	LOWELL F C, FRANKLIN W, MICHELSON A L, and SCHILLER I W.  A note on the association of emphysema peptic ulcer and smoki  N Engl J Med 254: 123-24, 1956.  LUPU D N G, VELICAN C and RACOVEANU C: Fibrosis pulmonar p tabaco. Prensa Med Argent 53: 646-51, 1966.	ng.	362 363
	MILLER G J, ASHCROFT M T, BEADNELL H M S G, WAGNER J C a PEPYS J: The lipoid pneumonia of blackfat tobacco smokers in Guyana. Quart J Med 15: 457-70, 1971.		364
	MILLER G J. BREADNELL H M S G and ASHCROFT M T: Diffuse pulmonary fibrosis and blackfat-tobacco smoking in Guyana.  Lancet 2: 259-60, 1968.		365
	MORK T: A comparative study of respiratory disease in England and Wales and Norway. Acta Med Scand 172 (Suppl 384): 1-100, 1962.		366
	NAKAJIMA T: A study on chronic bronchitis complicated with pulmona tuberculous patients. Kekkahu 41: 337-44, 1966.  PAVLIK I and CERMAKOVA Z: Casna kozni reakce na tabakovy extra		367
<b>)</b>	chorob dychadel. (Early skin reactions against tobacco extract in respiratory system disease). Rozhledy Tuberk Nem Plicnik 24: 629-35, 1964.		368
	PREISER F M, DONNER M W and VAN METRE T E Jr: Comparison of midplane, full-chest tomograms, duffusing capacities, sputu eosinophil content, and other parameters in patients with asthma and emphysema. J Allergy 44: 154-64, 1969.		369
•	ROSEN F L: Bronchial asthma in the young male adult. Ann Allergy 4: 247-60, 1946.		370
	ROSEN F L and LEVY A: Bronchial asthma due to allergy to tobacco in an infant. JAMA 144: 620-21, 1950.	smoke •	371
	ROWE A H and ROWE A Jr. Food allergy its role in emphysema a chronic bronchitis. Dis Chest 48: 609-12, 1965.	nd	372
•.	SEGAL MS and DULFANO MJ: Glomectomy in the treatment of chrobronchial asthma. N Engl J Med 272: 57-63, 1965.	nic	373
•	SHAH J R, WARAWADEKAR M S, DESHMUKH P A and PHUTANE P Institutional survey of pulmonary tuberculosis with special reference to smoking habits. Indian J Med Soi 13: 381-92,	•	374
ζ	SNIDER G L, DOCTOR L, DEMAS T A and SHAW A E: Obstructive disease in patients with treated pulmonary tuberculosis. Am 103: 625-40, 1971.	airway Rev Rosh Dis	375
•	SUSSMANN F H A: Die chronischen unspezifischen lungenerkrankunger sicht des Kreistuberkulosearztes. Z Tuberk 127: 191-202,		376

Source: https://www.industrydocuments

Bibliog	ography V-A	page 83	Repri
VISWA	ANATHAN R, MODY RK, PRASAD SS and SINHA SP: B	ronchial asthma	377
<u></u>			
WEISS	S W: Cigarette smoking and diffuse pulmonary fibrosis. Ar	ch Environ Health	
	14: 564-68, 1967.		378
WEISS	S W: Cigarette smoking and diffuse pulmonary fibrosis. A	m Rev Resn Dis	
***	99: 67-72, 1969.	III Icev Icesp Dis	379
-			4 ST . 2 5
WEISS	SS W: Cigarette smoking, asbestos, and pulmonary fibrosis.	Am Rev Resp Dis	
	104: 223-27, 1971.		380
·			
WINKE	KENWERDER W L, ALLAN W B and FISHER A M: A consi		ara Mira
<u>F</u>	etiology of obstructive emphysema, with especial emphase of allergy. Trans Am Clin Climatol Assoc 74: 1-14, 1		381
	Oraniergy. Itans in One One of the Francisco	. 702.	147 - No.
ZIVY :	P: Le pneumothorax spontane des fumeurs. Le poumon tab	agique.	A Tarting
	Rev Tuberc Pneumol 34: 125-58, 1970.		382
ZIVY	P and FOURNIER E: Tabagisme, emphyseme bulleux et p	- 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1 - 1	1494
	enontane (P.S. I. Vidionathique Ann Med Intern. 121. 1	040 56 1070	

The prevalence of respiratory symptoms among cigarette smokers has been investigated in several states. Some of the surveys consider the additional role of industrial hazards and of air pollution in the causation of symptoms which suggest the presence of chronic pulmonary disease.

disease among 1, 261 residents of Berlin, New Hampshire, was carried out (Ferris and Anderson, 1965; Anderson and Ferris, 1962; Anderson et al., 1963). It was observed that in cigarette smokers, chronic respiratory disease was substantially more prevalent than in nonsmokers of similar age and sex. A repeat study of the group in 1967 showed a lower prevalence of such disease than in 1961. The levels of air pollution were less in the second study than in the first, and it was suggested that this fall accounted for the decrease in prevalence of the disease and improvement in lung function (Ferris et al., 1971). Eckardt (1972) and Ferris et al., (1973) have exchanged correspondence with each other on the results that pulmonary function is better in males smoking 45 cigarettes /day than in those who smoke 1 to 4 cigarettes.

The above studies emphasize the shortcomings of morbidity statistics which attempt to correlate cigarette smoking with incidence of respiratory disease. If cigarette smoking is indeed an important cause of lung disease, continued smoking from 1961 to 1967 would have increased its incidence, but this did not occur. The effects of the reduction in air pollution appear to show that atmospheric impurity is a more important

1005051400

from workers in a pulp mill was compared with one of 124 men from a paper mill. Although the pulp mill workers were exposed to chlorine and sulfur dioxide, there was no significant difference between the two groups in regard to respiratory symptoms and ventilatory function. The working population of both mills together showed a lower prevalence of respiratory disease than did the male population of Berlin, N. H., suggesting that the working population may not be representative of the general population (Ferris et al., 1967).

2. Massachusetts. A sample of the population of Framingham has been examined periodically for symptoms of lung disease, chest x-ray and ventilatory function (Revotskie et al., 1962). The results do not show any correlation between smoking and incidence of symptoms or reduction in ventilatory function. The following groups were found to show a higher incidence of respiratory symptoms among smokers than nonsmokers: 5,000 registrants for military service (Andosca, 1966); 100 college seniors and (Peters and Ferris, 1967); 2,027 university graduates (Peters and Ferris, 1967). In the second group there was no statistically significant difference in vital capacity and forced expiratory volume between smokers and nonsmokers. In the third group, the higher incidence of respiratory symptoms in addition was accompanied by traumatic and genitourinary problems. The authors do not attribute the high incidence of trauma and genitourinary problems among smokers to cigarette smoking, yet they

ascribe the respiratory symptoms to it.

Two groups of workers were surveyed in Massachusetts.

Ferris et al. (1968) examined movie projectionists and noted no difference in ventilatory function between smokers and nonsmokers.

3. Connecticut. A group of 424 persons residing in Southern Connecticut were questioned for incidence of respiratory symptoms (Kelsey et al., 1968). Smokers were affected more frequently than nonsmokers. However, a statistically significant association was observed between severity of symptoms and migration from urban areas. The chronic bronchitics with severe symptoms were most likely to have left urban areas afflicted with air pollution in recent years.

A retrospective study was made of medical absenteeism among 226 male employees of a manufacturing company from 1956 through 1964 (Holcomb and Meigs, 1972). There was a significant association of high absenteeism with cigarette smoking and a more significant association of low absenteeism with cigar or pipe smoking. The authors proposed a hypothesis that adoption of the habit of cigar or pipe smoking at middle age occurred among men with constitutional and absentee behavioral characteristics tending to distinguish them from men with a lifelong cigarette smoking habit. The authors suggested caution in drawing causal conclusions regarding these associations because of considerable overlapping of individual absentee rates among groups divided by smoking habits.

4. New York. The following groups solicited by a questionnaire show a higher incidence of respiratory symptoms among smokers than among nonsmokers: 2,031 insurance policy holders (Short et al., 1939); 1,250 telephone men over 40 years of age (Holland and Stone, 1965; Comstock et al., 1970); and 12,604 postal and transit workers (Densen et al., 1967). In the last-mentioned study, there was a decrease of for ced expiratory volume with age, particularly among white men more than among nonwhite men (Densen et al., 1964, 1967). A similar racial difference was encountered in a survey of 6,314 workers for pulmonary emphysema (Robins et al., 1969).

Two surveys were made of industrial workers. A group of 55 grain handlers in Western New York State (Kleinfeld et al., 1968). There was / no significant difference in vital capacity between smokers and nonsmokers up to 19 years of age but a significant difference in vital capacity and forced expiratory volume between these two types of subject when 20 years old or more. A group of 1,744 individuals exposed to oil mist showed a higher incidence of respiratory symptoms for smokers than for nonsmokers (Ely et al., 1970).

5. Pennsylvania. Boucot et al. (1962) reported that among 6,137 males over 45 years of age there was a higher incidence of chronic cough among smokers than among nonsmokers. A similarity in cough pattern appeared in cigarette, cigar and pipe smokers.

suggesting that tobacco smoke in general rather than cigarette smoke

alone causes cough.

The other reports from Pennsylvania concern coal miners. In a group of 801 anthracite coal miners, smoking, aging and duration of exposure to coal dust increased the prevalence of pneumoconiosis (Tokuhata et al., 1970). Autopsies performed on a group of 144 Appalachian underground coal workers indicated no significant effect of smoking on lung structure in anthracite miners (Naeye et al. 1971). Among bituminous coal miners, the smokers had a greater degree of cor pulmonale and more emphysema and bronchiolar goblet cells. The authors suggested that the effect of smoking among anthracite miners may be obscured by the severe lesions primarily related to the pneumoconiosis. Another possibility is that above-ground forms of air pollution may partially obscure the influence of smoking. In response to a communication from Kibelstis (1972),/emphasized that it is not yet possible to quantitate the individual contributions to emphysema of mine dust, smoking, other air pollutants, infection and other factors of individual susceptibility.

6. West Virginia. The data collected from 258 bituminous miners show a higher incidence of respiratory symptoms among smokers than among nonsmokers (Hyatt et al., 1964). In another survey of 5,540 persons living in five communities, there was a higher incidence of respiratory symptoms among smokers than among nonsmokers (Higgins et al., 1968). The mean of forced expiratory volume of smokers and that of nonsmokers under 50 years of age were not different but these were lower for smokers and nonsmokers

1005051404

over 50 years of age.

That chronic obstructive disease can still occur among miners who are nonsmokers indicates the importance of exposure to coal dust as a causative factor (Rasmussen, 1971; Rasmussen and Nelson, 1971).

The reduction in the ventilatory function of coal miners who are smokers is particularly marked after the end of the working period (Lapp et al., 1972). The prevalence of respiratory symptoms in a mining community is the same both for miners and for nonminers, so that the contamination of the air influences the incidence of lung disease (Enterline and Lainhart, 1967).

7. Maryland. Stebbings (1971 a, b, c, d) has reported the results of examination of 736 white males, 35 to 64 years old, who were nonsmokers. Although the investigation did not include smokers, several problems relating to statistics became apparent. The prevalence of respiratory symptoms or illness as derived from the standard questionnaire does not discriminate adequately between upper respiratory allergic conditions, lower respiratory symptoms and cardiovascular symptoms. The pulmonary function values included sizable instrument and interviewer data that were statistically significant. Social class can influence respiratory symptoms and pulmonary function. Furthermore, the effect of duration of urban residence on pulmonary function vary.

Most of the shortcomings uncovered by Stebbings were overlooked in prevalence studies reported by the United States Public Health Service.

- 8. Southern States. The incidence of respiratory symptoms among smokers is higher than among nonsmokers in the following groups: 1,848 military cadets in Georgia (Finklea et al., 1971), and 408 cotton textile workers in North Carolina (Merchant et al., 1972) and in Georgia (Bouhuys et al., 1969). In Tennessee, the extent of air pollution has been shown to influence the incidence of respiratory symptoms (Zeidberg et al., 1964).
- 9. Michigan. In 1958 a group of 1, 317 men, ranging in age from 40 to 65 years, with industrial exposure in the Detroit area, were examined (Brinkman and Coates, 1962). Cigarette smoking was identified as the most important factor in the etiology of chronic bronchitis diagnosed by the occurrence of expectoration daily for at least six months. After a six-year period, the subjects diagnosed as chronic bronchitics did not show any adverse effect on health, working ability or ventilation even though they continued smoking (Brinkman and Block, 1966, 1972).

Additional surveys have appeared relating to the following:
6,500 residents of Tecumsch (Payne and Kjelsberg, 1964; Higgins and
Kjelsberg, 1967); 1,584 Detroit postal employees 40 years of age or
older (Coates et al., 1965); and 1,058 iron foundry workers (Clarke,
1972). A higher incidence of respiratory symptoms among smokers than
among nonsmokers was reported.

- 10. Minnesota. In a group of 450 railroad employees, there was a correlation between intensity of smoking and the expiratory volumes and incidence of symptoms (Blackburn et al., 1965). In subjects who stopped smoking, the ventilatory function was elevated to "supernormal" level. The authors have interpreted this as strengthening the concept that smokers differ from nonsmokers characteristics other than the smoking habit.
- 11. Midwestern states. The following surveys have revealed a higher incidence of respiratory symptoms among smokers than among nonsmokers: 1,887 white males 43 to 58 years old employed in an industrial plant in Illinois (Sharp et al., 1965); 136 white-collar workers 35 to 64 years of age in Ohio (Krumholz and Hedrick, 1973); 557 high school students in Oklahoma (Addington et al, 1970).
- 12. Colorado. In a group of 172 adult men and women over 40 years of age, ventilatory function in male nonsmokers was significantly better than in smokers, but there was no significant difference between female nonsmokers and smokers (Bower, 1961). In a prevalence study of 633 residents of Glenwood Springs, a strong association was found to exist between incidence of chronic bronchitis and cigarette smoking (Mueller et al., 1971 a, b). The authors reviewed interstudy variations geographic areas in the prevalence of respiratory disease in various /-that there are other factors which determine the incidence of chronic

obstructive pulmonary disease. In a group of 1,500 uranium miners, cigarette smoking did not influence the occurrence of shortness of breath or persistent coughing (Archer et al., 1964). The hazards of radiation exposure are more severe than other factors, such as aging, altitude and cigarette smoking.

12. California. The following surveys indicate a higher incidence of respiratory symptoms among smokers than among nonsmokers: 7,070 industrial workers in Los Angeles County (Balchum et al., 1962); 3,311 longshoremen in San Francisco bay and area (Goldsmith et al., 1962); 630 telephone workers in San Francisco and Los Angeles (Deane et al., 1965). In a group of 38 beryllium workers, ventilatory function was measured at 6-month intervals for 18 to 24 months. The day-to-day variation in values was greater for smokers than for nonsmokers (Dawson, 1966).

#### BIBLIOGRAPHY

#### V. NON-OCCURRENCE OF CHRONIC PULMONARY DISEASE IN SMOKERS

B. MORBIDITY STATISTICS IN THE UNITED STATES	**.
ADDINGTON W W, CARPENTER R L, DUNCAN K A and MOGG K: The	Reprin
Association of Cigarette Smoking With Respiratory Symptoms and Pulmonary Function in a Group of High School Students.  Oklahoma Med Ass J 63: 525-29, 1970.	384
ANDERSON D O and FERRIS B GJr: Role of Tobacco Smoking in the Causation of Chronic Respiratory Disease. NEngl J Med 267: 787-94, 1962.	385
ANDERSON D O, ZICKMANTEL R and FERRIS B G Jr: Response to a Respiratory Survey. Canad Med As 3 J 88: 590-602, 1963.	<b>3</b> 86
ANDOSCA J B: The Incidence of Smoking and the Smoking Habits in 5000  Registrants for Military Service. J Am Geriatr Soc 14: 631-32, 1966.	387
ARCHER V E, CARROL B E, BRINTON H P, and SACCOMANNO G: Epidemiological Studies of some Non-fatal Effects of Uranium Mining. In: Radiological Health and Safety in Mining and Milling of Nuclear Materials, Vol I, International Atomic Energy Agency: Vienna 21-36, 1964.	388
BALCHUM OJ, FELTON JS, JAMISON JN, GAINES RS, CLARKE DR and OWAN T: A Survey for Chronic Respiratory Disease in an Industrial City. Am Rev Resp Dis 86: 675-85, 1962.	389
BLACKBURN H, TAYLOR H L, PARLIN R W, KIHLBERG J and KEYS A: Physical Activity of Occupation and Cigarette Smoking. Arch Environ Health 10: 312-22, 1965.	390
BOUCOT K R, COOPER D A and WEISS W: Smoking and the Health of Older  Men. Arch Environ Health 4: 59-72, 1962.	391
BOUHUYS A, WOLFSON R L, HORNER D W, BRAIN J D and ZUSKIN E:  Byssinosis in Cotton Textile Workers. Respiratory Survey of a Mill with  Rapid Labor Turnover. Ann Intern Med 71: 257-69, 1969.	392
BOWER G: Respiratory Symptoms and Ventilatory Function in 172 Adults Employed in a Bank. Am Rev Resp Dis 83: 684-89, 1961.	<b>3</b> 93
BRINKMAN G L and BLOCK D L: The Prognosis in Chronic Bronchitis.  JAMA 197: 71-77, 1966.	394
BRINKMAN G L and BLOCK D L: Chronic bronchitis in a Working Population.  J Occup Med 14: 825-27, 1972.	395
BRINKMAN G L and COATES E O: The Prevalence of Chronic Bronchitis in an Industrial Population. Am Rev Resp Dis 86: 47-54, 1962.	396
CLARKE NE Sr.: Silicosis and Diseases of Retired Iron Foundry Workers.  Ind Mcd 41: 22-25, 1972.	397
COATES E O Jr. BOWER G C and Reinstein N: Chronic Respiratory Disease	

398

in Postal Employees. JAMA 191: 161-66, 1965.

Bibliography V-B page 94	Reprint
COMSTOCK G W. BROWNLOW W J, STONE R W and SARTWELL P E:  Cigarette Smoking and Changes in Respiratory Findings. Arch Environ  Health 21: 50-59, 1970.	399
DAWSON A: Reproducibility of Spirometric Measurements in Normal Subjects. Am Rev Resp Dis 93: 264-268, 1966.	400
DEANE M.GOLDSMITH JR and TUMA D: Respiratory Conditions in Outside Workers. Arch Environ Health 10: 323-331, 1965.	401
DENSEN P M, JONES E W, BASS H E and BREUER J: A Survey of Respiratory Disease Among New York City Postal and Transit Workers. Environ Res 1: 262-86, 1967.	402
DENSEN P M, JONES E W, BASS H E, BREUER J and REED E: A Survey of Respiratory Disease Among New York City Postal and Transit Workers. <u>Environ Res</u> 2: 277-96, 1964.	403
ECKARDT R E: Chronic Nonspecific Respiratory Disease, Berlin, New Hampshire 1961-1967: A Cross Section Study. Am Rev Resp Dis 105: 854, 1972.	404
ELY TS, PEDLEY SF, HEARNE FT and STILLE WT: A study of mortality, symptoms, and respiratory function in humans occupationally exposed to oil mist. J Occup Med 12: 253-61, 1970.	405
ENTERLINE PE and LAINHART WS: The relationship between coal mining and chronic nonspecific respiratory disease. Am J Public Health 57: 484-95, 1967.	406
FERRIS B G and ANDERSON D O: The prevalence of chronic respiratory disease in a New Hampshire town. Am Rev Resp Dis 86: 165-77, 1965.  FERRIS B G Jr, BURGESS W A and WORCESTER J: Prevalence of chronic respiratory disease in a pulp mill and a paper mill in the United States.  Br J Ind Med 24: 26-37, 1967.	407
FERRIS B G Jr. BURGESS W A and WORCEST ER: Chronic nonspecific respiratory disease in a movie projectionists union. Arch Environ Health 16: 886-91, 1968.	409
FERRIS B G Jr, HIGGINS L T T, HIGGINS M W, PETERS J M, VAN GANSE W I and GOLDMAN M D: Chronic nonspecific respiratory disease, Berlin, New Hampshire, 1961-1967: A cross-sectional study. Am Rev Resp Dis 104: 232-44, 1971.	F 410
FERRIS B G Jr., HIGGINS I T T, HIGGINS M W and PETERS J M: Chronic nonspecific respiratory disease in Berlin, New Hampshire, 1961 to 1967.  A follow-up study. Am Rev Resp Dis 107: 110-22, 1973.	411
FINKLEA J F, HASSELBLAD V, SANDIFER S H, HAMMER D I and LOWRIMORE G R: Cigarette smoking and acute non-influenzal respiratory disease in military cadets. Am J Epidemiol 93: 457-62, 1971.	412
GOLDSMITH J.R., HECHTER H.H., PERKINS N.M. and BORHANI N.O.: Pulmonary function and respiratory findings among longshoremen. Am Rev Resp Dis. 86: 867-74, 1962.	413
HIGGINS I T T, HIGGINS M W, LOCKSHIN M D and CANALE N: Chronic respiratory disease in mining communities in Marion County, West Virginia. Br J Ind Med 25: 165-75, 1968.	414

	Bibliography V-B page 95	Reprin
	HIGGINS MW and KJELSBERG M: Characteristics of smokers and nonsmokers in Tecumseh, Michigan. Am J Epidemiol, 86: 60-77, 1967.	415
$\alpha$	HOLCOMB III H S, and MEIGS J W: Medical absentecism among cigarcte, and cigar and pipe snokers. Arch Environ Health 25: 295-300, 1972.	416
	- HOLLAND W W and STONE R W: Respiratory disorders in United States east coast telephone men. Am J Epidemiol 82: 92-101, 1965.	417
<b>大学</b>	HYATT RE, KISTIN A D and MAHAN TK: Respiratory disease in southern West Virginia coal miners. Am Rev Resp Dis 89: 387-401, 1964.	418
	KELSEY J L, MOOD E W and ACHESON R M: Population mobility and epidemiolog of chronic bronchitis in Connecticut. Arch Environ Health 16: 853-61, 1968.	y 419
	KIBELSTIS J.A: Coal workers' pneumoconiosis. JAMA 221: 301-02, 1972.	420
1964	KLEINFELD M, MESSITE J, SWENCICKI R E and SHAPIRO J: A clinical and physiologic study of grain handlers. Indust Hyg Rev 10: 12-19, 1968.	421
	KRUMHOLZ R A and HENDRICK E C: Pulmonary function differences in normal moking and nonsmoking, middleaged, white-collar workers. Am Rev Resp Dis 107: 225-30, 1973.	<u>s</u> 422
	LAPP N L, HANKINSON J L, BURGESS D B and O'BRIEN R: Changes in ventilatory function in coal-miners after a work shift. Arch Environ Health 24: 204-8, 1972.	423
9	MERCHANT JA, KILBURN KH, O'FALLON WM, HAMILTON JD and LUMSDEN JC: Byssinosis and chronic bronchitis among cotton textile workers. Ann Intern Med 76: 423-33, 1972.	<b>4</b> 24
	MUELLER R E, KEBLE D L, PLUMMER J and WALKER S H: The prevalence of chronic airway obstruction, and respiratory symptoms in a Colorado city.  Am Rev Resp Dis 103: 209-28, 1971A.	425
	MUELLER R E, WALKER S H and PLUMMER J: An epidemiologic study of chronic bronchitis, chronic airway obstruction and respiratory symptoms in a Colorado city. Hartford Hosp Bull 26: 261-97, 1971B.	426
•	NAEYE R L: Coal workers' pneumoconiosis. J Am Med Ass 222: 87, 1972.	427
	NAEYE R L, MAHON J K and DELLINGER W S: Effects of smoking on lung structure of Appalachian coal workers. Arch Environ Health 22: 190-93, 1971	. 428
	PAYNE M and KJELSBERG M: Respiratory symptoms, lung function, and smoking habits in an adult population. Am J Public Health 54: 265-77, 1964.	429
•	PETERS J M and FERRIS B G Jr: Smoking, pulmonary function, and respiratory symptoms in a college-age group. Am Rev Resp Dis 95: 774-82, 1967.	430
e je same Projekt	PETERS J M and FERRIS B G Jr: Smoking and morbidity in a college-age group. Am Rev Resp Dis 95: 783-89, 1967.	431
<b>(</b>	RASMUSSEN D L: Impairment of oxygen transfer in dyspneic, nonsmoking soft coal miners. J Occup Med 13: 300-5, 1971.	432
	RASMUSSEN D L and NELSON C W: Respiratory function in southern Appalachian coal miners. Am Rev Resp Dis 103: 240-48, 1971.	. 433

Bibliography V-B page 96	Repr
REVOTSKIE N, KANNELL W, GOLDSMITH JR and DAWBER TR: Pulmonary function in a community sample. Am Rev Resp Dis 86: 907-11, 1962.	434
ROBINS A B, EPSTEIN H H, DIAMOND P H, PLISHNER M J and FINKELSTEIN H A project for the prevention of disability from chronic respiratory disease in working males. Am J Public Health 59: 77-85, 1969.	I: 43!
SHARP JT, PAUL O, LEPPER MH, McKEAN H, and SAXTON GAJr: Prevalence of chronic bronchitis in an American male urban industrial population. Am Rev Resp Dis 91: 510-19, 1965.	436
SHORT J J, JOHNSON H J, and LEY H A Jr: The effects of tobacco smoking on health. J Lab Clin Med 24: 586-89, 1939.  STEBBINGS JH Jr: Chronic respiratory disease among nonsmokers in Hagerstown,	43
Maryland. Design of the study and prevalence of symptoms. Environ Res 4: 146-62, 1971A.  STEBBINGS J H Jr: Chronic respiratory disease among nonsmokers in Hagerstown	438
Maryland. Problems in the estimation of pulmonary function values in epidemiological surveys. Environ Res 4: 163-92, 1971B.	439
STEBBINGS J H Jr: Chronic respiratory disease among nonsmokers in Hagerstown,  Maryland. Social class and chronic respiratory disease. Environ Res  4: 213-32, 1971C.	44(
STEBBINGS J H Jr: Chronic respiratory disease among nonsmokers in Hagerstown,  Maryland. Effects of urban residence on pulmonary function values.  Environ Res 4: 283-304, 1971D.	44
TOKUHATA GK, DESSAUER P, PENDERGRASS E P, HARTMAN T, DIGON E, and MILLER W: Pneumoconiosis among anthracite coal miners in Pennsylvania. Am J Public Health 60: 441-51, 1970.	44
ZEIDBERG L D. PRINDLE R A and LANDAU E: The Nashville air pollution study. III: Morbidity in relation to air pollution. Am J Public Health 54: 85-97, 1964.	44

#### V-C. MORBIDITY STATISTICS IN GREAT BRITAIN

chronic bronchitis is the more common form of chronic obstructive pulmonary disease in Great Britain. Fletcher (1963) and Reid et al. (1964) noted that there was little difference in the prevalence of productive cough between the United States (Berlin, surveyed New Hampshire) and all areas of Great Britain. However, the symptoms of more advanced bronchitis with infection and dyspnea were

/ more prevalent in British areas than in the United States.

Holland et al. (1965) arrived at a similar conclusion in comparing

the prevalence of symptoms of advanced bronchitis in London and three American cities: Westchester (New York), Baltimore and Washington. The prevalence rates for respiratory symptoms in British male migrants are no greater than for those in native-born Americans, so that the environment is more important than country of birth (Reid et al., 1966). An Anglo-Danish comparison has revealed a higher incidence of respiratory symptoms in England than in Denmark (Olsen and Gilson, 1960).

1. Influence of air pollution. Most of the early comparisons of prevalence rates of respiratory symptoms in Great Britain involved only a single community: 1,300 subjects in London (Oswald et al., 1953); 6,245 clerical workers also in London (Oswald et al., 1955); 631 residents of an agricultural community near Penarth (Higgins et al., 1956; Higgins, 1957); 147 young soldiers at Aldershot (Liebeschuetz, 1959); 2,081 men

over 60 years of age in Birmingham (Edwards et al., 1959), and 92 general practitioners (College of General Practitioners, 1961). These surveys showed that there is a higher incidence of respiratory symptoms among smokers than nonsmokers.

Subsequent observations have emphasized the importance of air pollution in influencing the prevalence of respiratory symptoms.

Higgins and Cochrane (1958) and Higgins (1959) noted a difference in incidence between a community (Annandale) in Scotland and the Vale of Glamorgan in Wales, i.e., between an urban and a rural community.

The extent of influence of cigarette smoking was determined by the amount of air pollution.

The urban factor in chronic bronchitis has been studied by Holland and Reid (1965), who surveyed 293 drivers and vehicle maintenance men in central London and 477 men in three rural areas in Southern England. Although personal smoking habits were shown to be closely related to the frequency of respiratory disturbance, the local levels of air pollution appeared to be the most likely cause of the difference in morbidity between urban and rural areas. A study of 2,342 families revealed a relationship between the prevalence of cough and environmental factors. In the fathers who were exposed to the hazards both of home and of occupation, smoking and social class influenced the prevalence of cough. It was not possible to incidence of specify the role of smoking alone in coughing (Colley and Holland,

2. Incidence in children and young adults. Approximately

10,000 children in secondary and grammar schools completed a
questionnaire relating to their smoking habits (Holland and Elliot,

1968). The frequency of respiratory symptoms was higher
among students who smoked more than five cigarettes a day -- about

30% coughing and 20% expectoration -- as compared with 2% and 5%
among those smoking less than one cigarette a day. Holland et al.

(1969 a, b, c) have summarized the analysis of their results and concluded
that environmental and personal factors are important in the onset of
chronic respiratory disease. Colley and Reid (1970) surveyed 10,000

children from 6 to 10 years old and noted a pronounced social class
gradient in the frequency of chronic cough. The occurrence of
respiratory symptoms in children is influenced by the smoking
habits of the families (Norman-Taylor and Dickinson, 1972).

Source: https://www.industrydocuments.ucsf.edu/docs/pfmk0000

3. Industrial hazards. In 1957 the British Medical Research Council's Pneumoconiosis Research Unit carried out a survey in Staveley, an industrial town (Higgins et al., 1959). The prevalence of respiratory symptoms was higher and ventilatory function was lower in miners exminers and men exposed to chemical fumes than in men who had worked in nondusty jobs. After five and nine years, the follow-up study showed that the annual decline of forced expiratory volume was greater in smokers than in nonsmokers or ex-smokers (Higgins and Oldham, 1962; Higgins et al., 1968, 1969). It was concluded that smoking is a more important factor in the development of respiratory disability than occupation.

Other investigators surveying miners have not identified smoking as a more important factor than industrial hazards in the causation of respiratory symptoms (Ashford et al., 1961; Caplan et al., 1966; Ashford et al., 1970; Ryder et al., 1970; Walker et al., 1971; Lyons et al., 1972).

Among the following industrial workers, smoking contributes to the prevalence of respiratory symptoms: alkaline dust workers (Chivers, 1959), flax workers (Elwood et al., 1965), spinners (Molyneux and Tombleson, 1970), as bestos workers (Regan et al., and Warwick 1971; Muldoon ., 1972) and steel foundrymen (Lowe et al., 1970, 1972; McCallum, 1972). In the last-mentioned type of workmen, the influence of air pollution in the prevalence of respiratory symptoms has been suggested by Lowe et al. (1968).

Source: https://www.industrydocuments.ucsf.edu/docs/pfmk0000

HIGGINS I T.T., COCHRANE A L., GILSON J C and WOOD C H: Population studies of chronic respiratory disease. A comparison of mines, foundryworkers, and others in Staveley, Derbyshire. Br J Ind Med 16: 255-68, 1959.

	page 102	
	HIGGINS IT T, GILSON J C, FERRIS B G Jr, WATERS M E, CAMPBELL H and	
	HIGGINS MW: Chronic respiratory disease in an industrial town: A nine-year follow-up study. Preliminary report. Am J Public Health	
	58: 1667-76, 1968.	459
, d	HIGGINS ITT, GILSON JC, FERRIS BG, WATERS WE, CAMPBELL H, and HIGGINS MW: Chronic respiratory disease in an industrial town: A	
	nine-year follow-up study. Respiration 26: 221-2, 1969.	460
	HIGGINS IT T and OLDHAM PD: Ventilatory capacity in miners. A five-year	
	follow-up study. Br J Ind Med 19: 65-76, 1962.	461
	HIGGINS IT T, OLDHAM PD, COCHRANE A L and GILSON J C: Respiratory  symptoms and pulmonary disability in an industrial town. Survey of a random	462
	sample of the population. Br Med J 2: 904-9, 1956.	462
	HOLLAND W W and ELLIOTT A: Cigarette smoking, respiratory symptoms, and anti-smoking propaganda. Lancet 1: 41-43, 1968.	463
The Control of the Co	HOLLAND W W, HALIL T, BENNETT A E and ELLIOTT A: Factors influencing the onset of chronic respiratory disease. Br Med J 2: 205-8, 1969A.	464
	HOLLAND W W, HALIL T, BENNETT A E and ELLIOTT A: Indications for	
	measures to be taken in childhood to prevent chronic respiratory disease.  Millbank Mem Fund Quart 47: 215-27, 1969B.	465
	HOLLAND W W, HALIL T and ELLIOTT A: The effect of environmental factors on ventilatory function in schoolchildren.  Pub Health Serv Pub	- 466
	1879: 259-72, 1969C.	
9	HOLLAND W W and REID D D: The urban factors in chronic bronchitis.  Lancet 1: 445-48, 1965.	467
	HOLLAND W W, REID D D, SELTSER R and STONE R W: Respiratory disease in England and the United States. Arch Environ Health 10: 338-48, 1965.	468
	HOWARD P: A long-term follow-up of respiratory symptoms and ventilatory function in a group of working men. Br Ji nd Med 27: 326-333, 1970.	469
	LAMBERT P M and REID D D: Smoking, air pollution, and bronchitis in Britain. Lancet 1: 853-57, 1970.	470
•	LEIBESCHUETZ H J: Respiratory signs and symptoms in young soldiers and their relationship to smoking. J Royal Army Med Corps 105: 76-81, 1959.	471
eri erika erikilari Gundari	LOWE CR, CAMPBELL H, and KHOSLA T: Bronchitis in two integrated steel works. III. Respiratory symptoms and ventilatory capacity related to atmospheric pollution. Br J Ind Med 27: 121-29, 1970.	472
- 1	LOWE CR and KHOSLA T: Chronic bronchitis in-ex-coal miners working in the steel industry. Br J Ind Med 29: 45-49, 1972.	473
	LOWE CR, PELMEAR PL, CAMPBELL H, HITCHENS RAN, KHOSLA T and KING TC: Bronchitis in two integrated steel works. I. Ventilatory capacity age and physique of non-bronchitic men. Br J Prev Soc Med 22: 1-11, 1968.	474
	LYONS J P, RYDER R, CAMPBELL H and GOUGH J: Pulmonary disability in coal workers' pneumoconiosis. Br Med J 1: 713-16, 1972.	475
	McCALLUM R I: Respiratory disease in foundrymen. Br J Ind Med 29: 341-44,	476

Diditography, e. -C

### V-D. MORBIDITY STATISTICS IN OTHER FOREIGN COUNTRIES

The prevalence of respiratory symptoms has been investigated in several countries. As in the United States and Great Britain, there are several causes of respiratory symptoms including cigarette smoking, air pollution and industrial hazards.

- 1. Canada. As in the United States, there is an increasing mortality from bronchitis and emphysema in Canada (Gauthier, 1970; The morbidity statistics are less extensive in Neri et al., 1970). Canada. Anderson and others (1965) surveyed 246 males and 311 females in Chilliwack, British Columbia. Chronic respiratory disease was found to affect 29.3% of men and 18% of women between 25 and 79 years of age. The authors concluded that cigarette smoking was the most important single factor associated with respiratory diseases. Smoking caused a higher prevalence of respiratory symptoms in the following groups: 200 men and women 70 to 89 years old (Aguzzi et al., 1966); 310 male physicians 25 to 74 years old (Lefcoe and Wonnacott, 1970); 1,015 male employees in asbestos mines and mills (McDonald et al., 1972; and 1,138 Eskimos (Beaudry, 1968). The respective roles of air pollution, respiratory infections and social status have not been elucidated in these reports.
- 2. Sweden. The twin registry in Sweden has provided an unusual opportunity to investigate the morbidity among monozygotic and dizygotic pairs. In a group of 71 monozygotic twins with discordant smoking habits, there were more complaints of coughing and bronchitis.

The following surveys indicate a higher incidence of respiratory illness in smokers than in nonsmokers: 339 men 50 years old (Wilhelmsen and Tibblin, 1966; Wilhelmsen et al., 1969), 240 iron mine workers (Jörgensen and Svensson, 1970), and 22,250 urban and rural residents (Irnell and Kiviloog, 1968). In the last two reports, the extent of air pollution influenced the morbidity from cigarette smoking.

3. Finland. In a rural population and among area pulp mill worke there is a difference in prevalence of respiratory symptoms both in smokers and in nonsmokers. The higher incidence in the industrial area may relate either to the pollution in the air or to the fact that it is in the Arctic area (Huhti, 1965, 1966; Huhti et al., 1970). In these surveys as well as in those conducted by Järvinen et al. (1966) and Ruikka et al. (1966), smoking was found to increase the

- 4. France. The higher prevalence of chronic bronchitis among smokers as compared with nonsmokers has been reported by Fréour et al. (1966), Kourilsky et al. (1966), Coudray et al. (1969), Brille (1969, 1970), Brille et al., 1970, Fournier and Zivy (1970), Golli (1970), Jancik (1970), and Vigy (1970). All of these reports confirm the observations previously made in the United States and Great Britain.
- 5. Germany. More than 30,000 questionnaires from 10 clinics of different towns in West Germany were analyzed (Ulmer and Reif, 1966). The nature of occupation influences the incidence of obstructive emphysema. Specific occupational types have been surveyed by others: asbestos workers (Hany et al., 1967), coal miners (Ulmer et al., 1968; Ulmer and Reichel, 1970; Ulmer, 1970) and chemical factory workers (Possner, 1970). The role of industrial exposure and cigarette smoking in the pathogenesis of pulmonary emphysema in Germany has been discussed by Wendel (1966).
- 6. Other Western European countries. In the Netherlands,
  Biersteker (1968, 1969) and Biersteker et al. (1969) reported an
  association between smoking and symptoms of chronic bronchitis
  among 1,000 male municipal employees. In Denmark, a group of 802
  males and females over 50 years of age showed a lower ventilatory
  function in smokers than in nonsmokers (Hagerup and Larsen, 1971).
  In Copenhagen, a group of 156 welders was compared with a group of
  152 other workers from the same plant (Fogh et al., 1969). The

Bibliography V-D	Reprint
PREGOWSKI W, BRYSIEWICZ, CESARZ-FRONCZYK M, KORDECKA J, LESZCBINSKI B, LUKJAN Z and SADOKIERSKA H: La bronchite chronique parmi les ouvriers d'une usine textile. Dépistage clinique et	
radiologique. Rev Tubere Pneum 34: 753-5, 1970.	569
PRINSLOO I and LAUBSCHER N F: Emphysema in South African gold miners.  In: Pneumoconiosis, edited by H A Shapiro, Cape Town, Oxford University Press, London, 380-5, 1970.	570
PUDELSKI J, OKLEK K and DELOFF L: Obraz epidemiologiczny przewlekłych nieswoistych chorób płuc u mieszkańców miasta zabrza. (Epidemiological	
situation of chronic non-specific diseases of the lung in inhabitants of the town of Zabrze.) Gruzlica 39: 604-9, 1971.	571
RODRÍGUEZ R L, ACHÓN P M, RODRÍGUEZ S H, DE LA PAZ E, PITA DE LA VEGA H, DURAND O R and SOLLET G R: Hábito de fumar y diagnóstico al alta. Estudio de 453 casos. Rev Cuba Med 8: 559-67, 1969.	572
RUIKKA I. SOURANDER L B and KASANEN A: The health of the aged in Turku.  Ann Acad Sci Fenn Series A, 120: 1-50, 1966.	<b>57</b> 3
SLUIS-CREMER G K and SICHEL H S: Ventilatory function in males in a Witwatersrand town. Am Rev Resp Dis 98: 229-39, 1968.	574
SLUIS-CREMER GK, WALTERS LG and SICHEL HS: Chronic bronchitis in miners and non-miners: an epidemiological survey of a community in the gold-mining area in the Transvaal. Brit J Indust Med 24: 1-12, 1967a.	575
SLUIS-CREMER GK, WALTERS LG and SICHEL HS: Ventilatory function in relation to mining experience and smoking in a random sample of miners and non-miners in a Witwatersrand town. Brit J Industr Med 24: 13-25,	576
STANEK V, FODOR J, HEJL Z, WIDIMSKY J, CHARVAT P, SANTRUCEK M,	
ZAJIC F and VAVRIK M: A contribution to the epidemiology of chronic bronchitis. Acta Med Scand 179: 737-46, 1966.	577
STANESCU D.C., PILAT L., GAVRILESCU N., TECULESCU D.B and CRISTESCU I.  Aspects of pulmonary mechanics in arc welders' siderosis. Brit J Industr  Med 24: 143-7, 1967.	i: 578
STROBEL M and GSELL O: Mortalität in Beziehung zum Tabakrauchen 9 Jahre Beobachtungen bei Arzten in der Schweiz. Helv Med Acta 32: 547-92, 1965.	579
SZYMCZYKIEWICZ K E, KUNSKI H and GIELEC L: Clinical evaluation of the respiratory system in cotton workers. Pol Med Sci Hist 13: 110-4, 1970.	580
TAKENOUCHI S: Epidemiological studies on chronic respiratory disturbances  among employees in a certain organization's workshop. J Nara Med Assoc  19: 749-63, 1968.	581
TSUMETOSHI Y, SHIMIZU T, TAKAHASHI H, ICHINOSAWA A, UEDA M, NAKAYAMA H, YAMAGATA Y and OHSHINO A: Epidemiological study of chronic bronchitis with special reference to effect of air pollution. International Arch Arbeitsmedizin 29: 1-27, 1971.	582
ULMER W T: Emphysema of the lung: clinical and experimental investigations of its pathogenesis. Geriatrics 25: 164-9, 1970.	583

		u,
	Dionography V-D	
		į,
	ULMER W T and REICHEL G: Zur Epidemiologie der chronischen bronchitis und	经
	deren Zusammenhang mit der Luftverschmutzung. (The epidemiology of chronic bronchitis and its relationship to air pollution.) Deutsche Med	
		84
	<u> </u>	
<b>A</b>	ULMER W T, REICHEL G and WERNER U: Die chronisch obstruktive bronchitis	
	des Bergmannes. Untersuchungen zur Häufigkeit bei der Normalbevölkerung und bei Bergleuten. Die Bedeutung der Staubbelastung und der Einflub des	1
	Rauchens. (Chronic obstructive bronchitis of the coal miner An	
	epidemiologic study of its incidence in the normal population and in coal	
	miners. The importance of dust and smoking. Int Arch Gewerbepath  Gewerbehyg 25: 75-98, 1968.	85
	ULMER W T and REIF E: Epidemiologische Untersuchungen zur klinischen	77
¥1.012-33	Bedeutung des chronisch obstruktiven Lungenemphysems. (Epidemiologic study of the clinical importance of chronic obstructive emphysema.)	- 14
		86
	VALIC F and ZUSKIN E: Byssinosis: a follow-up study of workers exposed to fine grade cotton dust. Thorax 27: 459-63, 1972.	87
		227. 1. 7.
	VIGY M: Tabac, bronchite chronique et emphyseme. Concours Med 3: 31-7, 1970.	88
	WYSYOCII I. Watah ara saurich nodminak kanamanani hyanchitiday a amfuramen	
	VYSKOCIL J: Vztah pracovnich podminek k onemocneni bronchiticou a emfyzemem plic. (Relationship between working-conditions, bronchiticodisease and	5 4 E
	lung-emphysema.) Prac Lek 20: 41-7, 1968.	89
	MATCHE MARGINET A. Designatory comptons and empling babits in	
and the second	WALSHE M M and HAYES J A: Respiratory symptoms and smoking habits in  Jamaica. Am Rev Resp Dis 96: 640-4, 1967.  5	91
) ·	WENDEL H: Beziehungen zwischen Nikotinabusus und Lungenemphysem.	9
	Z Gesamte Inn Med 21: 777-80, 1966.	
	WILHELMSEN L, ORHA I and TIBBLIN G: Decrease in ventilatory capacity	
	between ages of 50 and 54 in representative sample of Swedish men.	9
	Brit Med J 3: 553-6, 1969.	
	WILHELMSEN L and TIBBLIN G: Tobacco smoking in fifty-year-old men.	
•	1. Respiratory symptoms and ventilatory function tests. Scand J Resp	9
	Dis. 47: 121-30, 1966.	7
	WOOLCOCK A J, BLACKBURN C R, FREEMAN M H, ZYLSTRA W and	
	SPRING S R: Studies of chronic (nontuberculous) lung disease in New Guinea populations. Am Rev Resp Dis 102: 575-90, 1970.	9
	Guinea populations. Am Nev Ness Bis 102. 515-70, 1770.	
	ZWI S, SLUIS-CREMER GK and DU PREEZ L: A survey of pulmonary function	4
	in male office workers. Med Proceedings 13: 569-74, 1967.	9
		47 47
		, p.,
•		

VI: BRONCHOPULMONARY FUNCTION IN SMOKERS AND NONSMOKERS

There are several investigations comparing measurements of bronchopulmonary function in smokers and nonsmokers. The results are discussed in this section, as distinct from those contrasting patients having pulmonary emphysema or chronic bronchitis with healthy controls (see Sections II and III). The separation is not a common practice because of the widespread belief that in smokers there is an abnormal bronchopulmonary condition representing the early stages of chronic bronchitis or pulmonary emphysema. There is no clinical or experimental support for this assumption.

With regard to the literature on this topic, it is important to

point out that in the follow-up of smokers who show subnormal

bronchopulmonary functioning there is no report of a subject who

has developed chronic bronchitis and pulmonary emphysema. Most

of the available articles deal with a comparison of smokers and nonsmokers

executed once without a follow-up. The bibliographic list containing

articles which serve as a background to this section is as follows:

No. 13. Pulmonary functional test.

Several reports have appeared describing the effects of cigarette smoking on lung function. The conclusions are varied and depend on the manner of comparing smokers with nonsmokers. This discussion considers the importance of age, sex, height, environment and country of origin in determining the influence of cigarette smoking on pulmonary ventilation.

l. Age, sex, height and environment. A group of 302 subjects in Australia was investigated by Read and Selby (1961). The results were expressed as regression coefficients. for mean expiratory flow based on age and height. Smokers without symptoms did not differ significantly from nonsmokers. However, smokers who complained of coughing with or without expectoration had lower ventilatory function than nonsmokers. The authors concluded that smoking alone does not reduce ventilatory function and suggested that a combination of genetic and environmental factors would produce symptoms and signs of decreased ventilatory function.

Cotes et al. (1966) have calculated the ventilatory capacities standardized to the average age, height and weight. They found the regression coefficient calculated separately for nonsmokers, light-to-moderate smokers and heavy smokers to be practically the same. This manner of testing the significance of the difference between smokers and nonsmokers has been overlooked by others who have correlated ventilatory capacities with age only.

in the United States, specifying their age, height and smoking habits.

They noted that vital capacity, inspiratory capacity and forced expiratory rate were slightly smaller and the functional residual capacity and residual volume were slightly lower in smokers than they were in nonsmokers. The differences were explained by the difference in mean age of nonsmoking subjects, which was lower than that of any of those who smoked.

In Finland, 420 men and 608 women, ranging in age from 40 to 64 years, were examined by Huhti (1967). On the basis of regression coefficients of age and height, there was no difference in forced vital capacity, one-second forced expiratory volume and peak expiratory flow between smokers and nonsmokers. A group of 44 young male recruits in Roumania was investigated by Stanescu et al. (1968).

Lung volumes and ventilatory capacities of smokers did not differ from those of nonsmokers.

2. Age and environment. Flick and Paton (1959) reported the first comparison of ventilatory tests in a group of 222 male patients at a Veterans Hospital in the United States. The maximal expiratory flow was statistically different only between 60 and 70 years of age: the mean value for 20 nonsmokers was 378 l/min and that for 61 smokers was 258 l/min. Subjects from 20 to 50 years did not show a significant difference in their maximal expiratory flow.

by other investigators to reduce ventilatory function: Franklin

and Lowell (1961) in 376 male employees in Massachusetts; Catlett and Kidera (1969) in 257 male flight officers. Larson (1963) noted a difference in ventilatory function between smokers and nonsmokers, starting with the 30-year-old group in California. However, the difference cannot be attributed to smoking alone. The role of air pollution is difficult to assess in these reports, but one additional report has defined its influence. Anderson and Ferris (1965) compared the results of pulmonary function tests in a group in New Hampshire and another group in Canada. After controlling variations in age, sex and smoking habits, there were still significant differences in values of forced expiratory volume and peak expiratory flow rate. These differences could be explained by ethnic factors and atmospheric pollution.

physicians, ranging in age from 25 to 79 years, showed that smokers have lower ventilatory function than nonsmokers (McIlreath and Cohen, 1966). In another group of 410 volunteers in an American community, ranging in age from 20 to 103 years, there was a decrease in ventilatory function of smokers (Edelman et al., 1966). A similar conclusion was arrived at in the examination of the following: 350 males in Pennsylvania 50 years or over (Weiss et al., 1963), 20 medical students or doctors in Massachusetts 18 to 45 years (Zwi et al., 1964), 140 males and females in Connecticut 67 to 95 years (Kiss, 1966), 150 males in India 15 to 50 years of age (Mohanty and Gupta, 1968), 298 females in Canada 25 to 59 years of age (Woolf and Suero, 1971), and 365 students in Connecticut

Source: https://www.industrydocuments.ucsf.edu/docs/pfmk0000

occurrence of respiratory symptoms in the welders and in the controls

was associated with the cigarette smoking.

In Norway, Haenszel and Hougen (1972) concluded that the prevalence of respiratory symptoms was related to urban residence and to the amount of cigarettes consumed. In Spain, Bouhuys et al. (1969 a, b) noted that in a group of 216 hemp workers the associated disabling respiratory disease is attributable not to smoking habits but to prolonged exposure to hemp dust. In Switzerland, prospective studies of 3,479 doctors (Strobel and Gisell, 1965) and 1,885 residents of Basle (Mannhart, 1962) revealed a higher incidence of chronic lung disease among smokers than among nonsmokers.

7. Eastern Europe. In Poland, most surveys were conducted among workers with the following industrial connections: port and shipyard (Dobrzyński et al., 1970a), grain elevator (Dobrzyński et al., 1970b), wool industry (Brysiewicz et al., 1970), textile (Cierniak et al., 1970;, Szymczykiewicz et al., 1970), plaster works (Owsinski et al., 1971) and steel works (Council of Scientific Research, 1972). The workers who smoked cigarettes showed a higher rate of incidence of respiratory symptoms than nonsmokers. There are surveys which include residents of Bialystok (Pregowski et al., 1970), Warsaw and (Kucewicz, 1969 a, b), Lublin (Durda and Szafranski, 1971),/Zarze (Pudelski et al., 1971).

In Czechoslovakia, epidemiological studies have been conducted as follows: 473 men 60 to 64 years old in Prague (Stanek et al., 1966);

441 men 50 to 65 years old in Prague (Boudik et al., 1970); 112 min ers in Karvina (Pochmon et al., 1968); 3,466 hospital patients in Katowicach (Gasinska and Gburek, 1970); 2,376 persons in Brno (Vyskocil, 1968; Jancik and Jancik-Mak, 1972); and 95 men residing in Klin (Feuereisl et al., 1972). In the last-mentioned study, 5 men out of 51 who smoked 292,000 to 584,000 cigarettes during their lifetime showed no objective signs of chronic bronchitis.

The surveys in the U.S.S.R. (Petrova, 1956; Danovich et al. 1969; Futorainsky et al., 1971), Roumania (Jelea et al., 1964; Stanescu et al., 1967), and Yugoslavia (Kalacić, 1970; Valic and Zuskin, 1972) have revealed an association between smoking and symptoms of chronic lung disease.

8. Middle East and Africa. Workers in South African gold mines have been examined by several investigators. Chatgidakis (1960) noted at autopsy that smokers with silicosis had lungs characterized by enlarged bronchial mucus glands. The enlargement was also seen in nonsmokers. In a survey of 827 males, chronic bronchitis was more common in miners than in nonminers in the same smoking category (Zwi et al., 1967; Sluis-Cremer et al., 1967 a, b; Sluis-Cremer and Sichel, 1968). The severity of emphysema in gold miners detected at autopsy was not related to smoking habits (Prinsloo and Laubscher,

1970).

In Egypt, a group of 223 cement workers showed a significant correlation between cement exposure and wheezing, dyspnea and physical signs of chronic bronchitis. Cigarette smoking was related only to cough and expectoration (El-Sewefy and Awad, 1971). Among 89 bricklayers and 245 workers in the rolling mills, respiratory symptoms were more frequently encountered among smokers than among nonsmokers (El-Sewefy, 1970). In Israel, among 257 cotton mill workers and 64 orchestra players, smoking was a significant factor which increased the prevalence of respiratory symptoms (Chwat and Mordish, 1971). Results of a survey of 175 Indian wool workers are similar (Mathur and Misra, 1972).

9. Asian-Pacific countries. In Australia, the following types of industrial workers have been examined: coalminers (Outhred and Flynn, 1960; McKenzie et al., 1969); felt manufacturers (Gandevia and Milne, 1965), cotton workers (Barnes and Simpson, 1968).

Residents in New South Wales (Hong et al., 1967; Gandeva, 1969) and in Sydney (Lake, 1969) have also been surveyed. An association between cigarette smoking and prevalence of chronic lung disease is indicated.

The results of surveys conducted in Japan (Takenouchi, 1968;

Nishimoto et al., 1970; Tsumetoshi et al., 1971), in Singapore (Da Costa, 1972), in New Zealand (De Hamel et al., 1972) and in New Guinea

(Woolcock et al., 1970) have revealed a relationship between smoking

Source: https://www.industrydocuments.uesf.edu/docs/pfmk000/

10. South America and Caribbean. The epidemiological
1971;
studies, reported from Mexico (Paras Chavero et al., 1970/Celis
1968),
et al./. Cuba (Rodriguez Rivera et al., 1969), Chile (Oyanguren et al.,
1972) and Jamaica (Walshe and Hayes, 1967) reveal an association
between smoking and prevalence of respiratory symptoms.

V. NON-OCCURRENCE OF CHRONIC PULMONARY DISEASE IN SMOKERS

D. MORBIDITY STATISTICS IN OTHER FOREIGN COUNTRIES

· [발생] (10] [발생] [발생] [발생] [발생] [발생] [발생] [발생] [발생	Reprin
AGUZZI G, WOOLF C R and PATERSON J F: The prevalence and type of chronic obstructive bronchopulmonary disease in very old people. Canad Med Ass J	401
94: 932-9, 1966.	491
ANDERSON DO, FERRIS JR B C and ZICKMANTEL R: The Chilliwack respirator survey, 1963: Part IV. The effect of tobacco smoking on the prevalence of respiratory disease. Canad Med Ass J 92: 1066-76, 1965.	y 492
BARNES R and SIMPSON G R: Ventilatory capacity changes on exposure to cotton dust. Med J Australia 1: 897-900, 1968.	493
BEAUDRY PH: Pulmonary function survey of the Canadian Eastern Arctic Eskimo.  Arch Environ Health 17: 524-8, 1968.	494
BIERSTEKER K: Bronchitisklachten bij Rotterdams gemeentepersoneel. (Symptoms of bronchitis among municipal personnel in Rotterdam.) Ned T Geneesk 112: 1208-11, 1968.	495
BIERSTEKER K: Air pollution and smoking as cause of bronchitis. Arch Environ Health 18: 531-5, 1969.	496
BIERSTEKER K, VAN GEUNS H. A and VAN LEEUWEN P: Cough and peak flow rates of municipal employees in Rotterdam. Environ Res 2: 272-6, 1969.	497
BOUDIK F, GOLDSMITH JR, TEICHMAN V and KAUFMANN P-C: Epidemiology of chronic bronchitis in Prague. Bull WHO 42: 711-22, 1970.	<b>4</b> 98
BOUHUYS A, BARBERO A, SCHILLING R S and VAN DE WOESTIJNE K P:  Chronic respiratory disease in hemp workers. Am J Med 46: 526-37, 1969a.	499
BOUHUYS A, SCHILLING R S and VAN DE WOESTIJNE K P: Cigarette smoking, occupational dust exposure, and ventilatory capacity. Arch Environ Health 19: 793-7, 1969	500
BRILLE D: Fréquence de la bronchite chronique chez la femme rôle du tabac.  Rev Tuberc et Pneumol 33: 794-6, 1969.	501
BRILLE D: Possibilites du depistage au stade initial de la bronchite chronique et de l'emphysème pulmonaire. (Possibilities of detection in the early stages of chronic bronchitis and pulmonary emphysema.) Rev Med 11: 2201-10, 1970.	502
BRILLE D, COUDRAY P, ISRAEL-ASSELAIN R, LE MELLETIER J, MICHEL F : MIGUERES J, PHAM G T and VOISIN C: Colloques sur la bronchite chroni I Nosologie et épidémiologie. Rev Tuberc Pneumol 34: 723-48, 1970.	
BRYSIEWICZ K, BULUK H, CESARZ-FRONCZYK M, KORDECKA J, LESZCZYNSK B, ŁUKJAN Z and SADOKIERSKA H: Wpływ Pracy W Warunkach Zapylenia Na Częstość Występowania Przewlekłych Zapaleń Oskrzeli U Pracowników Zakładów Przemysłu Wełnianego Im. Sierżana W	Í

Biarymstoku. (The effect of occupation in dustiness on the prevalence of chronic bronchitis among the workmen of the Sierzan's establishments

of wool industry at Białystok.) Gruzlica 38: 657-61, 1970.

_		-يانورسىد
<u> </u>	Bibliography V-D	Reprint
,	CEDERLÖF R: Urban factor and prevalence of respiratory symptoms and "angina pectoris." Arch Environ Health 13: 743-8, 1966.	<b>5</b> 05
•	CEDERLÖF R, EDFORS M-L, FRIBERG L and JONSSON E: Hereditary factors, "spontaneous cough" and "smoker's cough". Arch Environ Health 14: 401-6, 1967.	506
	CEDERLOF R, FRIBERG L and HRUBEC Z: Cordiovascular and respiratory symptoms in relation to tobacco smoking. A study on American twins.  Arch Environ Health 18: 934-40, 1969.	507
24	CEDERLOF R, FRIBERG L, JONSSON E and KAIJ L: Morbidity among monozygotic twins. Arch Environ Health 10: 346-50, 1965.	508
	CEDERLOF R, FRIBERG L, JONSSON E and KAIJ L: Respiratory symptoms and "angina pectori" in twins with reference to smoking habits. An epidemiological study with mailed questionnaire. Arch Environ Health 13: 726-37, 1966.	509
(	CELIS S A, DÍAZ O J, GOROCICA D and CANO V F: Enfisema pulmonar y contaminacion aerea. Neumol Cir Torax 29: 135-41, 1968.	510
a:	CHATGIDAKIS CB: A study of the bronchial mucous glands in white South  African gold miners. Arch Environ Health 1: 335-42, 1960.	511
	CHWAT M. and MORDISH R: Respiratory symptoms and byssinosis among textile workers in Israel. Harefual 80: 559-62, 586-7, 1971.	512
	CIERNIAK E, JELONKIEWICZ J, SŁODCZYK M, CIERNIAK J and PAWLAK F: Przewlekłe Zapalenie Oskrzeli Wśród Pracowników Zakładów Włókienniczych. (Chronic bronchitis in the workmen of textile industry.) Gruzlica 38: 635-41, 1970.	513
	COUDRAY P, SERISE M and FRÉOUR P: Recherche épidémiologique sur les bronchites chroniques et les insuffisances respiratoires. Étude de prévalence dans un échantillon de femmes de 30 a 70 ans. Rev Tuberc Pneumol 33: 769-80, 1969.	514
	COUNCIL O F SCIENTIFIC RESEARCH ON CHRONIC NONSPECIFIC RESPIRATOR DISEASES IN CRACOW, POLAND: Przewlekle Nieswoiste Choroby Układ Oddechowego Wsrod Mieszakancow Krakowa. XVI. Wyniki Badania Przekrojowego w Hucie im Lenina. (Chronic nonspecific respiratory diseases in the inhabitants of Cracow, XVI. Results of a cross section study in the Lenin steel works.) Przeglad Epidemiologiczny 26: 125-34, 1972.	
1	DA COSTA J L: Chronic obstructive lung disease (COLD): A comparison between men and women. Singapore Med J 13: 74-8, 1972.	
1	DANOVICH B Z, KOROVINA O V, SMIRNOV I P and ANISIMOVA N A: On the epidemiology of chronic nonspecific diseases of the lungs. Teraper Ark 41: 63-8, 1969.	hiv 517
1	DEHAMEL F A and O'DONNELL T V: Smoking habits and respiratory symptoms of Dunedin public servants. New Zealand Med J. 76: 1-4, 1972;	518
1	DOBRZYŃSKI W, KISIELEWICZ J and KOCIECKA I: Analiza Kliniczno- Statystyczna Przewlekłych Nieżytow Oskrzeli U Pracowników Portu I Stoczni W Szczecinie. (Clinical and statistical analysis of chronic bronchitis in port and shipyard workers.) Medycyna Pracy 21: 294-	
	106 1970-	519

The state of the s	
Bibliography V-D	leprint
DOBRZYŃSKI W, KISIELEWICZ Jand KOCIECKA I: Przewiekłe Zapalenia Oskrzeli U Pracowników Portowego Elewatora Zhożowego. (Chronic bronchitis among the workmen of a harbour grain elevator.) Gruzlica 38: 651-6, 1970b.	520
DURDA M and SZAFRANSKI W: Przewlekłe nieswoiste choroby płuc wśród ludności wiejskiej woj. lubelskiego. (Chronic non-specific diseases of the lung among the rural population of the Lublin Voivodship.) Gruzlica 39: 616-20, 1971.	521
EL-SEWEFY A Z: Chest symptomatology in a Sheffield steel works (S. P. T.).  J Egyp Med Assoc 52: 578-81, 1970.	522
EL-SEWEFY A Z and AWAD S: Chest symptomatology in an Egyptian Portland cement factory. J Egyp Med Assoc 54: 457-64, 1971.	523
FEUEREISL R, KRZHIVANEK I, REIL I, KUBIK A and FELKEL K: Effect of smoking on the cardiopulmonary function. Klinicheskaya Meditzina 50: 82-4, 1972.	524
FOGH A, FROST J and GEORG J: Respiratory symptoms and pulmonary function in welders. Ann Occup Hyg 12: 213-8, 1969.	525
FOURNIER E and ZIVY P: Poumon et tabac. (Lung and tobacco). Poumon Coeur 26: 1109-25, 1970.	526
FREOUR P, COUDRAY P, ROUSSEL A and SERISE A: Les bronchites chroniques et l'insuffisance respiratoire dans l'agglomération de Bordeaux. J Med Bordeaux 143: 1865-79, 1966.	527
FUTORYANSKY YI, DETSIK YI, SHUFLAT AN, SIMONCHIK V N and BARANSKY S M: Effect of smoking on the ventilation function of the lungs. Klinicheskava Meditsina 49: 70-4, 1971.	528
GANDEVIA B: A productive cough upon request as an index of chronic bronchitis: the effects of age, sex, smoking habit and environment upon prevalence in Australian general practice. Med J Australia 1: 16-20, 1969.	<b>52</b> 9
GANDEVIA B and MILNE J: Ventilatory capacity on exposure to jute dust and the relevance of productive cough and smoking to the response. Brit J Industr Med 22: 187-95, 1965.	530
GAUTHIER H: L'insuffisance respiratoire au Canada. Laval Med 41: 992-4, 1970.	531
GASINSKA T and GBUREK Z: O Szkodliwości Przewlekłego Palenia Tytoniu.  (On the harmful effects of chronic tobacco smoking.) Wiad Lek 23: 2157-62, 1970.	532
GOLLI V: Étude épidémiologique des bronchopathies non spécifiques. Bronches 20: 313-30, 1970.	. 533
HAENSZEL W and HOUGEN A: Prevalence of respiratory symptoms in Norway.  J Chronic Dis 25: 519-44, 1972.	534
HAGERUP L and LARSEN M: Tobaksrygning og respiratoriske symptomer i en dansk population. (Tobacco smoking and respiratory symptoms in a Danish population.) Ugeskr Laeg 133: 1302-6, 1971.	<b>5</b> 35
HANY A, BURCKHARDT P and BUHLMANN A: Zur Klinik und Pathophysiologie  der Lungenasbestose. Schweiz Med Wochenschr 97: 597-603, 1967.	536

	*.
HONG CS, GANDEVIA B and LOVELL H: Ventilatory capacity in a series of male	
adults and the effect of respiratory symptoms, productive cough, smoking	
habit and occupation. Med J Australia 1: 169-72, 1967.	537
UHUTI E. Danielana of manimatory symptoms about heart has the	
HUHTI E: Prevalence of respiratory symptoms, chronic bronchitis and pulmonary emphysema in a Finnish rural population. Field survey of age group 40-64	
in the Harjavalta area. Acta Tuberc Pneumol Scand Suppl 61: 101 pp., 1965.	538
美麗報報 하는 사람들은 사람들은 사람들이 가장 하는 사람들이 되었다.	
HUHTI E: Chronic respiratory disease among old people in a Finnish rural area.	
Ann Med Int Fenn 55: 99-105, 1966.	539
HINTE E BUILANEN D WINDALA Hand MAKKINITH I Chang	St. Inc. of the
HUHTI E, RYHANEN P, VUOPALA U and TAKKUNEN J: Chronic respiratory disease among pulp mill workers in an arctic area in Northern Finland.	
Acta Med Scand 187: 433-44, 1970.	540
INTERNATIONAL SYMPOSIUM IN SAN JUAN: Twin registries in the study of	
chronic disease. With particular reference to the relation of smoking	
to cardiovascular and pulmonary diseases. Acta Med Scand Suppl. 523:	541
	•
IRNELL L and KIVILOOG J: Bronchial asthma and chronic bronchitis in a	
Swedish urban and rural population. With special reference to prevalence,	
respiratory function and socio-medical condition. Scand J Resp Dis (Suppl	
66: 86pp., 1968.	542
	* .
JANCIK E: Quelques données et corrélations provenant d'une étude épidémiologique sur l'incidence des symptômes caractéristiques des maladies respiratoires	
chroniques non spécifiques. Rev Tuberc Pneum 34: 749-52, 1970.	543
	Carrier Contract
JANCIK E and JANCIK-MAK M: Chronische unspezifische lungenerkrankungen.	
(Chronic non-specific lung diseases). Praxis Pneumologie 26: 69-81, 1972.	544
	.'
JARVINEN KA, PATIALA J and THOMANDER K: The role of chronic smoking	
in the etiology of obstructive pulmonary emphysema. A study of 3, 375 persons. Ann Med Intern Fenn 49: 307-311, 1966.	545
	343
JELEA A, ILIE E and GAVRILA F: Contribuții la studiul insuficienței pulmonare	
de etiologie tabagică. (Contributions to the study of pulmonary insufficiency	
of tabacic etiology.) Medicina Interna 16: 229-34, 1964.	. 546
JÖRGENSEN H and SVENSSON A: Studies on pulmonary function and respiratory	
tract symptoms of workers in an iron ore mine where diesel trucks are used	Į
underground. J Occup Med 12: 348-54, 1970.	547
KALAČIĆ I: Kronicni bronhitis i ventilacijska funkcija pluća u pušača i nepušača.	
(Chronic bronchitis and ventilatory lung function in smokers and nonsmokers	
Arh Hig Rada 21: 5-12, 1970.	548
KOURILSKY R, BRILLE D and HATTE M J: Etude statistique de la relation entre	
le tabac et la bronchite chronique. Bull Acad Nat Med 150: 318-29, 1966.	549
	•
KUCEWICZ J: Przewlekie Nieswoiste Choroby Układu Oddechowego Wśród	
· Mieszkańcow Krakowa, VIII. Metoda Dyskryminacyjna Fishera W Analizie	
Wprywu Palenia Tytoniu na Występowanie Objawów Chorobowych. (Chronic	
nonspecific respiratory diseases in the city of Cracow.) Przeg Epid 23:	550
<b>331-7</b> , 1969a.	250
KUCEWICZ J: Chronic nonspecific respiratory diseases in the city of Cracow.	
Epidemiol Rev 23: 159-65, 1969b.	551
	•
The second secon	
	<b>!</b> ~
"我没有是我多数,还是不要会的。"  "我们,我们也没有一种的特别。  "我们就是这个一个,我们还是一个一个一个,我们还是一个一个一个,我们就是这样的,我们就是	The second second

Bibliography V-D page	ll5 Repris	nt )
LAKE B: Morbid conditions at death in old men. J Chronic Dis 21: 761-79, 19	69. 552	
LEFCOE N M and WONNACOTT TH: The prevalence of chronic respiratory in the male physicians of London, Ontario. Canad Med Assoc J 102: 3		
LUNDMAN T: Smoking in relation to coronary heart disease and lung function  Acta Med Scand 180 (Suppl. 455): 1-75, 1966.	in twins. 554	
MANNHART M: Krankheiten in ihrer Beziehung zu Genubmitteln in der Bevolk von Basel. Enquête bei erwachsenen Personen 1956 und 1961. Zeitsch fur klinische Medizin 157: 240-57, 1962.		
MATHUR K C and MISRA S N: Incidence of pulmonary diseases among wool workers. Indian J Chest Dis 14: 172-8, 1972.	<b>5</b> 56	
MCDONALD J C, BECKLAKE M R, FOURNIER-MASSEY G and ROSSITER of Respiratory symptoms in chrysotile asbestos mine and mill workers of Quebec. Arch Environ Health 24: 358-63, 1972.		
MCKENZIE HI, GLICK Mand OUTHRED KG: Chronic bronchitis in coal miners: ante-mortem/post-mortem comparisons. Thorax 24: 527-35-1969.	5, 558	
NERI L C, MANDEL J S, O'CONNOR M L and LARSON M: A household sur of chronic obstructive pulmonary disease in Ottawa: Methodological problems and results of pilot survey. Canad J Pub Health 61: 407-12,		
NISHIMOTO Y, BURROWS B, MIYANISHI M, KATSUTA S, SHIGENOBU T a KETTEL L J: Chronic obstructive lung disease in Japanese poison ga workers. Am Rev Resp Dis 102: 173-9, 1970.		
OUTHRED K G and FLYNN M J: Some aspects of chronic respiratory disease coalminers in New South Wales, Australia. Dis Chest 37: 390-9, 1960		
OWSINSKI J, JANUS T and MYSIK M: Występowanie przewlekłych nieswoisty chorób układu oddechowego (p. n. ch. u. o.) wśród pracowników Zakładu Przemyslu Gipsowego, Dolina Nidy" w Gackach. (Occurrence of chrunspecific diseases of the respiratory system among workers in the prworks at Dolina Nidy in Gacki.) Przeg Lek 28: 476-80, 1971.	onic	
OYANGUREN H. DONOSO H. PRIETO L. VARGAS Sand PAZ S: Prevalen- la bronquitis cronica en un area urbana y una rural de Santiago y su r- con el grado de contaminacion atmosferica. (Prevalence of chronic b in urban and rural areas of Santiago: Its relation with the degree of a	elacio <del>n</del> ronchitis	
pollution). Rev Mcd Chile 100: 101-5, 1972.  PARAS C E, MERCADO C E and QUINTERO N A: Tabaquismo y Cardiopatia Arch Inst Cardiol Mex 40: 128-34, 1970.		
PARÁS CE, MERCADO E and QUINTERO A: Simposio sobre problemas m originados por el tabaquismo. II. Alteraciones cardiovasculares. Ne Cir Tórax Méx 32: 15-8, 1971.	edicos umol 565	[005]
PETROVA A: Influence of occupational hazards in tobacco fermentation factors Bulgaria (town Ploydiv) on the organism of the workers. 69-81, 19		05051
POCHMON V, SKARABELLOVA M and LEGOSTOV J: Chronic obstructive in miners of Karvine District. Prac Lekar 20: 366-71, 1968.	bronchitis 567	
POSSNER W: Betrachtungen und Ergebnisse zur Epidemiologie nichtspezifisc Erkrankungen der Atemwege. Z Ges Hyg 16: 545-52, 1970.	her	3

of smokers and nonsmokers were not matched, so that it is not possible to specify the age when smoking influences ventilatory function.

The technique of pairing a smoker with a nonsmoker of the same age and sex has been used by two groups of investigators. Wilson et al. (1960) reported 14 pairs in Texas all over 40 years old and noted a decrease in timed expiratory volume, total lung capacity and maximal breathing capacity. Carey et al. (1967) reported 7 pairs in Ireland ranging from 19 to 60 years of age. Ventilatory capacity measured daily was less in smokers than in nonsmokers. The size of the group does not permit the conclusion as to the age when smoking starts to influence ventilatory capacity.

4. Abstinence from smoking. In some smokers with abnormal ventilatory function, cessation of smoking caused an improvement (Peters and Miller, 1960; Wynder et al., 1967; Peterson et al., 1968; Schuman, 1971). Administration of bronchodilator drugs has also reduced the airway obstruction in smokers (Petty et al., 1970). The abnormalities in ventilatory function associated with smoking are reversible.

	page 123
BIBLIOGRAPHY	
VI. BRONCHOPULMONARY FUNCTION IN SMOKERS AND NONSMOKE	ers
A. VENTILATORY FUNCTION	Reprint
ANDERSON DO, and FERRIS BG Jr: Air pollution levels and chronic redisease. Arch Environ Health 10: 307-11, 1965.	spiratory 596
BOREN HG, KORY R C and SYNER J C: The Veterans administration-a cooperative study of pulmonary function. II: The lung volume and i subdivisions in normal men. Am J Med 41: 96-114, 1966.	
CAREY G C R, DAWSON T A J and MERRETT J D: Daily changes in ven capacity in smokers and in non-smokers. Br J Prev Soc Med 21: 86-89, 1967.	tilatory 598
CATLETT G F and KIDERA G J: Detection of respiratory impairment in part Med 40: 1252-57, 1969.	599
COTES JE, ROSSITER CE, HIGGINS ITT and GILSON JC: Average values for the forced expiratory volume in white Caucasian males.	normal
Br Med J 1: 1016-19, 1966.	600
EDELMAN NH, MITTMAN C, NORRIS AH, COHEN BH and SHOCK N  The effects of cigarette smoking upon spirometric performance of community dwelling men. Am Rev Resp Dis 94: 421-29, 1966.	w: 601
FLICK A L and PATON R R: Obstructive emphysema in cigarette smoker.  Arch Intern Med 104: 518-26, 1959.	602
FRANKLIN W and LOWELL F C: Unrecognized airway obstruction associ with smoking: a probable forerunner of obstructive pulmonary em  Ann Intern Mcd 54: 379-86, 1961.	atied physema. 603
HUHTI E: Ventilatory function in healthy nonsmokers and smokers. Scan Resp Dis 48: 149-55, 1967.	<u>d J</u> 604
KISS G T: Pulmonary survey in the aged. Dis Chest 49: 305-8, 1966.	605
LARSON R K: The chronic effect of cigarette smoking on pulmonary venti <u>Am Rev Resp Dis</u> 88: 630-35, 1963.	lation.
McILREATH F J and COHEN BM: Ventilatory performance of American  Am J Med Sci 252: 1-8, 1966.	physicians. 607
MOHANTY P and GUPTA T C: Smoking habit and pulmonary ventilation Adivasis. Indian J Physiol Pharmacol 12: 87-89, 1968.	in 608
PETERS G A and MILLER R D: Effect of smoking on asthma and emphys  Proc Mayo Clin 35: 353-58, 1960.	ema. 609
PETERSON D I, LONGERGAN L H and HARDINGE M G: Smoking and pufunction. Arch Environ Health 16: 215-18, 1968.	ilmonary 610
PETTY T L, BRINK G A, MILLER M W and CORSELLO PR: Objective functional improvement in chronic airway obstruction. Chest 57 216-23, 1970.	

## VI-B. AIRWAY RESISTANCE

That cigarette smoking reduces forced expiratory volume in certain individuals described in the preceding section indicates a bronchospastic component. The other techniques used to investigate airway resistance are discussed in this section.

1. Body plethysmograph technique. The measurement of air resistance or airway conductance by body plethysmograph technique is more sensitive than others applicable to man. All reports uniformly show an increase in resistance or a decrease in conductance in smokers as compared with nonsmokers: Pelzer and Thomson (1966) reporting from England on 82 subjects 17 to 82 years of age; Sewik (1967) from Czechoslovakia on 150 men 18 to 68 years of age; Guyatt and Alpers (1968) and Guyatt et al. (1970) from England on 752 men 38 to 67 years old; Allen and Sabin (1971) from Texas on 32 males and females 16 to 82 years old; Diamond et al. (1971) from Kentucky on 30 subjects 19 to 44 years old; Huttemann et al. (1971, 1972) from Germany on 20 subjects 20 to 40 years old. The extent of increase in resistance varies with the age of the individual and the country of origin. The bronchoconstrictio is readily reversed by abstinence from smoking for a few days (Cohen, 1966; Wilhelmsen, 1967; McFadden and Linden, 1972) or by the inhalation of a bronchodilator drug (Curtis et al., 1966; Crompton, 1968; Woolcock et al., 1969; Ruff et al., 1972).

various techniques. Immediately after inhalation of cigarette smoke, there is an increase in airway resistance in normal subjects, both smokers and nonsmokers (Attinger et al., 1958; Sterling, 1967; Chiang and Wang, 1970; Reintjes et al., 1972; Da Silva and Hamosh, 1973). The bronchoconstriction was reduced by prior administration of atropine, indicating that the vagus nerve participates in the response (Sterling, 1967). Filtration of the smoke to remove either the particulate or the vapor phase had a similar effect in reducing the bronchoconstrictor response, showing that both phases contribute to this response (Clarke et al., 1970). Inhalation of isoproterenol prevented the response, which reflects its reversibility (Nadel and Comroe, 1961).

The acute bronchoconstriction has been reported following inhalation of cigarette smoke in patients with chronic lung disease:

Simonsson (1962, 1965) reporting from Sweden; Cellereno and Billia (1962) from Italy; McDermott and Collins (1965) from Great Britain;

Johannsen (1966) from Poland; James (1970) from Ohio; and Votchal et al. (1970) from the Soviet Union. These effects are transient in nature and do not necessarily contribute to the abnormal ventilatory function.

3. Acute and chronic effects of inhalation of cigarette smoke in animals. The author of this review and his collaborators reported the effects of inhalation of cigarette smoke in several animal species:

dog (Aviado and Palecek, 1967; Klide and Aviado, 1968; Cho et al., 1968;

Carrillo and Aviado, 1970); cat (Palecek and Aviado, 1967); rabbit

(Palecek et al., 1967); and rat (Aviado and Carrillo, 1969; Shore and

Aviado, 1969; Aviado et al., 1970). The bronchoconstriction is brought

about by the following mechanisms: (a) stimulation of cholinergic

receptors in the airways; (b) stimulation of sensory receptors in

the lungs, which triggers reflex spasm and (c) release of histamine,

which produces bronchospasm (Aviado and Samanek, 1965; Samanek

and Aviado, 1965; Samanek et al., 1965). Simultaneously, bronchodilatation

is also produced by stimulation of adrenergic mechanisms arising from

nicotine contained in the cigarette smoke and by reflex mechanism.

Following chronic exposure to cigarette smoke, there is an increase in airway resistance in rats (Ito and Aviado, 1968; Aviado, 1972) and mice (Aviado, 1972). This effect is reversed by inhalation of sympathomimetic drugs. The results in animals serve to explain the reversibility of bronchoconstriction as seen in man and described in the above paragraphs.

B. AIRWAY RESISTANCE	
경기를 하고 있는데 보고 있다. 1920년 - 1921년 - 1922년	Reprint
ALLEN G W and SABIN S: Comparison of direct and indirect measurement of airway resistance. Am Rev Resp Dis 104: 61-71, 1971.	y 621
ATTINGER E O, GOLDSTEIN M M, and SEGAL M S: Effects of smoking upon the mechanics of breathing. I. In normal subjects. Am Rev Tuberc Pulm Dis 77: 1-16, 1958.	622
AVIADO D M: Bronchovascular effects of cigarette smoke. CTR Grant No. 599 Progress Report No. 4, June 1, 1971 - April 15, 1972, 19 pages	623
AVIADO D M and CARRILLO L R: Hormones and pulmonary effects of tobacco.  Arch Environ Health 18: 925-33, 1969.	624
AVIADO D M and PALECEK F: Pulmonary effects of tobacco and related substances. Arch Environ Health 15: 187-93, 1967.	625
AVIADO D.M. SADAVONGVIVAD C and CARRILLO L R: Cigarette smoke and pulmonary emphysema. Influence of bronchodilators and biogenic amines in experimental induction in rats. Arch Environ Health 20: 483-87,	626
AVIADO D M and SAMANEK M: Bronchopulmonary effects of tobacco and related substances. Arch Environ Health 11: 141-51, 1965.	627
AVIADO D M, WATANABE T and BELEJ MA: Bronchovascular effects of cigarette smoke. CTR Grant # 599 Prog. Rep. # 5, 27 pages, 1973.	628
CELLERINO A and BILLIA P: Volumi polmonari in fumatori sani e broncopatici prima e dopo uso di filtro chimico per sigarette. Minerva Med 53: 1516-21, 1962.	629
CHIANG S T and WANG B C: Acute effects of cigarette smoking on pulmonary function. Am Rev Resp Dis 101: 860-67, 1970.	630
CHO Y W, SAMANEK M and AVIADO D M: Differences in the effects of inhalation of sulfur dioxide and cigarette smoke.  Arch Environ Health 16: 651-55, 1968.	631
CLARKE B G, GUYATT A R, ALPERS J H, FLETCHER C M and HILL I D:  Changes in airways conductance on smoking a cigarette. A study of repeatability and of the effect of particulate and vapour phase filters.  Thorax 25: 418-22, 1970.	632
COHEN B M and ROCHESTER M D: Sympathomimetic amine aerosol administration in chronic obstructive ventilatory disease. Med Times 94: 355-59, 1966.	633
CROMPTON G K: A comparison of responses to bronchodilator drugs in chronic bronchitis and chronic asthma. Thorax 23: 46-55, 1968.	634
CURTIS J K, ASHLEY P L, RASMUSSEN H K and CREE E M: The bronchospastic component in patients with chronic bronchitis and emphysema. JAMA	4.5

197: 693-97, 1966.

page 129	reprint
DA SILVA A M T and HAMOSH P: Effect of smoking a single cigarette on the 'small airways'. <u>J Appl Physiol</u> 34: 361-65, 1973.	<b>6</b> 36
DIAMOND L, LIPSCOMB W and JOHNSON R: Acute pulmonary effects of smoking a reference cigarette. <u>Toxicol Appl Pharmacol</u> 18: 638-48, 1971.	637
GUYATT A R and ALPERS J H: Factors affecting airways conductance: a study of 752 working men. J Appl Physiol 24: 310-16, 1968.	638
GUYATT A R, BERRY G, ALPERS J H, BRAMLEY A C and FLETCHER C M:  Relationship of airway conductance and its immediate change on smoking habits and symptoms of chronic bronchitis.  Am Rev Resp Dis  101: 44-54, 1970.	639
HUTTEMANN U, OSWALD P, LODE H and HUCKAUF H: Veranderungen der lungenfunktion jugendlicher als folge langjahrigen zigarettenrauchens.  (Changes in pulmonary function of young people as a result of cigarette smoking over many years).  Deutsche Med Wochenschrift 96: 1791-93, 1971.	640
HUTTEMANN U, OSWALD P, LODE H and HUCKAUF H: Uber den einfluss langjahrigen zigarettenrauchens auf die lungenfunktion jugendlicher normalpersonen. (Lung function studies in young apparently healthy cigarette smokers versus nonsmokers). Respiration 29: 270-87, 1972.	641
ITO H and AVIADO M: Pulmonary emphysema and cigarette smoke. Experimental induction and use of bronchodilators in rats. Arch Environ Health  16: 865-70, 1968.	642
JAMES R H: Prior smoking as a determinant of the distribution of pulmonary ventilation. Am Rev Resp Dis 101: 105-7, 1970.	643
JOHANNSEN Z and PREIBISZ J: Bezposredni wplyw palenia papierosow na spirogram u zdrowych i chorych z rozedma pluc i przewleklym niezytem oskrzeli. (A direct effect of smoking on spirometric values in normal subjects and patients with pulmonary emphysema and chronic bronchitis). Pol Arch Med Wzwn 36: 783-86, 1966.	644
KLIDE A M and AVIADO D M: Carotid receptors and bronchomotor responses.  Effects of cigarette smoke, lobeline, and cyanide. Arch Environ Health 17: 65-70, 1968.	645
McDERMOTT M and COLLINS M M: Acute effects of smoking on lung airways resistance in normal and bronchitic subjects. Thorax 20: 562-69, 1965.	<b>64</b> 6
McFADDEN ERJr and LINDEN DA: A reduction in maximum mid-expiratory flow rate. A spirographic manifestation of small airway disease. <u>Am J Med</u> 52: 725-37, 1972.	647
MILLER J M and SPROULE B J: Acute effects of inhalation of cigarette smoke on mechanical properties of the lungs. Am Rev Resp Dis 94: 721-26, 1966.	<b>64</b> 8
NADEL JA and COMROE JH Jr: Acute effects of inhalation of cigarette smoke on airway conductance. <u>J Appl Physiol</u> 16: 713-16, 1961.	649
PALECEK F and AVIADO D M: Pulmonary effects of tobacco and related substance II. Comparative effects of eigarette smoke, nicotine, and histamine on the anesthetized cat. Arch Environ Health 15: 194-203, 1967.	es. <b>6</b> 50

Bibliography VI-B page 130	Reprint
PALECEK F, OSKOUI M and AVIADO D M: Pulmonary effects of tobacco and related substances. III. Inhibition of synthesis of histamine in various species. Arch Environ Health 15: 204-13, 1967.	651
PELZER A M and THOMSON M L: Effect of age, sex, stature, and smoking habits on human conductance. <u>J Appl Physiol</u> 21: 469-76, 1966.	652
REINTJES M, SEIERENGA J and BOGAARD J M: Effect of smoking one cigarette on airway resistance. Scand J Resp Dis 53: 129-34, 1972.	653
ROBERTSON D G, WARRELL D A, NEWTON-HOWES J S and FLETCHER C M:  Bronchial reactivity to cigarette and cigar smoke.  Br Med J  3: 269-71, 1969.	654
RUFF F, SALEM A, BUSY F, de VERNEJOUL P, EVEN P and BROUET G:  La fermeture des voies aeriennes peripheriques son augmentation chez les fumeurs. (Airway closure and its increase in smokers).  Rev Tuberc Pneumol 36: 308-11, 1972.	655
SAMANEK M and AVIADO D M: Bronchopulmonary effects of tobacco and related substances. II. Bronchial arterial injection of nicotine and histamine. Arch Environ Health 11: 152-59, 1965.	656
SAMANEK M, AVIADO D M and PESKIN G W: Bronchopulmonary effects of tobacco and related substances.  Arch Environ Health 11: 160-76, 1965.	657
SEVCIK M: Vliv koureni na funkci plic a bronchospasmus zjistovany pomoci farmkodynamickeho testu. (Influence of smoking on pulmonary function and bronchospasm ascertained by a pharmacodynamic test).  Scripta Med Fac Med Brunchis 40: 265-77, 1967.	658
SHORE SR and AVIADO DM: Hormones and pulmonary effects of tobacco.  II. Progesterone. Arch Environ Health 19: 59-69, 1969.	659
SIMONSSON B: Effect of cigarette smoking on the forced expiratory flow rate.  Am Rev Resp Dis 85: 534-39, 1962.	<b>6</b> 60
SIMONSSON B G: Clinical and physiological studies on chronic bronchitis.  III. Bronchial reactivity to inhaled acetylcholine. Acta Allergol  20: 325-48, 1965.	661
STERLING G M: Mechanism of bronchoconstriction caused by cigarette smoking.  Br Med J 3: 275-77, 1967.	662
VOTCHAL B E, LOZINSKY L G, GRIBOVA O A, TAGAI V A and SHEFER Y M:  Condition of bronchial patency in young people. Terap Arkhiv 42: 39-43, 1970.	663
WILHELMSEN L: Effects on bronchopulmonary symptoms, ventilation and lung mechanics of abstinence from tobacco smoking. Scand J Resp Dis 48: 407-14, 1967.	664
WOOLCOCK A J, VINCENT N J and MACKLEM P T: Frequency dependence of compliance as a test for obstruction in the small airways.  J Clin Invest 48: 1097-1105, 1969.	665

## VI-C. BRONCHIAL CLEARANCE

Cigarette smoking retards the clearance mechanism in certain experimental conditions. The effects of cigarette smoking on ciliary motility and mucus secretion are part of the defense mechanisms in the lung against inhalants in general and air pollutants in particular. (Gross, 1967; Gross and Hatch, 1963; Kilburn, 1967, 1968).

1. Mucociliary transport in man. The first observations relating to the effect of cigarette smoking on the respiratory epithelium were made in vitro (Rakieten et al., 1912). Strips of the nasal mucosa were examined under the microscope. The solution through which cigarette smoke was passed did not influence ciliary activity for a 5-to 10-min period. Longer exposure caused cessation of ciliary activity. These results were difficult to relate to the usual manner of smoking. Ballenger (1960) introduced another in vitro technique, using fresh mucosa scraped under general anesthesia from the trachea of children. The ciliotoxicity from the smoked solution is also difficult to relate to the usual form of smoking.

The introduction of radioactive particles for measurement of bronchial clearance has made it possible to investigate the influence of cigarette smoking in man. The results so far have not been consistent. Albert et al. (1969 a) used monodisperse ferric oxide aerosols (0.8 to 7.9  $\mu$ ): bronchial clearance was impaired in 8 out of 14 cigarette smokers. Pavia et al. (1970, 1971) and Thomson and Pavia (1973) used radioactive polystyrene particles 5 µ in diameter. There was no demonstrable difference in mucociliary transport between smokers

and nonsmokers. Lourence et al. (1971) used particles 2 µ in size and noted that the clearance was delayed in smokers as compared with nonsmokers. Camner and Philipson (1971) and Camner et al. (1971, 1973) used 6 to 7 µ particles and observed a faster clearance immediately following smoking. However, habitual smokers showed an increase in clearance three weeks after abstaining. There appear to be two kinds of effects produced by cigarette smoking, an acute type which accelerates and a chronic type which decelerates clearance. Mucociliary transport in the nasal cavity is either retarded by smoking (Evert, 1965) or not influenced by it (Quinlan et al., 1969).

In 10 monozygotic pairs of twins that were discordant with regard to cigarette smoking, half of the smoking pairs had a markedly slower clearance than the nonsmokers, whereas the other half had a similar clearance (Camner and Philipson, 1972; Camner et al., 1972). It has not been possible to obtain a consistent effect from cigarette smoking even by comparing a smoker with a nonsmoker who is constitutionally similar.

2. Mucociliary transport in the rabbit. The results in animals are also variable. There was no effect from two hours' exposure to cigarette smoke on ciliary activity in vivo in the rabbit (Dalhamn, 1964) and on bronchial clearance of radioactive particles in the same species (LaBelle et al., 1966). A depression has been reported by others, using different techniques for measuring ciliary activity (Krueger and Smith, 1958) and clearance (Holma, 1969).

The functional activity of the ciliated epithelium can be influenced by oxygen tension (Dalhamn and Rosengren, 1968), particulate matter (Falke et al., 1959), and acetylcholine (Kordek et al., 1952). Cigarette smoke can exert its influence through any one or more of these mechanisms and this may account for the varied results reported in the rabbit. Kensler and Battista (1963), by using smoke filters, concluded that the ciliotoxic activity is in the gaseous phase. The following constituents of smoke depress ciliary activity in the rabbit: carboxylic acid and aldehydes (Wynder et al., 1965), phenol (Dalhamn, 1966) and acrolein (Scharbort, 1967).

3. Mucociliary transport in the cat. The observations in the cat have consistently shown a depression of bronchial clearance by exposure to cigarette smoke (Dalhamn, 1964, 1966; Carson et al., 1966; Kaminski et al., 1968). The tar content of the cigarette smoke influenced its ciliotoxicity (Nakashima, 1938; Dalhamn and Rylander, 1964, 1965, 1967, 1968). This effect was not simulated by nicotine but could be prevented by oxolamine citrate (Dalhamn, 1968, 1969). role of the volatile constituents in cigarette smoke has been debated (Kensler and Battista, 1966; Dalhamn and Rylander, 1967, 1970; Kensler, 1967). Some chemical constituents of cigarette smoke have been shown to interact in their effect on mucociliary transport, might explain the varied action reported for cigarette smoke (W

4. Mucociliary transport in other animal species. The following animal species have been used in investigation of the influence of cigarette smoke: rat (Dalhamn and Rhodin, 1956; Dalhamn, 1959; Ferin et al., 1966; Guillerm et al., 1961); guinea pig (Hilding and Filipi, 1966); cow (Hill, 1928; Mendenhall and Shreeve, 1937; Hilding, 1956 a, b); monkey (Kensler and Battista, 1966); chicken (Battista and Kensler, 1970 a, b); donkey (Albert et al., 1969 b, 1973; Frances et al., 1970); and invertebrates (Boche and Quilligan, 1959; Wynder et al., 1963, 1965; Walker and Kiefer, 1966; Dalhamn, 1967; Weiss, 1968; Krahl and Bulmash, 1969; Kennedy and Elliott, 1970). Ciliary activity consistently shows a depression by cigarette smoke but bronchial clearance is influenced in a variable manner, the latter by ciliary activity and mucus formation, as discussed in Section III.

# BIBLIOGRAPHY

VI. BRONCHOPULMONARY FUNCTION IN SMOKERS AND NONSMOKERS

## C. BRONCHIAL CLEARANCE

學學하면 가장하면 되었다. 그 그 사람이 그리고 아니는 하는 그 회사는 이 그를 가장하는 그	Reprint
ALBERT R E, ALESSANDRO D, LIPPMANN M and BERGER J: Long-term smoking in the donkey. Arch Environ Health 22: 12-19, 1971.	<b>6</b> 66
ALBERT R E, LIPPMANN M and BRISCOE W: The characteristics of bronchial clearance in humans and the effects of cigarette smoking. Arch Environ Health 18: 738-55, 1969.	667
ALBERT R E, LIPPMANN M, PETERSON H T Jr, BERGER J, SANBORN K and BOHNING D: Bronchial deposition and clearance of aerosols.  Arch Int Med 131: 115-27, 1973.	668
ALBERT R E, SPIEGELMAN J R, SHATSKY S and LIPPMANN M: The effect of acute exposure to cigarette smoke on bronchial clearance in the miniature donkey. Arch Environ Health 18: 30-41, 1969.	<b>6</b> 69
BALLENGER JJ: Experimental effect of cigarette smoke on human respiratory cilia. N Engl J Med 263: 832-35, 1960.	670
BATTISTA S P and KENSLER C J: Use of the nonimmersed in vitro chicken tracheal preparation for the study of ciliary transport activity.  Arch Environ Health 20: 318-25, 1970. A.	671
BATTISTA S P and KENSLER C J: Mucus production and ciliary transport activity.  In vivo studies using the chicken. Arch Environ Health 20: 326-33,  1970 B	672
BOCHE R D and QUILLIGEN J J Jr: The effects of air pollutants on tissue cultures.  Fed Proc Bull 18: 559, 1959.	673
CAMNER P and PHILIPSON K: Humanstudier av tracheobronchiell clearance med monodispersa terflonpartiklar markta med 18 <sub>F</sub> och 99 mTc.  (Tracheobronchial clearance in man studied with monodispersed teflon particles tagged with 18 <sub>F</sub> and 99 mTc. Nord Hyg Tidskr 52:	
1-9, 1971.	674
CAMNER P and PHILIPSON K: Tracheobronchial clearance in smoking-discordant twins. Arch Environ Health 25: 60-63, 1972.	675
CAMNER P. PHILIPSON K and ARVIDSSON T: Cigarette smoking in man. Short- term effect on mucociliary transport. Arch Environ Health 23: 421-26, 1971.	<b>6</b> 76
CAMNER P. PHILIPSON K and ARVIDSSON T: Withdrawal of cigarette smoking.  A study of tracheobronchial clearance.  1973.  Withdrawal of cigarette smoking.  Arch Environ Health 26: 90-92,	677
CAMNER P. PHILIPSON K and FRIBERG L: Tracheobronchial clearance in twins. Arch Environ Health 24: 82-87, 1972.	678
CARSON S, GOLDHAMER R and CARPENTER R: Mucus transport in the respiratory tract. Am Rev Resp Dis 93: 86-92, 1966.	679

page 136	
CRESS HR, SPOCK A and HEATHERINGTON DC: The localization of succinic dehydrogenase, cytochrome oxidase and adenosine triphosphatase in ciliated respiratory epithelium. J Histochem Cytochem 13: 677-83, 1965.	<b>6</b> 80
DALHAMN T: The effect of cigarette smoke on ciliary activity in the upper respiratory tract. Arch Otolaryngol 70: 166-68, 1959.	681
DALHAMN T: Studies on tracheal ciliary activity. Am Rev Resp Dis 89: 870-77, 1964.	682
DALHAMN T: Effect of cigarette smoke on ciliary activity. Am Rev Resp Dis 93: 108-14, 1966.	683
DALHAMN T: Effect of different doses of tobacco smoke on ciliary activity in cat. Variations in amount of tobacco smoke, interval between cigarettes, content of "tar", nicotine, and phenol. Natl Cancer Inst Monogr 28: 79-87, 1968.	684
DALHAMN T: The anticillostatic effect of cigarettes treated with oxolamine citrate. Am Rev Resp Dis 99: 447-48, 1969.	685
DALHAMN T: Ciliostatic effect of phenol and resorcinol. Arch Otolaryngol 84: 325-32, 1966.	<b>6</b> 86
DALHAMN T and RHODIN J: Mucous flow and ciliary activity in the trachea of rats exposed to pulmonary irritant gas. Br J Ind Med 13: 110-13, 1956.	687
DALHAMN T and ROSENGREN A: The effect of oxygen lack on the tracheal ciliary activity. Arch Environ Health 16: 371-73, 1968.	688
DALHAMN T and RYLANDER R: Ciliastatic action of smoke from filter-tipped and non-tipped cigarettes. Nature 201: 401-2, 1964.	<b>68</b> 9
DALHAMN T and RYLANDER R: Ciliastatic action of cigarette smoke.  Arch Otolaryngol 81: 379-82, 1965.	<b>6</b> 90
DALHAMN T and RYLANDER R: Tar content and ciliotoxicity of cigarette smoke.  Acta Pharmacol Toxicol 25: 369-72, 1967.	691
DALHAMN T and RYLANDER R: Reply. Arch Environ Health 14: 372-73, 1967.	692
DALHAMN T and RYLANDER R: Ciliotoxicity of cigarette smoke and its volatile components. Am Rev Resp Dis 98: 509-11, 1968.	693
DALHAMN T and RYLANDER R: Ciliotoxicity of cigar and cigarette smoke.  Arch Environ Health 20: 252-53, 1970.	694
DALHAMN T, RYLANDER R and SPEARS A W: Differences in ciliotoxicity of cigarette smoke. Am Rev Resp Dis 96: 1078-79, 1967.	<b>6</b> 95
EVERT G: On the mucous flow rate in the human nose. Acta Otolaryngol 300: 1- 1965.	<b>6</b> 96
FALK H L, TREMER H M and KOTIN P: Effect of cigarette smoke and its constituents on ciliated mucus-secreting epithelium.  J Natl Cancer Inst 23: 999-1012, 1959.	697
FERIN J: The mechanism of elimination of deposited particles from the lungs.  Ann Occup Hyg 10: 207-16, 1967.	<b>6</b> 98

Bibliography VI-C	page 137	Reprint
FERIN J, URBANKOVA G, VICKOVA A and REICHRTOVA E:  of cold and cigarette smoke on the elimination of dust from	The effect	•
<u>Pracov Lek</u> 18: 263-64, 1966.		699
FRANCES R, ALESSANDRO D, LIPPMANN M, PROCTOR D E  Effect of cigarette smoke on particle transport on nasocil  of donkeys. Arch Environ Health 21: 25-31, 1970.		<b>7</b> 00
The Company of the Co	- 33. 1315 37	
GROSS P: Le mecanisme du transport alveolaire, <u>Poumon Coeur</u> 1967.	23: 1213-27,	701
GROSS P and HATCH T: Pulmonary clearance: its mechanism a to pulmonary disease. J Occup Med 5: 191-94, 1963.	nd its relation	702
GUILLERM R, BADRE R and VIGNON B: Effects inhibiteurs de tabac sur l'activite ciliaire de l'epithelium respiratoire e	t nature des	
composants responsables. <u>Bull Acad Natl Med</u> 145: 416	-23, 1961.	703
HILDING A C: On cigarette smoking, bronchial carcinoma and ci III. Accumulation of cigarette tar upon artificially produc islands in the respiratory epithelium. Ann Otololaryng		704
1956A.		•
HILDING A C: On cigarette smoking, bronchial carcinoma and ci	gs the deposition	
of tar in the bronchial tree and removal by ciliary action. 254: 1155-60, 1956B.	N Engl J Med	705
HILDING A C: The role of the respiratory mucosa. Minn Med 1967.	50: 915-19,	706
HILDING A C and FILIPI A N: A method of determining nasal fi cigarette smoke. Ann Otol 75: 714-24, 1966.	ltration of	<b>7</b> 07
HILL L: The ciliary movement of the trachea studied in vitro. 2: 802-5, 1928.	Lancet	<b>7</b> 08
HOLMA B: The acute effect of cigarette smoke on the initial cour clearance in rabbits. Arch Environ Health 18: 171-73,		709
KAMINSKI E J, FANCHER O E and CALANDRA J C: In vivo st ciliastatic effects of tobacco smoke. Arch Environ Halti		
188-93, 1968.	•	710
KENSLER C J. Cigarette smoke and ciliastasis. Arch Environ 14: 371-73, 1967.	Health	711
KENNEDY JR and ELLIOTT AM: Cigarette smoke: Residue of structure. Science 168: 1097-98, 1970.	n mitochondrial	712
KENSLER C J and BATTISTA S P: Components of cigarette sm depressant activity. N Engl J Med 269: 1161-66, 196		713
KENSLER CJ and BATTISTA S P: Chemical and physical factor mammalian ciliary activity. Am Rev Resp Dis 93: 93-	rs affecting	714
KILBURN K H: Cilia and mucus transport as determinants of the lung to air pollutants. Arch Environ Health 14: 77-91,	-	715
KILBURN K H: A hypothesis for pulmonary clearance and its impartment of the control of the contr	olications.	716

Bibliography VI-C page 138	Reprint
KORDICK P, BULBRING E and BURN J H: Ciliary movement and acetycholine.  Br J Pharmacol 7: 67-79, 1952.	717
KRAHL V E and BULMASH M H: Studies on living ciliated epithelium.  Am Rev Resp Dis 99: 711-17, 1969.	718
KRUEGER A P and SMITH R F: Effects of gaseous ions on tracheal ciliary rate. Proc Soc Exp Eiol Med 98: 412-14, 1958.	719
LaBELLE C W. BEVILACQUA D M and BRIEGER H: The influence of cigarette smoke on lung clearnace. Arch Environ Health 12: 588-96, 1966.	720
LOURENCE R V, KLIMER M F and BOROWSKI C J: Deposition and clearance of 2µ particles in the tracheobronchial tree of normal subjects smokers and nonsmokers. J Clin Invest 50: 1411-20, 1971.	721
MENDENHALL W L and SHREEVE K E: The effect of cigarette smoke on the tracheal cilia. J Pharmacol Exp Ther 60: 111-12, 1937.	722
MOOLTEN S E: Pulmonary lymphatics in relation to pulmonary clearance interstitial fluid and the pathogenesis of emphysema. Pub Health Serv 1787: 191-221, 1968.	723
NAKASHIMA T: Pharmacological studies on the ciliary movement. Jap J Med Sci Sect Pharmacol 11: 42, 1938.	724
PAVIA D, SHORT M D and THOMSON M L: No demonstrable long term effects of cigarette smoking on the mucociliary mechanism of the human lung.  Nature 226: 1228-31, 1970.	<b>7</b> 25
PAVIA D, THOMSON M L and POCOCK S J: Evidence for temporary slowing of mucociliary clearance in the lung caused by tobacco smoking.  Nature 231: 325-26, 1971.	<b>7</b> 26
QUINLAN M F, SALMAN S D, SWIFT D L, WAGNER H N Jr, and PROCTOR D F:  Measurement of mucociliary function in man.  99: 13-23, 1969.	727
RAKIETEN N, RAKIETEN M L, FELDMAN D and BOYKIN M J Jr: Mammalian ciliated respiratory epithelium. Arch Otolaryngol 56: 494-503, 1912.	728
SCHARBORT J C: Lactic dehydrogenase from human lung inhibition by certain water-soluble ciliastatic components of tobacco smoke. J South Africa Chem Inst 20: 103-112, 1967.	729
THOMSON M L and PAVIA D: Long-term tobacco smoking and mucociliary clearance. From the human lung in health and respiratory impairment.  Arch Environ Health 26: 86-89, 1973.	<b>7</b> 30
WALKER TR and KIEFER JE: Ciliastatic components in the gas phase of cigarette smoke. Science 153: 1248-50, 1966.	731
WEISS W: Cigarette smoke gas phase and paramecium survival. Arch Environ Health 17: 62-64, 1968.	732
WEISSBECKER L, CREAMER R M and CARPENTER R D: Cigarette smoke and tracheal mucus transport rate. Am Rev Resp Dis 104: 182-87, 1971.	733
WYNDER E L, GOODMAN D A and HOFFMANN D: Ciliatoxic components in cigarette smoke. II. Carboxylic acids and aldehydes. Cancer 18:	734

WYNDER E L, GOODMAN D A and HOFFMANN D: Ciliatoxic components in cigarette smoke. III. In vitro comparison of different smoke components.

Cancer 18: 1652-58, 1965.

735

WYNDER E L, KAISER H E, GOODMAN D A and HOFFMANN D: A method for determining ciliastatic components in cigarette smoke. Cancer 16: 1222-25, 1963.

726

## VII. CARDIOPULMONARY FUNCTION IN SMOKERS AND NONSMOKERS

This section is a continuation of Section VI on the differences between smokers and nonsmokers. Measurements of cardiopulmonary function include ventilation-perfusion relationship, pulmonary circulation, cardiac function, pulmonary function, surfactant and other constituents of the lung. It is important to point out that animal experiments have served to clarify the mode of action by which the constituents of cigarette smoke influence the heart and lungs. The following bibliographic lists contain articles on general aspects of the topic under consideration.

No. 14. Constituents of cigarette smoke.

No. 15. Cardiopulmonary effects of cigarette smoke.

Measurement of pulmonary diffusion capacity shows a reduction in smokers compared with nonsmokers: Martt (1962) reporting from Missouri; Krumholz et al. (1964, 1965) from Indiana; Rankin et al. (1965) from Wisconsin; and Van Ganse et al. (1972) from the Netherlands.

Anderson and Shephard (1969) reported from Canada that among 43 healthy adult men, by serial multiple regression analysis, pulmonary diffusion capacity correlated with age and weight. By using the predicted equation for diffusion coefficient, smoking habits had no substantial effect on diffusion capacity. Bates (1955) and McGrath and Thomson (1959) arrived at a similar conclusion with a limited number of subjects. It has also been suggested that the presence of carboxyhemoglobin in blood of smokers would account for the lower value for diffusion capacity.

Cigarette smokers show some inadequacy of gas mixing in the lung as compared with nonsmokers (Ross et al., 1967; Stanesce et al., 1968).

There is also an alteration in the ventilation-perfusion ratio in smokers
(Williams and Anderson, 1968 a, b; Anderson and Williams, 1969). The change has not been confirmed by others and the participation of pulmonary blood vessels in the phenomenon has not been proven.

Inhaling cigarette smoke by the use of a smoking device that brought more smoke into the lungs with each breath than would be the case in ordinary smoking did not significantly change blood gas exchange in normal subjects or patients with a moderate degree of emphysema (Motley and Kuzman, 1958). However, patients in California with severe emphysema showed decreases in arterial exygen tension and saturation.

A similar hypoxemia was observed to arise from smoking by patients with chronic lung disease in Ireland (Chapman, 1965) by patients with chronic bronchitis in England (Penman and Howard, 1966), and by patients with chronic bronchitis in Massachusetts (Strieder and Kazemi, 1967; Strieder et al., 1968).

Hypoxemia is characteristic of patients in whom it does not necessarily relate to smoking but to pathological lesions in the lung (Pain et al., 1967; Weiss and Dulfano, 1968; Poppius, 1969).

Postoperative complications are reported to be higher among nonsmokers than smokers in Great Britain (Morton, 1944; Piper, 1958; Wightman, 1968). Smokers from Sweden do not show a higher incidence of pulmonary postoperative complications (Wiklander and Norlin, 1957). It has been suggested that postoperative hypoxemia is related to the appearance of pulmonary complications among smokers in Australia (Morton, 1969).

	ANDERSON T W and SHEPHARD R J: Normal values for single-breath diffusing capacity - the influence of age, body size and smoking habits. Respiration 26: 1-7, 1969.	<b>7</b> 37
-	지수 내가 되었다. 그는 사이에는 가장에 가는 사고를 잃어지는 아이들이 되었다. 중요	
1	ANDERSON W H: Acute exposure to cigarette smoke as a cause of hypoxia. Chest 59: 33S-34S, 1971.	738
	ANDERSON W H and WILLIAMS J B: Effects of cigarette smoke on distribution of pulmonary perfusion. In: Current Research in Chronic Respiratory Diseases. Proceedings of the 11th Aspen Emphysema Conference, Aspen, Colo. U.S. Department of Health, Education, and Welfare, Public Health Service Publication 1879: 75-9, 1969.	<b>73</b> 9
	BATES DV, BOUCOT NG and DORMER A E: The pulmonary diffusing capacity in normal subjects. J Physiol 129: 237-52, 1955.	<b>7</b> 40
	BURROWS B: Pulmonary diffusion and alveolar-capillary block. Med Clin N Am 51: 427-38, 1967.	741
	CHAPMAN TT: The acute effect of cigarette smoking on pulmonary function. J Irish  Med Assoc 56: 72-4, 1965.	742
	DORMANDY T L: Abnormal oxygen-dissociation curves. Lancet 2: 80-1, 1965.	743
	FORSTER R E, ROUGHTON F J, CANDER L, BRISCOE W A and KREUZER F:  Apparent pulmonary diffusing capacity for CO at varying alveolar O <sub>2</sub> tensions.  J Appl Physiol 11: 277-89, 1957.	744
	HOLLAND J, MILIC-EMILI J, MACKLEM P T and BATES D V: Regional distribution of pulmonary ventilation and perfusion in elderly subjects. J Clin Invest 47: 81-92, 1968.	745
	KRUMHOLZ RA, CHEVALIER R B and ROSS J C: Cardiopulmonary function in young smokers. A comparison of pulmonary function measurements and some cardio-pulmonary responses to exercise between a group of young smokers and a comparable group of nonsmokers. Ann Internal Med 60: 603-10, 1964.	746
	KRUMHOLZ RA, CHEVALIER R B and ROSS J C: Changes in cardiopulmonary functions related to abstinence from smoking. Studies in young cigarette smokers at rest and exercise at 3 and 6 weeks of abstinence. Ann Internal Med 62: 197-207, 1965.	<b>7</b> 47
	MARTT J M: Pulmonary diffusing capacity in cigarette smokers. Ann Internal Med 56: 39-45, 1962.	<b>74</b> 8
	McGRATH M W and THOMSON M L: The effect of age, body size and lung volume change on alveolar-capillary permeability and diffusing capacity in man. J Physiol 146: 572-82, 1959.	749
	MORTON A: Postoperative hypoxaemia. Med J Australia 2: 341-2, 1969.	750
	MORTON H J: Tobacco smoking and pulmonary complications after operation. Lancet 1: 368-70, 1944.	751
	MOTLEY H L and KUZMAN W J: Cigarette smoke. Its effects on pulmonary function measurements. California Med 88: 211-20, 1958.	752

Bibliography VII-A	Repri
PAIN M C, GLAZIER J B, SIMON H and WEST J B: Regional and overall inequality of ventilation and blood flow in patients with chronic airflow obstruction. Thorax	
22: 453-61, 1967.	753
PENMAN R W: Regional and overall ventilation-perfusion ratios in normal subjects and patients with chronic lung disease. In: Current Research in Chronic Respiratory Diseases. Proceedings of the 11th Aspen Emphysema Conference, Aspen, Colo.  U.S. Department of Health, Education, and Welfare, Public Health Service Publication 1879: 63-73, 1969.	754
PENMAN R W and HOWARD P: Distribution of pulmonary ventilation and blood flow in normal subjects and patients with chronic bronchitis. Clin Sci 30: 63-78, 1966.	<b>7</b> 55
PIPER DW: Respiratory complications in the post-operative period. Scottish Med J 3: 193-8, 1958.	<b>7</b> 56
POPPIUS H: Expiratory CO <sub>2</sub> curve in pulmonary diseases. Scand J Resp Dis 50: 135-46, 1969.	757
RANKIN J, GEE J B and CHOSY L W: The influence of age and smoking on pulmonary diffusing capacity in healthy subjects. Medicina Thoracalis 22: 366-74, 1965.	<b>7</b> 58
ROSS J C, LEY G D, KRUMHOLZ R A and RAHBARI H: A technique for evaluation of gas mixing in the lung: studies in cigarette smokers and nonsmokers. Am Rev Resp Dis 95: 447-53, 1967.	<b>7</b> 59
STANESCU D C, TECULESCU D B. PACURARU R and GAVRILESCU N: Chronic effect of smoking upon pulmonary distribution of ventilation in healthy males.  Respiration 25: 497-504, 1968.	760
STRIEDER D J and KAZEMI H: Hypoxemia in young asymptomatic cigarette smokers.  Ann Thoracic Surg 4: 523-31, 1967.	761
STRIEDER D J, MURPHY R and KAZEMI H: Hypoxemia in asymptomatic smokers. <u>Clin Res</u> 16: 376, 1968.	<b>7</b> 62
STUART-HARRIS C H: Pulmonary hypertension and chronic obstructive bronchitis.  Am Rev Resp Dis 97: 9-17, 1968.	763
VAN GANSE W, DEPOORTER A M, MEHEUS A, EYLENBOSCH W, VANHOORNEEN M, VUYLSTEEK K, DE VREEZE G and RAVEEL E: Bijdrage van de pulmonaire diffusiecapaciteit (transferfactor van de long) in het kader van bevolkingsonderzoek naar CARA. (The role of the pulmonary diffusion capacity (transfer factor of the lung) in screening for CARA) Nederlands Tijdschrift voor Geneeskunde 116/28: 1167-72, 1972.	764
WEISS E B and DULFANO M J: Quantitative acid-base dynamics in chronic pulmonary disease. Ann Internal Med 69: 263-72, 1968.	765
WIGHTMAN J A: A prospective survey of the incidence of postoperative pulmonary complications. Brit J Surg 55: 85-91, 1968.	766
WIKLANDER O and NORLIN U: Effect of physiotherapy on post-operative pulmonary complications. A clinical and roentgenographic study of 200 cases. Acta Chirurgica Scand 112: 246-54, 1957.	767
WILLIAMS J B and ANDERSON W H: Acute effects of cigarette smoke on pulmonary ventilation-perfusion relationships. Am Rev Resp Dis 98: 145-6, 1968a.	768
WILLIAMS J B and ANDERSON W H: Acute effects of cigarette smoke on distribution	769

#### VII-B. PULMONARY CIRCULATION

The pulmonary vascular lesions in pulmonary emphysema have been summarized by Liebow (1959). The lesions in the small pulmonary arteries are severe enough to cause elevation of pulmonary arterial blood pressure (Hicken et al., 1965). Electron microscopy has revealed damage to the endothelium, filling of the capillary lumina with collagen, and recanalization of the damaged vessels (Martin and Boatman, 1965).

Crenshaw (1968) has reviewed the hypothesis that pulmonary emphysema has a vascular etiology. However, none of the vascular lesions seen in human pulmonary emphysema have been reproduced in animals by cigarette smoking. The lungs of healthy cigarette smokers show fibrous thickening which is also seen in the aging lungs of nonsmokers (Auerbach et al., 1968).

The acute effect of smoking one cigarette was studied in 67 Indian patients suffering from chronic bronchopulmonary disease (Viswanathan, 1965). There was a significant rise in mean and diastolic pressure in the pulmonary artery, but no significant change in systolic. The change in diastolic pressure was interpreted to mean vasoconstriction, although there was no measurement of cardiac output nor of left atrial pressure to allow estimation of pulmonary vascular resistance. The fall in pulmonary compliance seen among smokers does not relate to changes in the pulmonary circulation (Krumholz et al., 1965).

There are some observations relating to pulmonary circulation in animals exposed to cigarette smoking. Larson et al. (1965) used nicotine intravenously and noted a biphasic rise and then fall in pulmonary vascular resistance in the dog. Samanek and Aviado (1966) noticed a rise in pulmonary arterial blood pressure following inhalation of cigarette smoke, but the

response could not be repeated in the same animal for purposes of eliciting the mechanism of the rise. Nicotine injection was utilized and the following mechanisms were identified: vasoconstriction by release of catecholamine stores in the lung and the release of histamine.

### BIBLIOGRAPHY

# VII. CARDIOPULMONARY FUNCTION IN SMOKERS AND NONSMOKERS

#### B. PULMONARY CIRCULATION

그 그 어머니의 집에 가게 들어 보고 있는데 아버지는 사람들이 가장 그 것은데 이렇게 되는데 되었다.	Kepri
AUERBACH O, HAMMOND E C and GARKINKEL L: Thickening of walls of arteric and small arteries in relation to age and smoking habits. New Engl J Med 2	oles 78:
	770
CRENSHAW G L: Vascular etiology of pulmonary emphysema. 1968.	771
HICKEN P, HEATH D, BREWER D B and WHITAKER W: The small pulmonary arteries in emphysema. <u>J Path Bacteriol</u> 90: 107-14, 1965.	772
KRUMHOLZ RA, CHEVALIER R B and ROSS J C: A comparison of pulmonary con	<b>m-</b>
pliance in young smokers and nonsmokers. Am Rev Resp Dis 92: 102-7, 1965	773
LARSON R K, FUKUDA P and MURRAY JF: Systemic and pulmonary vascular ef	fects
of nicotine in anesthetized dogs. Am Rev Resp Dis 91: 556-64, 1965.	774
LIEBOW A A: Pulmonary emphysema with special reference to vascular changes.  Am Rev Resp Dis 80: 67-93, 1959.	<b>7</b> 75
MARTIN H B and BOATMAN E S: Electron microscopy of human pulmonary emphys	sema.
Am Rev Resp Dis 91: 206-9, 1965.	776
SAMANEK M and AVIADO D M: Cardiopulmonary effects of tobacco and related sul III. Pulmonary vascular effects of cigarette smoke and nicotine. Arch Environment	bstances. on Health
12: 717-24, 1966.	777
VISWANATHAN R: Acute effects of cigarette smoking on pulmonary artery pressur- Indian J Med Res 53: 421-7, 1965.	es. <b>77</b> 8

# 1005051464

#### VII-C. CARDIAC FUNCTION

Since cigarette smoking has been identified as the most important cause of pulmonary emphysema by the United States Public Health Service, it has been implicated in cor pulmonale (Parade, 1969; Behnke et al., 1970).

In a group of 150 patients with cor pulmonale in India, 28% were nonsmokers (Sanghvi and Kotia, 1963). In another group of 134 patients in India, 39% were nonsmokers (Khosla et al., 1969). In the United States the proportion of nonsmokers is lower among patients with cor pulmonale; Cullen et al. (1970) reported from New York only 3 out of 60 patients. Patients dying of cor pulmona have manifested failure of the left ventricle but the role of cigarette smoking has not been proven (Mitchell et al., 1968; Rao et al., 1969; Strong set al., 1969).

#### BIBLIOGRAPHY

# VII. CARDIOPULMONARY FUNCTION IN SMOKERS AND NONSMOKERS

# · C. CARDIAC FUNCTION

			R	eprı
pathologic stu	MERLEN JT, DAOUD A and dy of the lungs and heart in chre 190-204, 1970.		ase. Am Rev	<b>7</b> 79
•				
	MMISSION FOR HEART DISEAS			
	art disease. Circulation 41: A			<b>7</b> 80
	OTRA A N and ANKLESARIA			A10
pulmonale in I	Northern India J Indian Med A	ssoc 53: 529-34, 1969.		<b>7</b> 81
MITCHELL BS WA	LKER SH and MAISEL JC:	The causes of death in ch	ronic airman	
	I. Myocardial infarction. Am			<b>7</b> 82
PARADE D: Cor pul	monale und Rauchen. Med Klin	n 64: 80-4, 1969.		<b>7</b> 83
RAO BS, COHN K	E, ELDRIDGE F E and HANCO	CK E'W: Left ventricula	r failure	
-	chronic pulmonary disease. <u>In</u>		The state of the s	
	iscases. Proceedings of the ll			
	U.S. Department of Health, E e Publication 1879: 129-33, 1969			<b>7</b> 84
	OTIA K C: Heart failure in pa Dis Chest 44: 67-78, 1963.	tients with chronic pulmo	nary	<b>7</b> 85
	ARDS M L, MCGILL H C, EGO ation of cigarette smoking with	•		
	Res 10: 303-17, 1969.			786

#### PULMONARY SURFACTANT AND OTHER CONSTITUENTS

Stability of the pulmonary alveoli is partially due to changes in surface tension. The surface active component of lung extract which is responsible for maintaining alveolar distension has been called surfactant. Bondurant (1960), Miller et al (1960) and Miller and Bondurant (1962) reported that the application of cigarette smoke to lung extract caused a decrease in surface tension. They also reported that lung extract prepared from rats which have been exposed to smoke in life have lower surface tension than that of unexposed rats. They concluded that these results may relate to the pathogenesis of pulmonary emphysema.

Subsequent reports have confirmed the reduction of surfactant in bronchial washings of the dog by in vitro application (Webb et al., 1967) and in the dog, cat and guinea pig by in vivo inhalation of cigarette smoke (Giammona, 1967; Sekulic et al., 1968). Two recent reports include the incorporation of radioactive labeled palmitic acid into pulmonary surfactant. Giammona et al. (1971) used unanesthetized dogs exposed repeatedly to cigarette smoke daily for 37 to 87 days. They noted no change in surface activity or in rate of time of incorporation of palmitic acid in these dogs. Balint et al. (1971) used anesthetized dogs and administered cigarette smoke by positive ventilation from 90 to 220 min. They reported a decrease in choline and phosphate incorporation in the lecithin of the lung. The results in the unanthestetized dog are similar to the effects obtained by the usual use of tobacco in man, whereas those in the anesthetized dog are toxicological

Measurement of surfactant activity in man has been made on bronchial washings. Cook and Webb (1966) noted a decrease of surfactant action in nonsmokers with lung disease and in chronic smokers with or without demonstrable pulmonary disease. A reduction in the white layer of the lavage fluid after centrifugation has been reported in smokers (Pratt et al., 1969). There is a deficit in total lipid content, particularly of lecithin, in the lavage from smokers as compared with that from nonsmokers (Finley and Ladman, 1972). These observations have been commented upon by Clements (1972), Sachs (1972) and Finley (1972) in relation to the shortcomings of using lavage fluid in interpreting the events occurring in the alveolar lining.

There are other reported changes in the lung of patients with chronic lung disease. The reduction in tissue content of cadmium and zinc in patients with chronic bronchitis or pulmonary emphysema by Szadkowski et al. (1969). Lewis et al. (1969), and Morgan (1971) have not been examined in the lungs of smokers without lung disease. Inhalation of cigarette smoke causes a release of histamine from lung tissue (Aviado et al., 1966) and from the guinea pig lung (Saindelle et al., 1968; Lewis and Nicholls, 1972). Exposure to cigarette smoke causes an increase in protein-mucopolysaccharide complexes in the connective tissue of the guinea pig lung (Lupu and Velican, 1962), an increase in hydroxyproline content of the mouse lung (Rosenkrantz et al., 1969) and a change in electron spin resonance of the rabbit lung (Rowlands et al., 1967). The significance of these observations in relation to lung disease has not been elucidated.

802

macrophages and pulmonary surfactant (?) obtained from the lungs of numan smokers and nonsmokers by endobronchial lavage. Anat Rec 163: 497-507, 1969.

# VIII. COMMENTARY ON SELECTED ARTICLES AND THEIR PUBLICITY

# VIII-A. COMMENTARY ON U.S. PUBLIC HEALTH SERVICE PUBLICATIONS

The author of this review has prepared a page by page critique of the seven documents distributed by the U.S. Public Health Service. A few examples of misquotation, omission and misinterpretation have been selected and are appended in this report. The reports have been identified by the year of their appearance. The pages pertinent in the subject of pulmonary disease are as follows:

·(1)	1964:	Smoking and Health. pp 259-315.		810
(2)	1967:	The Health Consequences of Smoking.	pp 87-124.	811
(3)	1968:	The Health Consequences of Smoking.	pp 63-86.	812
(4)	1969:	The Health Consequences of Smoking.	pp 35-52.	813
<b>(</b> 5)	1971:	The Health Consequences of Smoking.	pp 137-230.	814
(6)	1972:	The Health Consequences of Smoking.	pp 35-56.	815
(7)	1973:	The Health Consequences of Smoking.	pp 43-91.	816

#### Conclusions of the 1964 Report (59)

1. Cigarette smoking is the most important of the causes of bronchitis in the United States, and increases the risk of dying from chronic

chitis in the United States, and increases the rest of a many states bronchitis.

2. A relationship exists between pulmonary emphysema and cigarette smoking but it has not been established that the relationship is causal. The smoking of cigarettes is associated with an increased risk of dying from pulmonary emphysema.

3. For the bulk of the population of the United States, the importance of cigarette smoking as a cause of chronic bronchopulmonary disease is much greater than that of atmospheric pollution or occupational exposures. tional exposures.

 4. Cough, sputum production, or the two combined are consistently more frequent among cigarette smokers than among nonsmokers.
 5. Cigarette smoking is associated with a reduction in ventilatory function. Among males, cigarette smokers have a greater prevalence of breathlessness than nonsmokers.
 6. Cigarette smoking does not appear to cause asthma.
 7. Although death certification shows that cigarette smokers have a moderately increased risk of death from influenza and pneumonia, an association of cigarette smoking and infectious diseases is not otherwise substantiated. wise substantiated.

#### HIGHLIGHTS OF THE 1967 REPORT (57)

1. New data confirm and to some extent strengthen the conclusions of the Surgeon General's 1964 Report.

Cigarette smoking is the most important of the causes of chronic non-neoplastic bronchopulmonary diseases in the United States. It greatly increases the risk of dying not only from both chronic bronchitis but also from pulmonary emphysema.
 Cessation of smoking is followed by a reduction in mortality from chronic bronchopulmonary disease relative to the mortality of those who continue to smoke

who continue to smoke.

4. Even relatively young cigarette smokers frequently have demonstrable respiratory symptoms and reduction in ventilatory function.

#### SUMMARY AND CONCLUSIONS 1971

1. Cigarette smoking is the most important cause of chronic obstructive bronchopulmonary disease in the United States. Cigarette smoking increases the risk of dying from pulmonary emphysema and chronic bronchitis. Cigarette smokers show an increased prevalence of respiratory symptoms, including cough, sputum production, and breathlessness, when compared with nonsmokers. Ventilatory function is decreased in smokers when compared with nonsmokers.

2. Cigarette smoking does not appear to be related to death from bronchial asthma although it may increase the frequency and severity of asthmatic attacks in patients already suffering from this diseasc.

3. The risk of developing or dying from COPD among pipe and/ or cigar smokers is probably higher than that among nonsmokers while clearly less than that among cigarette smokers.

4. Ex-cigarette smokers have lower death rates from COPD than do continuing smokers. The cessation of cigarette smoking is associated with improvement in ventilatory function and with decrease in pulmonary symptom prevalence.

- 5. Young, relatively asymptomatic, cigarette smokers show measurably altered ventilatory function when compared with non-smokers of the same age.
- 6. For the bulk of the population of the United States, the importance of cigarette smoking as a cause of COPD is much greater than that of atmospheric pollution or occupational exposure. However, exposure to excessive atmospheric pollution or dusty occupational materials, and cigarette smoking may act jointly to produce greater COPD morbidity and mortality.
- 7. The results of experiments in both animals and humans have demonstrated that the inhalation of cigarette smoke is associated with acute and chronic changes in ventilatory function and pulmonary histology. Cigarette smoking has been shown to alter the mechanism of pulmonary clearance and adversely affect ciliary function.
- 8. Pathological studies have shown that cigarette smokers who die of diseases other than COPD have histologic changes characteristic of COPD in the bronchial tree and pulmonary parenchyma more frequently than do nonsmokers.
- 9. Respiratory infections are more prevalent and severe among cigarette smokers, particularly heavy smokers, than among nonsmokers.
- 10. Cigarette smokers appear to develop postoperative pulmonary complications more frequently than nonsmokers.

In the 1964 report, cigarette smoking is "the most important of the causes of chronic bronchitis" and for pulmonary emphysema "it has not been established that the relationship is causal."

By 1967, the two conditions were combined and cigarette smoking becomes "the most important of the causes of chronic non-neoplastic bronchopulmonary diseases."

In the latest summary, which appeared in 1971, cigarette smoking is identified as "the most important cause of chronic obstructive bronchopulmonary disease."

The change in terminology should be noted - from a separate identification of chronic bronchitis and pulmonary emphysema, finally to the all-inclusive name of chronic obstructive bronchopulmonary disease. This last term is used also for morbidity and mortality statistics, although death rates for pulmonary emphysema as distinct from chronic bronchitis are available. The reviewer is now in the process of evaluating the significance of the prevalence of bronchitis and emphysema separately.

1967 Auerbach and his associates (2,0) tracheostomized to annual and, in an attempt to approximate human smoking more closely, exposed them to eigarette smoke through the trucheostomy tube. Five dogs died during this experiment and the remaining five were sacrificed after approximately 14 months of exposure. Other beagles were kept as controls; two had tracheostomy openings. These control dogs were sacrificed at the time the last five smoking dogs were sacrificed. Lungs of the dogs exposed to cigarette snoke showed microscopically the presence of diluted air spaces, especially beneath the pleural surface. Here the alveolar septa showed a fibrous thickening of the walls with areas of rupture and dilated air sacs. Padlike attachments to alveolar septa were found. These zones of connective tissue surrounding dilated air sacs were also visible macroscopically as white areas on the lung surface. There was no thickening of the walls of small arteries and arterioles within the lung. The lungs of the control dogs were normal in appearance with none of these changes. These abnormalities approximate but are not fully concordant with some of the typical pathological findings in human emphysema. This experiment does indicate that inhaled eigerette smoke apparently can damage the pulmonary parenchyma of dogs. Other findings (6) as yet unpublished, indicate that abnormalities of the bronchial epithelium resulted that approximate many of the histopathologic findings of human chronic bronchitis.

The investigations of Auerbach and his coworkers (15, 16, 8\$) have demonstrated by the use of both light and electron microscopy that dogs who inhale cigarette smoke through tracheostomas develop progressively more severe lesions of the bronchi and parenchyma with increased exposure to cigarette smoke. In electron microscopic studies of specimens taken from the lungs of dogs thus exposed to cigarette smoke, the following changes were observed: In 5 dogs sacrificed after only 44 days of smoking exposure, there was a proliferation of goblet cells as well as a partial loss of cilia in the lining cells, and in 5 dogs sacrificed after 420 days or more of exposure, the number of cell layers in the bronchial epithelium was found to be twice that of the nonsmoking dogs. Goblet cells and ciliated columnar cells were no longer present; instead, the surface was lined with columnar and cuboidal cells with stubby projections in place of cilia. Mitotic figures were frequently observed in the basal cells. These findings may be relevant to carcinogenesis as well as to the development of COPD.

In a long-term experiment, carried out by the same group, dogs were exposed to varying doses of cigarette smoke. Details of the experimental procedure have been outlined in the section on Pulmonary Carcinogenesis. The animals were separated into non-smoker, filter-tip cigarette, nonfilter-light, and nonfilter-heavy exposure groups. The dogs were "smoked" for 875 days, or approximately 29 months. The animals which died during the experiment and the animals sacrificed after day 875 were examined for pulmonary parenchymal changes as well as for bronchial epithelial alterations. As seen in figures 1 and 2, dose-related pathological changes, including fibrosis and emphysema, were found in the lung parenchyma of the exposed dogs. These changes were similar to those seen in the lungs of humans with COPD.

The experiments which failed to show that cigarette smoking does not produce pulmonary emphysema have not been mentioned in the reports. Some of the work from one laboratory has been quoted to describe details that do not relate to pulmonary emphysema directly:

1968 Aviado and his co-workers (2, 3, 4, 19, 38, 45, 46, 47, 48) have continued their studies on bronchoconstriction and bronchodilation in animals and recently have further investigated the role of histamine in a study of inhibitors for histamine decarboxylase in rabbits, dogs, and cats (39). These species have variations in response to cigarette smoking as previously noted. Cats have a uniphasic bronchoconstrictor response to inhaled eigarette smoke (somewhat like man's) and dogs have a biphasic response. Rabbits were observed to behave differently than cats or dogs. Histamine has been implicated as mediating part of the bronchoconstrictive effect of cigarette smoke. The rabbit does not respond to histamine by bronchoconstriction. This study (39) suggests that the rabbit lacks a histamine sensitive system in the airways, in contrast to cats and dogs. Alpha-hydrazino histidine, which inhibits the enzyme histamine decarboxylase, was demonstrated to prevent much of the bronchoconstrictive effect in cats and dogs. By analogy, this suggests the possibility that histamine may mediate some of the bronchoconstrictive response to inhaled tobacco smoke noted in humans. Pretreatment with atropine has been shown to block the bronchoconstriction caused by eigarette smoke (26) and by histamine inhalation in humans (7, 8, 52).

Aviado and coworkers have performed a series of experiments on live animals and in heart-lung preparations to study the effect of cigarette smoke on pulmonary physiology and structure (18, 19, 20, 21, 22, 179, 180, 199, 200, 201, 202). The authors observed that cigarette smoke causes acute bronchoconstriction both by the release of histamine and the stimulation of parasympathetic nerve pathways in the lung. Bronchial arterial injections of nicotine were found to cause reactions similar to those observed after cigarette smoke inhalation. The bronchoconstriction was usually followed by bronchodilatation which the authors attributed to sympathetic stimulation. As mentioned in the Chapter on Cardiovascular Diseases, nicotine has been shown to induce the release of catecholamines.

Experiments by Aviado and coworkers as well as other authors (66, 99) using guinea pigs showed that exposure to cigarette smoke was associated with increased bronchopulmonary resistance and decreased pulmonary compliance. The authors related these changes to the bronchoconstriction of terminal ventilatory units.

Similar experiments in dogs showed that the increase in resistance following either cigarette smoke exposure or intravenous nicotine could be blocked by pretreatment with atropine. As a parasympathetic blocker, atropine would decrease the acute bronchoconstrictive phase.

1973-Aviado and coworkers have studied the effects of hormones on the pulmonary response to cigarette smoke inhalation and intravenous nicotine injection. Subcutaneous progesterone administration, prior to nicotine or smoke exposure, reduced the bronchoconstrictor response in rats (64). A similar experiment involving pretreatment of dogs with glucocorticoids resulted in variable bronchoconstrictor responses after exposure to cigarette smoke (12).

creased numbers of enlarged air spaces and increased pulmonary resistance when compared with animals who underwent only

In the 1971 quotation, there is no mention that the attempt to induce emphysema in rats by exposure to cigarette smoke failed, although other procedures caused the lesions. The failure to elicit emphysema appeared in the conclusions of two articles which the documents overlooked.

#### Summary of Ito and Aviado (1968)

tracheal ligation.

Rats exposed to cigarette smoke daily for ten weeks did not develop signs of pulmonary emphysema. The rats subjected to experimental ligation of the trachea and intratracheal injection of papain developed signs of pulmonary emphysema but exposure to cigarette smoke did not uniformly influence the various signs. Functional residual capacity was not markedly elevated although histological examination showed a higher percentage of air spaces in emphysematous rats exposed to cigarette smoke. Cigarette smoke caused an increase in pulmonary resistance only in the rats with experimental emphysema. The increase in resistance is interpreted to mean an increase in bronchomotor tone. Isoproterenol injection induces bronchodilatation, more intensely in rats exposed to cigarette smoke.

#### Summary of Aviado et al. (1970)

This is the second reported attempt to demonstrate that eigarette smoke can induce pulmonary emphysema in rats. The negative results have been scrutinized to determine the significance of the changes in airway resistance and content of biogenic amines during exposure to effarette smoke. These changes in the lung are not important in determining the appearance or disappearance of pulmonary emphysema in rats, in the same animal species, tracheal ligation and endotracheat injection of papain produce pulmonary emphysema.

3. Evaluation of air pollution. The U.S. Public Health documents have arrived at the conclusion that cigarette smoking is the most important factor in the causation of chronic bronchitis and pulmonary emphysema. They have interpreted articles relating to air pollution and industrial hazards as indicating that these are less important than cigarette smoking. A re-examination of 48 articles indicates just the opposite conclusion made by the authors and details are outlined in Sections III and IV. One example is the report of Phelps and Koike (1962). The 1964 report has the following citation:

In the "Tokyo-Yokohama Asthma" studies, a severe asthma-like disease, presumed to be caused by air pollution, affected eigarette smokers predominantly (155). The absence of smoking data on unaffected members of the same population leaves the question of an additive effect of eigarette smoking unanswered. One study suggests that non-smokers may have a slightly greater prevalence of asthma than smokers; the possibility of bias due to self-selection of the base population could not, however, be excluded in this study (84).

Phelps and Koike concluded that air pollution is the cause of the disability. Removing the patients from the area caused a dramatic relief. The quotations are as follows:

The quotations are as follows:

As the disease progresses (usually in the patient with symptoms who remains in this area for several years or who returns to the area after having had the disease during a previous tour), more shortness of breath is manifested. The patient by this time is usually receiving adrenal steroids and has been frequently hospitalized. It is at this time that all of the symptoms and signs of emphysema are present. This is more likely to occur in persons who are heavy eigarette smokers. Although the study is by no means complete, it is believed that there is evidence that air pollution, of unknown content, is the cause of much disability which, under certain conditions, may progress rapidly to emphysema.

The follow-up studies revealed that those patients who were transferred to the United States early in the course of the disease had dramatic relief of their symptoms. All pulmonary function studies returned to normal within one to two months after the arrival of these patients in the United States. Their function studies took longer to return to normal than their symptoms. Those persons who had their disease for a long period of time (two to three years) took much longer to become asymptomatic, and it took even longer for their pulmonary function studies to become normal. Most of the patients were improved so far as coughing and wheezing were concerned, but they still had shortness of breath. The first sign of recovery appeared to be an increased response to IPPB with Isuprel.

Page			Reprint
162	(8)	FIESER L F: More on smoking and health. Chemistry 19: , 1964	817
164	(9)	KAMPMEIER R H: Smoking and noncancerous pulmonary disease.  Southern Med J 57:981-3, 1964.	818
165	(10)	EDITORIAL. West Virginia Med J 60:51-2, 1964	819
· 166	(11)	RAVENHOLT R T: Cigarette smoking: magnitude of the hazard.  Am J Public Health 54:1923-26, 1964.	820
167	(12)	WAKERLIN G E: Cigarette smoking and the role of the physician. <u>Circulation</u> 29:651-6, 1964.	821
168	(13)	FISHBEIM M: Medical progress in 1964.	822
169	(14)	U.S. DEPARTMENT OF HEALTH, EDUCATION AND WELFARE.  Three kinds of evidence. Public Health Service Publication 1103-F:1-2, 1965.	823
171	(15)	FOGARTY J E: Delivered to the Task Force on chronic bronchitis and emphysema. NTA Bulletin 3-6, 1967.	824
172	(16)	SHAMBAUGH G E: Additional data on smoking. Arch Otolaryng 85: 589-90, 1967.	825
173	(17)	WHATLEY J.L: Battle for breath. Today's Health 42-71, 1967.	826
. 174	(18)	ABELSON P H: Changing attitudes toward smoking. Science 161: 319, 1968.	827
175	(19)	DELARUE N C: Health hazards of smoking. Modern Med Canada 23:37-42, 1968.	828
176	(20)	HARKEN DE: Smoking The Masochistic massacre. Dis Chest 54: 445-9, 1968.	829
177	(21)	HUTCHISON G.B: The nature of epidemiologic evidence: smoking and health. Bull NY Acad Med 44:1471-5, 1968.	830
178	(22)	NYLAN L H: Air pollution and emphysema. Connecticut Med 32:706 -7, 1968	831
179	(23)	OCHSNER A: Prevention of executive obsolescence. J Am Geriatrics	Soc 832

age				Reprin
180	(24)	SCHUMAN L M: Epidemiology of smoking related diseases v		
· · · · · · · · · · · · · · · · · · ·		physicians encounter in their office practice. Dis Cl 54: 176-9, 1968.	<u>nest</u>	833
181	(25)	JACOBS E C: Smoking - insidious suicide and personal air p Military Med 135:678-81, 1970	ollution.	834
182	(26)	BENNETT J S: Smoking - time to practice what we preach.  Med Assoc J 106:276, 278, 1972.	Canad	835

You may have seen headlines about, the report of the Surgeon General's Advisory Committee on Smoking and Health (Chemistry, February 1961; page 2), but I doubt that you have waded through the 387-page document or the lengthy newspaper accounts of its findings. Since I was the only chemist on the 10-man advisory committee, I could hardly decline a request from the editor of Chemistry for a summary of the report addressed to Chemistry's readers. You may care to know what it all boils down to, so that you can make a wise decision for yourself and so that you can offer sound advice to others.

Countless weekend committee meetings were held in Washington in which we worked over a Friday-Saturday-Sunday stretch, continuing each day from \$:50 a.m. to about \$ p.m. One problem was that the medical men, who properly predominated on the committee, would not use a simple term such as "cancer of the mouth" when the more dignified medical term "carcinoma of the buccal cavity" was available.

The unanimous opinion of our committee was: "Cigaret smoking is a health hazard of sufficient importance in the United States to warrant appropriate remedial action."

A few of the findings on which this judgement was based are as follows:

• Cigaret smoking is causally related to lung cancer in men; the magnitude of the effect of eigaret smoking for outweighs all other facters. The data for women, though less extensive, point in the same direction.

• The risk of developing lung cancer increases with duration of smoking and the number of cigarets smoked per day and is diminished by discontinuing smoking.

• Cigaret smoking is the most important of the causes of chronic bronchitis and increases the risk of dying from chronic bronchitis and from pulmonary emphysema.

• Male cigaret smokers have a higher death rate from coronary artery disease than nonsmoking males.

The death rate for smakers of cigarets is about 70% higher than that for nonsmokers.

These are strong words and they call for strong actions. It is not yet clear what remedial actions will be instituted by governmental agencies, but there is no question about the course young people should take. Don't smoke eigencts. Don't let yourself fall into the habit of smoking. It is a habit which many people regret but find difficult to break. You young people are in a position to make a decision in advance and so save yourselves from concern about remedial measures at a later, critical stage.

1005051479

 $\alpha$ 

The author of this Editorial has combined the two pulmonary diseases to read "increases the risk of dying from chronic bronchitis and from pulmonary emphysema." In the 1964 statement, the two diseases were covered in separate sentences: "cigarette smoking is the most important of the causes of chronic bronchitis" and "a relationship exists between pulmonary emphysema and cigarette smoking but it has not been established that the relationship is causal."

I Though there have been numerous domonstrations of impairment of pulmonary function in heavy eigarette smokers as compared to control groups, the main evidence of the hazard of cigarette smoking in terms of chronic bronchitis and eventual emphysema comes from epidemiologic studies. Everyone recognizes that air pollution, in cities especially and in certain occupations, can not be divorced from inhaled cigarette smoke when considering the ill effects of inspired air. Yet, other than asthma and familial tendencies to emphysema, chronic cough and infection are accepted as important factors in the development of emphysema. No one denies the "smokers' cough" and few would deny the airway resistance which is part of inlialed smoke, both unquestioned potentials in the development of emphysema. The pathologic evidence of alterations in the bronchial mucosa in prolonged cigarette smoking is so overwhelming that few argue about this anymore. The symptoms of this chronic bronchitis as a role in emphysema need no more than a mention.

What it all adds up to, is that no one has been able to show unquestioned causal relationship between eigarette smoking and chronic bronchitis and/or empliysema even approaching that of its relationship to bronchogenic carcinoma. But no matter how good a case the apologist for tobacco may make for chronic infection and air pollutants other than cigarette smoke as a cause for bronchitis and for emphysema, the fact remains that eigarette smoke is a pollutant which in its own right can impair pulmonary function and therefore has a deleterious effect on pulmonary structures over and above the other unavoidable factors in the air of our environment and the bacterial flora of our respiratory, tracti. It would be rather nonsensical to discard as accidental increased mortality rates in heavy cigarette smokers over those of control groups of nonsmokers in epidemiologic studies of ebronic brouchitis and emphysema.

The application of epidemiologic studies to pulmonary emphysema and chronic bronchitis is mixed up with pathologic evidence. The prevalence of respiratory symptoms does not refer to incidence of these two separate disease entities. Although the 1964 statement has made a distinction between the influence of cigarette smoking to symptoms of bronchitis, others have interpreted the symptoms as indicating the positive presence of the direase.

1005051481

An association between eigarette smoking and chronic lung disease has been established beyond medical argument. Cigarette smoking accounts for the majority of cases of emphysema and emphysema deaths now outnumber tuberculosis deaths. It has been estimated that smoking ten eigarettes per day will lead to a diminution of four years inlife expectancy for the average American. Cancer of the lung is from 20 to 30 times more prevalent in the eigarette smoker than the nonsmoker. This is our most deadly cancer with a salvage rate of only 5 per cent. A study of British physicians shows that at age 35 the chances of dying within the next ten years are 1-23 for the heavy eigarette smoker and 1-90 for the nonsmoker.

The opening sentence refers to <u>association</u> between cigarette smoking and chronic lung disease but the second setence condemns smoking because it <u>accounts</u> for the majority of cases of emphysema. A causal relationship between cigarette smoking and emphysema has not been established, according to the 1964 report.

(11) RAVENHOLT R T: Cigarette smoking: magnitude of the hazard. Am J Public Health 54: 1923-26, 1964.

The censensus of findings of many studies indicate that approximately nine-tenths of lung cancer doubs in the United States today are caused as smaking; and this consistent fact may be used in conjunction with the above findings (Table 1) to achieve a rough measure of total tobacco disease mortality:

1	Deaths from lung cancer in total population	122	1
	Excess deaths from all	723	- 6
	causes among smokers		

Hence, excess mortality from all causes among smokers in an Americantype population is probably about six times greater than total lung cancer mortality in that population.

During 1962, 41,376 deaths were seribed to lung cancer in the United States 10; and by application of the lungcancer index of one-sixth, one can estimate that there were roughly a quarter million excess deaths due to smokingwhich was 14 per cent of all deaths (1,756,720) from all causes. The relative importance of tobacco despecially cigarettes) as a hazard to life in the United States is indicated by the following balance of causes of death. (Figure 1.) The sum of all deaths from accidents, injection, suicide, homicide, alcohol, and stomach cancer only equals the quarter-million deaths from tobacco disease in the United States during 1962.

American Balance of Hazards to Life, 1962

Quarter Million Deaths from Tobacco
*Lung Cancer 37,000
10ther Cancer Deaths 23,000 (Of laryax, bladder, etc.)
†Coronary Disease 119,000
Other Vascular Disease 35,000 (Strokes, etc.)
Other Tolsacco Diseases 36,000 (Emphysema, bronchitis, etc.)

97,139 40,594 1,254 5,161 49,920
40,804 1,254 5,161
,
93,745 9,506 66,999 15 41 17,183
20,207
9,013
<b>9</b> ,959
19,378

9 Non-traitie of all (41,376). lung cancer deaths, § Number estimated from lung cancer index, {Total europary forms deaths-\$27,581; total deaths from all cancer-\$1,750,770.)

This article illustrates how portions of the 1964 statement has been used to equalize "excess death among smokers" with hazards to life. The death rate from emphysema and bronchitis is multiplied by lung cancer index and those compared with deaths from selected causes.

(21) HUTCHISON G B: The nature of epidemiologic evidence: smoking and health. Bull NY Acad Med 44: 1471-5, 1968.

#### Epidemiologic Evidence

An extensive literature on associations between smoking and cardio-'vascular disease is now available. I propose to comment on some of the studies in this literature in which the unit of observation is a person and in which information is collected to characterize persons as to state of health. These are the studies which are commonly referred to as epidemiologic studies. When these studies are interpreted in terms of cause-effect relations the cause-effect concept is usually the pragmatic one.

The proof of cause-effect would be established or rejected by deliberately manipulating the smoking habits of such persons and observing the results. No such studies have been made to my knowledge. Controversy, when it occurs, rests on differences of opinion as to what constitutes proof in the absence of such a manipulative study. There are those who feel that proof is never established without such human experimentation. Others feel that proof can be established through observational studies, but there is little agreement as to what sorts of observations are required.

A number of schemes have been suggested for assuring such proof. Koch's postulates are an early attempt, but are applicable only when the causative event is exposure or nonexposure to living organisms. The surgeon general's committee has proposed a number of conditions which must be satisfied. These include consistency, strength, specificity, and coherence of the association. The ultimate decision, however, is said to be judgmental.

The discussion of epidemiology is followed by the conclusion that the evidence is adequate to support the cause and effect relationship between lung cancer and cigarette smoking. If the same principle is applied to pulmonary emphysema, a cause and effect relationship has not been established.

(22) NYLAN L H: Air pollution and emphysema: Connecticut Med 32: 706-7, 1968.

About 20,000 Americans will die of emphysenia this year, seven times the number that succumbed to it a decade ago. It disables one out of fourteen American workers over forty-five years of age. E. Cyler Hammond,! Vice President of the American Cancer Society told a senate subcommittee that a recent study of 126,000 men indicates that a combined atmosphere of eightette smoke and occupational pollution creates heigthened risk of emphysenia. A nonsmoker expend to occupational pollution (smog, pollen, dust, exhaust fumes or other irritants) has one chance in ninety of developing emphysema. Among heavy smokers expect to the same air pollution, one in seven develops this crippling disease. These findings are not new and are supported by those of others that air pollution and eigerette smoking often leads to destruction of. flung tissue that characterizes emphysema.

The risk of smoker exposed to air pollution cited by Hammond in his testimony is one of 10 possible figures. The other reports do not indicate such a disparity between smokers and nonsmokers.

Severe emphysema is a condition that can be considered worse than cancer. The patient with cancer of the lung either dies (only about 6 per cent can
be cured) or gets well, whereas the patient with severe emphysema has a
living death. According to the United States Public Health statistics, more
than 12 million people in the United States today are completely disabled by
severe emphysema caused directly by cigarette smoking. In 1965 in the United
States, there were 77 million man-days lost from work because of tobacco,
an additional 88 million man-days of sickness in bed as the result of use of
tobacco, and 360,000,000 man-days of restricted activity because of the effects
of tobacco. On the basis of these statistics, and using minimal figures of \$40 a
day for those totally disabled and \$20 a day for those partially disabled, the
loss of income from disability caused by tobacco in 1965 amounted to
\$15,800,000,000. The only reason that tobacco does not cause senescence more
often is that frequently it causes early death—the dubious advantage of being
spared early obsolescence.

The Surgeon General's report has been used to discuss tobacco and senescence. The author, a foremost surgeon, has accepted all the statistics and figures in 1967 document.

(24) SCHUMAN L M: Epidemiology of smoking related diseases which physicians encounter in their office practice. Dis Chest 54: 176-9, 1968.

Chronic bronchopulmonary diseases were cauly implicated in the 1964 Advisory Committee port to the Surgeon Ceneral. The evidence v deemed adequate for a causal relationship between eigarette smoking and chronic bronchitis and was strongly implicated as increasing the risk death from pulmonary emphysima. Despite \* recent attempts at more precise definitions of chr. le bronchitis and pulmonary emphysema, the staof such definitions over the immediate past in t own country and those in other countries mai trends and comparisons difficult if we attempt separate the two entities. Thus, mortality and ma bility analyses are better applied to a combination of the two-dironic bronchitis and/or emphysema. As a problem in mortality, chronic bronchitis and/ or emphysema presently claim half as many deaths as does lung cancer. However, in the 16-year peried, 1950-65, deaths from these two non-neoplastic pulmonary diseases have increased over six-fold from 3,157 deaths reported in 1950 to 23,432 rereported in 1905;2.5 This increase has been greater than the similarly appalling toil from lung cancer. The most dramatic rise has been for the emphysems component for which the age-adjusted mortality rates rose from 1.3 per 100,000 in 1950 to almost 130 in 1935. Deaths from chronic broachitis rose less spectacularly, only doubling in the same period. Although some of these increases must represent diagnostic improvement, it is difficult to ascertain the degree to which this operated. The causal/relationship with eigarette smoking and the use of eigenettes in this period would suggest that a major part of this rise in mortality must be real.

The explanation for combination of chronic browhitis and emphysema in mortality statistics is not a convincing one. Since 1964, the diagnosis of the two disease entities has advanced to the point of widespread application. It would be reasonable to analyze the death rates from the two diseases separately.

(12) WAKERLIN G E: Cigarette smoking and the role of the physician. Circulation 29: 651-6, 1964.

#### Smoking and Other Diseases

Epidemiologie, experimental, and clinical studies have demonstrated adverse relationships between eigenette smoking and lung cancer, chronic bronchitis, and emphysema. One of the complications of emphysema is cor pulmonale. Cigarette smoking has also been shown to be associated with an increased incidence of arterioscleresis of the pulmonary arterioles and small arteries.

Increased morbidity and mortality among cigarette smokers have been reported for certain other forms of cancer and for peptic ulcer.

#### Smoking and Over-all Death Rates

Cigarette smokers show adverse changes in structure and function of the lungs and bronchial tree. 15, 19 Pulmonary function changes include decreased vital capacity and decreased forced expiratory volume. 20 Some of these changes revert toward normal in those who stop so oking. 21

In addition to the greater risk of illness and death from disease, recent studies suggest a reduced sense of well-being among cigarette smokers. Thus, compared with absence of the eigarette habit, eigarette smoking is more frequently associated with cough, shortness of breath, decreased appetite, and increased nervous tension." An adverse association of eigarette smoking with body function appears to begin with fetal life, since more prenature infants are born to mothers who smoke eigarettes. 23.24

#### Recent Actions with Reference to Smoking

The statements are individually accurate. The implication to the reader is unavoidable: pulmonary effects of cigarette smoking is responsible for deaths which in turn has led to recent actions against smoking.

The rise in cancer of the lung was associated with a considerable increase in emphysema in many parts of the world, particularly in the U.S., Canada, and England. One survey indicated that about 25% of men over 45 have some degree of emphysema. The condition is from seven to 13 times more frequent among cigarette smokers than among nonsmokers. Both cigarettes and air pollution have been considered as causative factors. Great attention is now focused on determining the exact relationships among these factors and diseases.

Looking back at 1964, a few peaks of interest become apparent in each area of medical research and medical practice. Heart disease, cancer, and high blood pressure with stroke received major medical attention as the leading causes of death. Infectious diseases were controllable as never before, and more of the infectious diseases of childhood became wholly amenable to prevention and cure.

The survey showing that 25% of men over 45 have some degree of emphysema refers to prevalence of coughing and expectoration. The whoted 7 to 13 times more frequent among cigarette smokers is also based on incidence of respiratory symptoms. The mortality ratio for pulmonary emphysema among smokers compared to nonsmokers is about 3 times.

#### SURGEON GENERAL'S COMMITTEE

By 1962 scientific research had implicated the cigarette in several major diseases. In addition to chronic bronchitis and emphysema, these included heart disease and lung cancer. That year, the Surgeon General of the Public Health Service appointed an advisory committee of 10 eminent scientists to review the accumulated evidence. The committee had advice and consultation from nearly 200 specialists representing every related field of interest.

#### THREE KINDS OF EVIDENCE

The Surgeon General's Committee reviewed three kinds of research evidence.

#### 1. Animal Experiments

Animals exposed to tobacco smoke show severe damage to the cells lining the air tubes—the same kind of damage seen in heavy smokers. (A pack a day or more is considered "heavy smoking.")

#### 2. Reman Tissue

When the cells of the bronchi of smokers and nonsmokers are compared, the smokers cells show characteristic damage: loss of the cilia and irregular shapes. These changes have been noted in thousands of studies, both in living patients and at autopsy.

#### 3. Population Studies

Extensive and repeated studies of large numbers of people have consistently show that cigarette smokers are more likely to have symptoms of respiratory disease than non-smokers. They cough more, bring up more phlegm, and complain more of breathlessness.

#### SMOKING AND HEALTH REPORT

In January 1964, the Surgeon General's Committee finished the most comprehensive study of smoking and health ever undertaken. Studies involving more than a million men had been reviewed. For cigarette smokers, the death rate from chronic bronchitis and emphysema was found to be six times greater than for nonsmokers. A 500-percent difference! Ex-smokers had a death rate from these diseases about 40 percent greater than for nonsmokers. Men who began smoking before the age 20 had a higher death rate than those who started later.

# Commentary (14) U.S. Department of Health, Education and Welfare, Cont.

This widely distributed leaflet contains selected statements, each one quoted from the 1964 report. The combination of the three kinds of evidence do not relate to chronic bronchitis or pulmonary emphysema mentioned above and below the quotation. The animals were not sick with emphysema or bronchitis, the human tissue was obtained from smokers who were not suffering from emphysema or bronchitis, and the population studies surveyed respiratory symptoms which do not necessarily signify emphysema or bronchitis.

(15) FOGARTY J E: Delivered to the Task Force on chronic bronchitis and emphysema. NTA Bulletin 3-6, 1967.

What is the cause-or should I say causes-of emphysema, the condition that begins innocently enough with a mild cough and seems to proceed to increasing shortness of breath until the normal activities of the day become increasingly burdensome?

Is cigarette smoking a factor? We are told in the report of the Surgeon General's Advisory Committee on Smoking and Health that cigarette smoking is an important cause of chronic bronchitis and that a "relationship" exists between cigarette smoking and emphysema. Has that relationship as yet been defined? Despite the accumulating evidence that cigarette smoking and emphysema are intimately linked, our society is heedlessly smoking six per cent more cigarettes a year than in 1964, when the report of the Surgeon General's committee was released.

What about air pollution? In all honesty, we do not need experts to tell us that air pollution is an insult to our respiratory tract, yet until now our national complacency has permitted the continued spewing of poisons into the air.

The relationship of cigarette smoking to chronic bronchitis and pulmonary emphysema is differentiated in this article, according to the 1964 report. The term cause is unfortunate; risk factor would have been more appropriate.

7 million

(16) SHAMBAUGH G E: Additional data on smoking. Arch Otolaryng 85: 589-90 1967.

THE NATIONAL Center for Health Staistics in October 1986 published the Mortality Statistics From Diseases Associated With Smoking, from 1950 to 1964. Two groups of liseases are considered: those causally related to smoking, namely cancer of the lung. ancer of the larynx, cancer of the lip, and hronic branchitis, and those associated with smoking but where the evidence is not yet ufficient to establish a causal relationship. The latter group comprises arteriosclerotic heart disease including coronary disease, cirthosis of the liver, emphysema, ulcer of the stomach, cancer of the esophagus, cancer of the oral cavity other than the lip, cancer of the bladder and other urinary organs, and noncoronary cardiovascular disease. JAMA (199:24 [Jan 23]: 1967): carries an interesting study of cigarette smoking in dogs. After more than 420 days of smoking through a tracheostomy tube there was marked cardiac enlargement compared to controls, and pulmonary abrosis and emphysema in all the smoking dogs, not found in any of the controls. Interestingly some of the dogs showed a liking of cigarette smoking as indicated by tail-wagging and jumping into the smoking box voluntarily. With this evidence perhaps emphysema should be reclassified as causally related to smoking. rather than associated with smoking.

The widely publicized "smoking dogs" has undoubtedly led to the reversal of the statement from no causal to causal relationship between cigarette smoking and pulmonary emphysema. There are other attempts in animals which failed to show a cause and effect relationship but these were never publicized.

Emphysema and chronic bronchitis are the fastest rising causes of death in the country. Almost 20,000 people will die this year from emphysema or its complications. Whether this increase is due to better methods of diagnosis, or to the greater longevity of our population, we do not know. Certainly, part of the increase is due to modern environmental conditions (exhaust fumes, smog, cigarette smoke, etc.).

Emphysema is about 13 times more prevalent among cigarette smokers than among non-smokers, but many who have never smoked contract the disease. Men develop emphysema five to 10 times more frequently than women, probably because women are "abdominal breathers" (they use the diaphragm and abdominal muscles more), whereas men are "chest breathers" (they use the chest muscles more).

At present, we don't know the exact cause of pulmonary emphysema, but we do know quite a bit about it. We know that it is an overinflation of the lung due to loss of that organ's natural elasticity. The smaller airways (bronchial tubes) narrow from spasm, irritation, infection, or a combination of the three. This narrowing traps the air in the small air sacs (alveoli), and they dilate and lose their elasticity, or ability to recoil.

The 13 times prevalence of emphysema among cigarette smokers is based on incidence of respiratory symptoms which does not necessarily signify the disease. The author avoids stating that emphysema is caused by cigarette smoking. Even though the author does not know the exact cause of pulmonary emphysema, the hurried reader will undoubtedly regard the 13 times prevalence of cigarette smoking as a causative factor.

(18) ABELSON P H: Changing attitudes toward smoking. Science 161: 319, 1968.

The attitudes of the people of the United States toward eigerette smoking are undergoing a major change. In the 30-year period prior to 1964, per capita consumption of eigerettes tripled. During the last 4 years, per capita consumption of eigerettes has leveled off, and currently it may be dropping. The 1964 Surgeon General's Advisory Committee report entitled "Smoking and Health" was a major milestone. The report set in motion developments that ultimately will have a profound effect. One consequence was the stimulation of research bearing upon the effects of smoking. A recent compendium lists 364 projects in 36 states, the District of Columbia, and 25 foreign countries.

Out of this research activity has come substantial evidence that the effects of smoking are even more serious than was stated in the 1964 report. Highlights of the findings have been presented in a new U.S. Public Health Service document. The most striking finding is that "the life expectancy for a two-pack a day, or more, smoker at age 25 is 8.3 years less than [that for] the corresponding non-smoker."

The 1964 report probably has influenced the smoking habits of physicians. At one time many of them were heavy smokers. A recent surveyt indicates that 100,000 physicians have stopped smoking cigarettes. Of all the physicians surveyed, 35 percent had never smoked, 36 percent had smoked at one time but had stopped, and only 29 percent were still smoking. In contrast, 52 percent of the general adult male population currently smoke cigarettes.

The survey also indicates that physicians are deeply concerned over the effects of smoking. More than 90 percent stated that they associate smoking with chronic bronchitis, lung cancer, and emphysema, and almost as many associate it with peripheral vascular disease and coronary artery disease. The responses of physicians who have stopped smoking are revealing. When asked on the survey questionnaire why they stopped, 60 percent checked "Protect my future health," 47 percent checked "Occurrence of certain symptoms," and 43 percent checked "Scientific reports convinced me."

Following the appearance of 1967 Report, a second wave of Editorials appeared. The smoking habits of physicians are discussed here. It is important to point out that 43% who stopped smoking were convinced of the scientific reports summarized in the U.S. Public Health documents.

2. Chronic Bronchitis and Emphysema.

There are now 80,000 annual deaths in the United States from these diseases and although they both may develop in non-smokers, the risk of death in cigarette smokers is increased from eight to 15 times as compared to, the non-smoking population. In the five-year period between 1959 and 1964 deaths from chronic bronchitis increased 90 per cent and over the same period, deaths from emphysema increased 104 per cent. There were 421,000 new cases of bronchitis recorded in 1965, and 267,000 new cases of emphysema defined in the same period. At that time, it was estimated that there were more than three million cases of chronic bronchitis in the United States.


This special article appeared in a Canadian journal. It can be noted that United States statistics are quoted here, even though there is information directly from Canada. Other countries have relied on the U.S. Public Health documents for statistics relating to chronic bronchitis, although its incidence varies from country to country.

The combination of seven prospective studies to seals that the mortality rates of cigarette smoken over non-smokers for all causes of death was 7 per cent higher in smokers.

Let us look at some cold facts on mortality in a combination of seven prospective mortality records of smokers versus non-smokers:

Cancer of lung	(10.8)	1000 per cent (lead- ing cancer in men)		
Bronchitis and		•		
emphysema	(61)	600 per cent (and leading cause of		
•	-	disability)		
Cancer of the		All Light Add Art 1		
brynz	(5.4)	•		
Oral cancer	(4.1)			
Cancer of the	` '			
esophagus	(34)			
Peptic ulcer	(27)			
Atherosclerotic	• ,			
discuso	(26)			
Coronary artery	• - /			
discuse	(17)	70 per cent		
•		1 <b>6</b>		

Although death rates of chronic bronchitis separately from pulmonary emphysema are available, the 1967 report combines the two diseases, presumably to improve the statistical significance. Reference to the mortality ratio for smokers to nonsmokers appearing in articles such as the one under consideration do not question the practice.



#### Facts

Smoking remains the foremost preventable cause of disease, disability and death in the aited States. 1-13 Smoking causes coronary artery disease, lung cancer, emphysema, thrombo-angiitis obliterans and the like, annually producing hundreds of thousands of premature deaths and many hundreds of thousands of cripples. Tobacco diseases are causing many millions of man-days lost annually in this country. Tobacco diseases can shorten the smoker's life by eight or more years, 12 can cause a death every 1.5 minutes, 13 and cost the Nation between \$4 billion and \$20 billion annually. 13

The statements in the U.S. Public Health documents are abbreviated in this short paragraph. It can be noted that smoking is stated to <u>cause pulmonary</u> emphysema. The phrase "no causal relationship has not been established" has been omitted.

Health by the Royal College of Physicians (London) was discounted by the tobacco industry, as contributing Little but "statistical association". The College's new report "Smoking and Health Now" (1971) points out that the criticism of "statistical evidence only" cannot be upheld as a valid objection. Much of actiology, morbidity incidence, etc., on human disease is based on statistical evidence: the pro-smoking faction suggests that, as far as smoking is concorned, no laboratory evidence is evailable to show its responsibility for lung cunter. However, the recent work of Auerbach and Hammond on invasive lung cuncer in dogs should convince the Doubting Thomas. Statements from the U.S. Department of Health, Education and Welfare (1971) are:

- 1. Cigarette smoking is the major cause of lung cancer.
- Cigarette smaking is the most important cause of chronic obstructive broncho-pulmonary disease, principally chronic bronchitis and pulmonary emphysema.
- J. Cigarette smoking is an important risk factor in the development of coronary heart disease and, by accelerating damage already present as a result of coronary heart disease, may contribute to sudden death.

Isn't it time that the medical profession became a little more active in the efforts to reduce one of the major health hazards of our time. We tend to get worked up about pollution but do little about one of the greatest contaminants of all the tobacco smoker. 1005051499

The 1971 U. S. Public Health document and the 1971 report from the Royal College of Physicians are cited in this article. The widely publicized smoking dogs are cited as the most important laboratory evidence to support the statement that cigarette smoking is the major cause of lung cancer and the most important cause of chronic bronchitis and pulmonary emphysema.

### VIII-C. PUBLICITY RELATING TO SMOKING AND HEALTH

The widespread distribution of the seven documents on "Smoking and Health" has led to the appearance of secondary publications. Almost all of them accept the conclusions appearing in the seven documents that cigarette smoking is the most important cause of chronic bronchitis and pulmonary emphysema. A few articles contest the statement but none has appeared in recent years. The articles are listed in the following order:

ilmonary	emphysema. A few articles contest the statement but none h	as
opeared i	in recent years. The articles are listed in the following order	: Page
C-(1).		rage
) 	statement. (total number - 62)	184
C-(2).	Articles relating to pulmonary emphysema.	
	(total number - 52)	187
C-(3).	Articles relating to chronic bronchitis. (total number-17)	190
C-(4).	Articles relating to mortality statistics.	
	(total number - 38)	191
C-(5).	Articles relating to morbidity statistics.	
	(total number - 25)	193
C-(6).	Articles relating to bronchopulmonary and cardiopulmonary	
•	effects of cigarette smoking. (total number - 14)	195
C-(7).	Antismoking campaign. (total number - 95)	196

C-(8). Arguments against Surgeon General's statement.

(total number - 21)

		Reprin
America	n Cancer Society: 1968 cancer facts and figures. pp 1-32, 1967.	836
	Answers to cigarettes quiz.	837
ANON:	U.S. report reaffirms cigarets are hazardous.	838
ANON:	Women face hazards of smoking. p. 44	839
ANON:	World conference on smoking.	840
ВЕСКЕ	RMAN S.C: The health consequences of smoking. J. Maine Med Assoc. 58: 222, 1967.	841
	CT: Pusenje, bolest i zdravstveno prosvjecivanje. <u>Lijec Vjesn</u> 86: 1409-11, 1964.	842
British	Medical Journal: Menace of smoking. Br Med J 3: 753-54, 1967.	843
British :	Medical Journal: The smoking disease. 1 (5740):61-62, 1971.	844
	ING R H: The physician and the cigarette. Education concerning dangers of smoking essential to render habit socially unacceptable.  RI Med J 50: 735-56, 1967.	. 845
Bulletin	of the Los Angeles County Medical Association: The health consequences of smoking. A public health review: 1967. Bull Los Angeles Co Med Association: Bull Los Angeles Co Med Association: Bull Los Angeles Co Med Association: 98: 1-2, 57-58, 1968.	g 846
CRALL	EY L J: Presidents page. Am Ind Hyg Assoc J 30: 2, 1969.	847
DAVIS I	A L: Current facts concerning smoking and pulmonary disease and cardiovascular disorders. Memphis Mid-South Med J 73-75, 1966.	<b>84</b> 8
DEODH	AR S D: Smoking and health. Indian J Med Sci 26: 327-37, 1972.	849
DIEHL	HS: Cancer 110.	850
DIEHL	H S: Opportunity unlimited. Min Med 699-701, 1969.	851
DOYLE	JT: Tobacco and heart disease. Postgrad Med 44: 188-92, 1968.	852
El Dia l	Medico: La costumbre de fumar. El Dia Med. 265, 1967.	853
EVANG	K: Sigarettoyking og helse. Pp 300-4, 1964.	854
EVANG	K: Tupakointi terveyden vaarana. Suom Laakarilehti 24: 3108-11, 1969.	855
FLETC	HER C.M: Cigarettes and respiratory disease. NTRDA Bull 12-14, 1968.	856


		- 27	
	. •	_	Bibliography VIII-C-(I) page 185 Reprin
Artis Sin Society			FLETCHER C M: Cigarettes and respiratory disease. Inhalation Ther 14: 45-50, 1969.
	•	•	GARCIA ECHEVERRIA J.P. Algo mas sobre el habito de fumar.  pp 1563-66.  858
		ز آن الوخيل ج	GODBER G: The case against smoking. pp 20-25. 859
			GUTHRIE E H: Smoking and healthone year late, J Sch Health 53-57.
			HAMMOND E C: The multiple health hazards of cigarette smoking.  Presented at: 92nd Annual Meeting of the American Public  Healt Association. pp 1-26, 1964.  861
		"信"	HAMMOND E C, BERGLUND L and KASLOW R: Smoking habits and disease.  Minn Med 44.  862
			HOFFMANN F P: Smoking 110-11, 1967
	•		Journal of Iowa Medical Society: The month in Washington, J Iowa Med Soc 58: 918, 21, 1968.
46			KOLLER S: Lebensverkurzung durch lungenkrebs und kreislaufschaden als folge des zigarettenrauchens. DA 4: 191-206, 1968.
			LISTER J: Smoking and health now. N Engl J Med 284: 656-57, 1971. 866  LUNDAR J: Tobakksroking og lungesykdommer. Tidsskr Nor Laegeforen 85: 437-42, 1965. 867
	O		McMILLEN S I: Coronary and cancer by the carton. In: Cancer by the Carton.  Fleming H Revell Company: Westwood, NJ pp 11-22.
		ينم. الرقي	McMILLEN S I: The case against cigarettes: some summaries. In: Cancer by the Carton. Fleming H Revell Company: Westwood NJ pp 42-59.
			Medical News: Emphysema and smoking link new evidence. Med News 1969.
			Medical Tribune: Dr. Terry warns of cigarette danger.
	• •		Medical Tribune: Women smokers: effects near those in men or greater.  Med Tribune 13 (7): 1, 24, 1972.  872
granit na	•	·	Medical Tribune: W. Coast survey of doctors shows 39% still smoke.  873
	٠.•		Medical World News: How goes the war on smoking? Med World News 46-51, 1969. 874
er oan Gestaar		•••	Medical World News: PHS heats up warfare on smoking. Med World News 29-30, 1967.
M€ රුදා		•	. Medicina: Salud y tabaco. Medicina 28: 413-14, 1968. 876
	É		MERRILL M H: States take action to combat smoking. Johper 10, 1965. 877

Bibliography VIII-C-(1)	page 186	Reprint
MEURMAN L: Tobaksronkning forgiftning eller stimulans?		878
the gradient of the second		
MURRAY D: Our fastest-growing health menace. Readers Digest 90: 111-14, 1967.		879
OCHSNER A: Health hazards from tobacco. 267.	81.	880
PHILLIPS A J: The relationship between smoking and health a review of the evidence. Can J Public Health 55: 12-15, 1964.	<b></b>	881
POPPIUS H: Klinisktfysiologiska studier. <u>Lakartidningen</u> 67: 397-401,		882
	•	• See 2
Public Health (London): Special report: eight years after. Public Health 85: 133-36, 1971.		883
RANDHAW I H: Smoking and its injurious effects on health. 13: 187-93, 1963.		884
RASTELLI M: Tabacco e salute. Clin Ter 49: 85, 1969.		885
ROSENBLUM L: Experts discuss smoking. 14, 19	64.	886
Royal Society of Health Journal: Smoking and health now. R Soc Health J 91: , 1971.		887
SCOTT R B and WALKER R M: Smoking and respiratory disease. In:  The Medical Annual, John Wright and Sons Ltd, Bristol, 1965, pp 380-83.		888
TERRY L L: Cigarette smoking and respiratory disease. Presented at: Albert Einstein Medical Seminar. pp 1-23, 1968.		889
The Medical Journal of Australia. Oh. Mr. Snedden! 1315-6, 1971.		890
U.S. Department of Health, Education and Welfare: Cigarette smoking and health characteristics. United States July 1964-June 1965.		
National Center for Health Statistics pp 1-65, 1967.		. <b>891</b>
VINCENT R.G.: A fence or an ambulance. J. Sch. Health 37: 369-72, 1		892
U.S. Medicine: New report on smoking provides ammo for anti-tobacco in U.S. Med 30, 1969.	orces.	893
VOORST VADER PJAV: Bertrekkingen tussen roken en de strefte in Ne gedurende 1963. Ned Tijdschr Geneeskd 109: 601-8, 1965.	derland	894
WARE G W: PHS Report on smoking and cancer. Med Ann DC 33: 120	, 1964.	. 895
WEISS W: Cigarette smoking: suicide in slow motion. Arch Environ Heal 16: 1-3, 1968.	<u>th</u>	896
WYNDER E L and HOFFMANN D: Less harmful ways of smoking. J Na	tl Cancer I	nst

48: 1749-58, 1972.

19: 612, 1969.

Bibliography VIII-C-(2) page 188	Reprint
Journal of the Air Pollution Control: Institutional notes. J Air Pollut Control As 16: 392, 1966.	915
Journal of the American Medical Association: Emphysema: several factors may cloud diagnosis. JAMA 193: 24, 1965.	916
Journal of the American Medical Association: Panelists discuss emphysema's etiology, detection, treatment. JAMA 197: 20-21, 1966.	917
Journal of the American Medical Association: US report of rare syndrome.  JAMA 199: 35, 1967.	918
Journal of the American Medical Association: Emphysema warning indicated.  JAMA 208: 2255, 65, 1969.	919
LARSON R K: Smoking and emphysema. Calif Sch Health 4-11, 1966.	920
LAURELL C B: Is emphysema in alpha -antitrypsin deficiency a result of autodigestion? Scand J Clin Lab Invest 28: 1-3, 1971.	921
Medical Tribune: Cigarette smoking is found common to six families reporting emphysema. Med Tribune 24, 1966.	922
Medical Tribune: Cancer from cigarettes. Smoking dogs get invasive type tumor. Med Tribune 1, 20, 1970.	923
Medical Tribune: Smoking is held top habit factor in emphysema. Med Tribune 1963.	924
Medical Tribune: Cigarette-cancer dog tests challenged by tobacco group.  Med Tribune 1970.	925
Medical Tribune: Emphysema: Only one thing sure ailment is on rise.  Med Tribune	926
Medical Tribune: Lung emphysema: histologic trials only certain clue.  Med Tribune	927
Medical World News: Alarming rise in emphysema: Challenge to physicians.  Med World News 32-40, 1965.	928
Medical World News: Smoking plus air pollution multiphy emphysema rate.  Med World News 8: 29, 1967.	929
Medical World News: Who are the 5% whose genes invite emphysema?  Med World News 11: 15, 1970.	930
Nebraska State Medical Journal: Chronic bronchitis and emphysema.  Nebr State Med J 53: 446-47, 1968.	931
OSWALD N C: BCG vaccination. Practitioner 201: 701, 1968.	932
PANZER F: Emphysema and smoking. N Engl J Med 286: 1321, 1972.	933
PAUL O: Smoking and youth. Ind Med Surg 287-88, 1968.	934
PETERS G A and PRICKMAN L E: Smoking is detrimental in asthma and emphysema. Minn. Med 40: 116-20, 1957.	<b>9</b> 35
Physical Therapy: Emphysema now major killer of women. Smoking scored.  Phys Ther 47: 238-39, 1967.	936



# . VIII-C-(3). ARTICLES RELATING CHRONIC BRONCHITIS

	Reprin
British Medical Journal: Beginnings of bronchitis. 2: 190-91, 1970.	950
CARNOW B W: Pulmonary disease and air pollution. Chicago Med 71: 581-86, 1968.	951
DOLL R: Investigation into cigarette smoking and atmospheric pollution in the aetiology of lung cancer. Methods Inf Med 2: 13-19, 1963.	952
ESPINO V J: Tabaquismo y corazon. Arch Inst Cardiol Mex 37: 539-40, 1967.	953
GOLDSMITH JR: Air pollution medical research. Science 154: 1588-91, 1966.	954
GOLDSMITH JR: Synthesis, perspective, and future applications. Arch Environ Health 18: 710-14, 1969.	955
GREGG I: Chronic bronchitis and occupation. Br Med J 1: 675-76, 1966.	956
Journal of the American Medical Association: A causal role for bronchitis? 193: 26-27, 1965.	957
La Presse Medicale: Pollution atmospherique, tabac et bronchite chronique. 78: 1072, 1970.	<b>9</b> 58
Medical Times: Human costs of polluted air. 98: 156-70, 1970.	<b>9</b> 59
MITCHELL RS: Describes ramifications of smoking. Mich Med. 550-51, 1966.	<b>9</b> 60
OCHSNER A: The health menace of tobacco. Am Sci 59: 246-52, 1971.	961
STURGIS K R: Is surgery for lung cancer worthwhile?	962
	963
TAYLOR I: The physician and air pollution. Pa Med 74: 39-40, 1971.  Z31-36, 1963.	964
The Medical Officer: The "chesty" child, chronic bronchitis, and cigarettes. 161, 1969.	965
U.S. Medicine: Air pollution held major U.S. problem. 2, 17-18, 1968.	. • 966 ∢
	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1

#### BIBLIOGRAPHY

## VIII-C-(4). ARTICLES RELATING TO MORTALITY STATISTICS

	Reprint
American Cancer Society News Service Release: Film on "Smoking: Past and	
present" traces the history of smoking. 189, 1968.	967
Asian Medical Journal: Smokers' chance for lung cancer said II times higher.	968
	. ,
BARCLAY RS: Worldwide chest medicine. Chest medicine and cardiovascular conditions in Great Britain. Dis Chest 49: 559-60, 1966.	<b>9</b> 69
BORGATTA EF and EVANS RR: Smoking, health, and behavior. Aldine Publishing Company: Chicago, 1968.	970
BURCH G E: Trends in the incidence of disease in the United States.  World Med J 8: 439-40, 1961.	971
BURGESS A M: Mortality from diseases associated with smoking: United States 1950-1964. RI Med J 192-93.	972
Canadian Journal of Public Health: News notes: United States, 56: 453-54, 1965.	973
Canadian Medical Association Journal: The annual meeting of the Manitoba Medical	
Association (Canadian Medical Association Manitoba Division), 91: 1083, 1964.	974
Canadian Medical Association Journal: Intelligent debate or demented fiddling? 96: 679-80, 1967.	975
Cancer News: New ACS report traces smoking effects in women. 18	976
CECILIONI V A: Lung cancer in Hamilton. Can Family Physician 1970.	977
FONTANA R S: Smoking and health.	978
FLETCHER C M: Cigarettes and respiratory disease. NTRDA Bulletin 12-14, 1968.	<b>9</b> 79
GARFINKEL L: Several studies show higher death rates for smokers than for nonsmokers. Public Health News 211-14, 1963.	980
HAMMOND E C: Some preliminary findings on physical complaints from a prospective study of 1,064,004 men and women. Am J Public Health 54: 11-23, 1964.	981
HAMMOND E C: Competitive risks: lung cancer, emphysema and coronary heart disease. Cancer Soc News Serv 1-10, 1967.	982
HAMMOND E C: La relacion entre el habito de fumar y las tasas de mortalidad.  Orientacion Med 20: 985-92, 1971.	983
HELMHOLZ H F Jr: On diagnostic errors. Ann Allergy 26: 517-18, 1968.	984
HIGGINS ITT: Smoking. In: Aging of the Lung. Edited by L Cander and J H Moyer, Grune and Stratton: New York, pp 227-38.	• 985

Bibliography VIII-C-(4)	Reprint 3
HYATT CL: Physicians and smoking. J Arkansas Med Soc 184.	986
.JONGKEES L B W: Sigaretten en spinazie. Ned Tijdschr Geneeskd 109: 530, 1965.	987
Journal of the American Medical Association: Cigarette smoking may cause small hearing loss, 210: 1845, 1969.	988
KLEIN R C: A hole in the whole case. Med Counterpoint 61, 1970.	989
LANDAU E and MORTON J: An epidemiologic view of chronic pulmonary insufficiency in the United States. Am Rev Resp Dis 83: 405-7, 1961.	990
L'Union Medicale du Canada: Augmentation du nombre de cancers pulmonaires.	
97: 1850, 1968.	991
McKEE K T: The occurrence and progression of pulmonary emphysema.  South Med J 56: 129-32, 1963.	992
Medical Bulletin: Warning: cigarettes are dangerous to your health. 45-47, 1968.	993
Medical Bulletin on Tobacco: Heart deaths lead cigarette effects: 5: 1967.	<b>9</b> 94
Medical Tribune: Lung cancer seen often in smokers in Czechoslovakia.	995
Medical Tribune: Firm ties denied between cancer and air pollution. 8-	996
Medical World News: Bronchitis and emphysema increase in female smokers.  35, 1967.	997
MITCHELL R S: Outlook in emphysema and chronic bronchitis. N Engl J Med 280: 445-46, 1969.	998
Public Health Service Publication No. 1103: Cigarette smoking and cancer. 1963, 1-32.	<b>9</b> 99
RAVENHOLT R T: Cigarette smoking: magnitude of the hazard. CA 187-88, 1965.	1000
RENZETTI A D Jr, McCLEMENT J H and LITT B D: Respiratory diseases.  Nebr State Med J 488-89, 1966.	1001
Nebr State Med J 488-89, 1966.  RYDER C F: Chronic respiratory diseases a mushrooming challenge.  JAMWA 21: 132-36, 1966.  The Journal of School Health: Cigarette smoking and cancer. A new booklet by the American Cancer Society. 266-68.	1002
The Journal of School Health: Cigarette smoking and cancer. A new booklet by the American Cancer Society. 266-68.	1003
WADDELL JA: Bronchitis and emphysema. Br J Clin Pract 20: 477-80, 1966.	1004

·····································	Reprint
ANDERSON K N: Doctor with one foot in space. Today's Health	
32-35, 66-69, 1964.	1005
솔루바리 4	
BOIN J K: Expert MD opinions vary on major health hazard facing coal miners.	
<u>U.S. Med</u> 4, 18, 1969.	1006
BOU COT KR: Smoking as a predisposing factor in respiratory disease.	7007
Arch Environ Health 16: 459, 1968.	1007
Canadian Medical Association Journal: Smekers who cough. 89: 1099-1100,	
1963.	1008
Canadian Medical Association Journal: The lung. 100: 349-51, 1969	1009
- · · · · · · · · · · · · · · · · · · ·	
CHRISTINE B W: Effects of cigarette smoking on Connecticut residents.	
Conn Health Bull 85: 23-28, 1971.	1010
College Health: The association presents its views about smoking.	1011
<b>2-3, 1965</b>	1011
FERRIS B G Jr, SPEIZER F E, WORCESTER J and CHEN H Y: Adult mortality	,
in Berlin, NH, from 1961 to 1967. Arch Environ Health 23: 434-39,	
1971	1012
FORD R M: Smoking, lung cancer and asthma. Br Med J 3: 372, 1971	1013
HEPPER N G G: Cigarette smoking and chronic respiratory disease.	1014
Minn Med 52: 1373-75, 1969.	1014
HORN D: Cigarette smoking in the high schools. Calif Sch Health	•
17-21, 1966	1015
HORTON R J M: Discussion. Epidemiologic studies of chronic respiratory	
diseases. Arch Environ Health 10: 344-45, 1965	1016
JOULES H:: A preventive approach to common diseases of the lung. Br Med J	1017
<b>2:</b> 1259-63, 1954	1011
Journal of Iowa Medical Society: The teen-age smoker. 220, 1969	1018
The term and the second of the	
Journal of the national Medical Association: Statistics on health and cigarette	•
smoking. 356, 1967.	1019
•	
LORENZONI L: <u>Med Lav</u> 58: 769-71, 1967	1020
MASON R P: Cancer and the environment. 111, 1970	1021
MASSON KI. Cancel and the environment.	
Medical News: Emphysema worse out of town. 1968	1022
	n 6 304
Medical World News: Doctors push black lung fight to surface. 10:	
14-16, 1969	1023
MITCHELL R S, MAISEL J C, DART G A and SILVERS G W: The accuracy	
of the death certificate in reporting cause of death in adults.	
Am Rev Resp Dis 104: 844-50, 1971.	1024
<del></del>	**

page 194

		•	77.	· · · · · ·					
•									•
		•						en e	
						•			
				The second				- ged	
								- AB	
			•						and the second
	•					e garanta antara da santa antar ♣			The Mark Co.
	•	•					i e e e e e e e e e e e e e e e e e e e		
				•	•			E Tomas de la companya de la company	

Bibliography VIII-C-(5)

disease.

in smokers.

86: 47, 1964

. Public Health Reports 78: 743-48, 1963

16, 1968

Natl Cancer Inst Monogr 19, 405-19, 1966

#### BIBLIOGRAPHY

VIII-C-(6). ARTICLES RELATING TO BRONCHOPULMONARY AND CARDIOPULMONARY EFFECTS OF CIGARETTE SMOKE

	Reprint
CLOUGH PW: The irritative effect of smoking on the respiratory mucous membranes. Ann Intern Med 45: 1235-41, 1956.	1030
DEVINE K D: Pathologic effects of smoking on the larynx and oral cavity.  Proc Mayo Clin 35: 349-51, 1960.	1031
Diagnosis News: Pulmonary emphysema is a "vascular disease".  2: 16-17, 19	1032
GUILLERM R, BADRE R, SAINDELLE A, HEE J and FLAVIAN N: Acquisitions recentes sur la toxicologie de la fumee de tabac. Gazette Hopitaux 140: 861-62, 1968  Medical Tribune: Nicotine found to intensify calcific disease in rabbits. 29, 1968	1033 1034
NTRDA Bulletin: Blood vessel damage caused by smoking important in emphysema. 12,	1035
OLSEN A M, FONTANA R S, PEABODY H D Jr, and ORDSTRAND H S V Relationship of smoking to chronic respiratory disease. Dis Chest 54: 189-90, 1968	1036
RAMIREZ E A: Protective factors from atherosclerosis in Puerto Rican veterans? Bol Asoc Med PR 161-62, 1969	1037
RANKIN J: The effect of smoking on pulmonary function 24-27	1038
SACHS DP and FINLEY TN: Smoking and pulmonary macrophages. N Engl J Med 286: 788-89, 1972	1039
SHUBIN H: Pulmonary emphysema and the heart. Pa Med 69: 52, 1966	1040
TAYLOR R M: Le probleme de la cigarette. Med Hyg 895: 1450-51, 1969	1041
Tobacco Reporter: Some theories about smoking. 94: 33, 1967	1042
Tobacco Reporter: Dryden gives institute's views on health issues.  54-55, 1967	1043

7-8, 10-11, 1970

page 1	48
Bibliography VIII-C-(7)	Reprint
JONES H: Smoke, chokeTeenagers fight back with smoke-out campaign.  NTRDA Bull 10-12, 1970.	1082
Journal of the American Medical Association: Psychiatrists shrink from giving up cigarettes. 219: 823, 826-28, 1972.	1083
Journal of the Mississippi State Medical Association: Mississippi M. D. 's heed Dr. Ochsner. 56	1084
LONG PH: Doctor, can you stop smoking? Med Times 92: 407-8, 1964	1085
LUBAN-PLOZZA B: De quelques possibilites de deshabituer les fumeurs.  Med Hyg 803: 1337-41, 1967	1086
LUDWIG A M and PYLE R L: Danger potential of commonly abused drugs.  Wis Med J 68: 216-18, 1969	1087
LYNCH HT: Antismoking clinic. Med Tribune	1088
McFARLAND J.W, GIMBEL H.W, DONALD W.A.J and FOLKENBERG E.J: The five-day program to help individuals stop smoking. Conn Med 28: 885-90, 1964	1089
Medical Arts and Sciences: Cigarette labeling law. 132-33, 1965	1090
Medical Tribune: Antismoking leaders hail FCC move; battle looms. 10: 1, 24 1969.	1091
Medical Tribune: Albany College faculty supports ban on radio, TV cigarette advertising. 4, 1969.	1092
Medical Tribune: Hearings opened on FTC rulings on cigarette ads. 8. 1969.	1093
Medical Tribune: California doctors tie illnesses to cigarettes.  3,	1094
MESERVE W G: Congressional action on smoking and health. 257-62	1095
MILLER R K: Summary of proceedings of house of delegates of the Oregon Medical Association. Northwest Med 385, 388-89, 391-93, 1964.	1096
MOERSCH Hill: Counselifor the patient who smokes. Dis Chest 54: 211-20, 1968	1097
MOORE GE and VINCENT RG: Where do we go from here in research.  1-10,	1098
MORGAN KR: Landlord, fill the flowing bowl (and while you're at it, bring me a pack of camels). Consultant 18-19, 1968	1099
MUELLER M: Anti-smoking forces gain ground. Science 165: 569, 1969	1100
MYDLIL F and TOMANEK A: Koureni cigaret a plicni choroby. Rozhledy <u>Tuberk Nemocech Plincnich</u> 24: 649-50, 1964	1101

#### BIBLIOGRAPHY

VIII-C-(8). ARGUMENTS AGAINST SURGEON GENERAL'S STATEMENT	Reprin
American Druggist: In Defense of cigarettes. 83-84, 89, 1970	1139
BARACH A L: Storage of pleasures: the physician's role. Geriatrics  22: 73-74, 1967  BARACH A L: Tobacco smoking and the humanities. JAMA 212: 1525, 1970	1140 1141
BELL J A'E: Mortality rates of smokers.	1142
BROWNLEE KA: A review of "smoking and health" 722-39	1143
CHADWICK DR: Chronic respiratory diseases something can be done.  NTA Bull 9-11, 1967.	1144
DELARUE N C: Gigarette smoking: A clinical and public health challenge.  Can Med Ass J 87: 961-69, 1962.	1145
HAHN MJJr, WESTINGHOUSE WD, SCHREIBER SJ, BRILLMAN CR,  LOMBARD R T and EISENBERG M: Should a strong attempt be made to  get old people to change habits like smoking and overeating. Geriatrics  25: 68, 71, 1970	1146
HUKILL PB: Opening wedge. N Engl J Med 269: 988, 1963	1147
KOTIN P: University of Kentucky tobacco and health workshop proceedings, 1967. 28-38	1148
KOTIN P: The cigarette lung cancer imbroglio and public opinion. JAMA 211: 506, 1970	1149
LEVIN M L: Recent epidemiological research in the relationship between cigarette consumption and health at Roswell Park Memorial Institute.  1-9	1150
MALLAN L: It is safe to smoke. Hawthorn Books Inc: New York City 9-23	1151
MATOUSEK W: Smoking: a basis for advice to the patient. Am Practitioner 8: 1226-28, 1957	1152
New England Medical Journal: Bulletin on tobacco. 268: 1087, 1963	1153
OVERHOLT R H: The physician's obligation in the smoking issue. NY State J Med 64: 1297-1300, 1964.	1154
ROLLINS I M: Two viewpoints a factual presentation of the smoking controversy.  In: Charlotte Postgraduate Seminar, 1969. 1-15	1155

#### IX. RECOMMENDATIONS

The reviewer is submitting the following recommendations to .
the Council for Tobacco Research:

- 1. Publication. That cigarette smoking is one of several risk factors and not the most important cause of chronic pulmonary disease should be publicized to the medical profession and the general public.

  Portions of this review can be expanded and modified for publication and distribution.
- 2. Epidemiology. The expert advice of a statistician is needed to review mortality statistics, specifically to analyze the death rate for pulmonary emphysema separately from that for chronic bronchitis.

  The morbidity statistics should also be examined for regional differences in the United States and comparison between countries.
- 3. Clinical investigation. A five-year follow-up of smokers should be conducted to determine whether those with early signs of ventilatory function develop pulmonary emphysema or chronic bronchitis.
- 4. Pathological investigation. A registry of persons who have been smokers for 50 years would be helpful in determining the influence of cigarette smoking.
- 5. Animal investigation. Attempts to induce pulmonary emphysema or chronic bronchitis in animals should continue. The emphasis should be given to determining the relative importance of air pollution, infection and cigarette smoking in the same laboratory model.

# Additional Bibliographic List No. 1

2 02MONANCI EMPHISEMEN: ETIOLOGY AND PATHOGENESIS	Rej	prir
ADELMAN J U: A review and reappraisal of emphysema. Dis Chest 51: 156-61, 1967.	A	1
BISHOP W R: Pulmonary emphysema. Proc Med Sec Amer Life Convention 51st annual meeting-June 24-26, p. 118-22, 1963.	·-A	2
EBERT R V. and PIERCE J A: Pathogenesis of pulmonary emphysema. Arch Intern Med 111:34-43, 1963.	A	3
GOUGH J. The pathogenesis of emphysema. In: The Lung, edited by Averill A. Liebow and David E. Smith, The Williams and Wilkins Co., Baltimore, 109-33, 1968.	A	4
ISHIKAWA S, FATTAL G A, ZYLAK C, CHERNIACK R M and WYATT J P: A  postmortem study of viscoelastic properities of lung in emphysema (correlation with quantitative morphometry). Pub Hlth Serv No 1787: 325-37, 1968.  KLEIN R C: The etiology of emphysema. Med Counterpoint 2: 13-17, 1970.	A	
MITCHELL R S, TOLL G and FILLEY G F: The early lesions in pulmonary emphysema.  Amer J Med Sci 243: 409-18, 1962.	A	7
RODMAN T and STERLING F H: Etiology and pathogenesis. In: Pulmonary Emphysema and Related Lung Diseases, The C. V. Mosby Co., St. Louis, 3-23, 1969.	A	8
THOMPSON W T: What is emphysema? Virginia Med Month 97: 65-7, 1970.	Å	9
TOMASHEFSKI J F and PRATT P C: Pulmonary emphysema: pathology and pathogenesis.  Med Clin N Amer 51: 269-81, 1967.	<b>. A</b>	10
WRIGHT G W and KLEINERMAN J: A consideration of the etiology of emphysema in terms		

#### List No. 2

RYDER R C. THURLI	BECK M and GOUGI	IJ: A study of interobserver	variation in the
assessment of	the amount of pulm	onary emphysema in paper-mo	
· lung sections.	Amer Rev Resp	Dis 99: 354-64, 1969.	
•			

A 28

SADOUL P: Conceptions actuelles sur l'emphysème pulmonaire. Vie Med 49: 909-34, 1968. A 29

THURLBECK WM: The anatomy and histology of chronic bronchitis and all forms of emphysema. In: Emphysema in Industry, Medical Series Bulletin No. 10, Pittsburgh, p. 1-20, 1966.

30

THURLBECK W'M: A pathologist looks at respiratory failure due to obstructive lung disease. Chest 58 (4 Suppl. 2): 408-14, 1970.

31

THURLBECK W.M.: Present concepts of the pathology and pathogenesis of pulmonary emphysema. Path Microbiol 35: 130-33, 1970.

A 32

WORTH, G, MUYSERS K and SMIDT U: Das Lungenemphysem. Fortschr Med 86: 361-64, 1968.

A 33

WRIGHT RR: Chronic pulmonary emphysema. Arch. Path. 85: 231-6, 1968.

A 34

WYATT J P and FISCHER V: Emphysema and ... Industr Med Surg 33: 213-19, 1964.

A 35

		Repr	cint
BANYAI A L: Pulmonary emphysema. Geratrics 17: 548-59, 1962.	in sangalija	- A	36
BANYAI A L: Pneumoclasia. Denominador común del enfisema pulmonar.  Presna Med Argentina 53: 110-13, 1966.		A	37
BATES D V: The challenge of pulmonary emphysema. Med Coll Va Quart 1: 34-41, 1965.		A	38
BATES D V: The diagnosis of emphysema. Pathol Microbiol 35: 134-37, 1	970.	A	39
BAUMANN H.R.: Zur behandlung des Lungenemphysems. Therap Umschau 592-98, 1968.	25:		40
BURROWS B: Emphysema. Presented at Chicago Medical Society Clinical Conference February 23, 1968.		А	41
CARIELLO M: Interesse medico-social dell'indagine spirografica di routin	ie.		
organisti kalendari (k. 1900). 1984 - Albertan Johnson (k. 1900).		A	42
COLP C, PARK S S and WILLIAMS M.H., Jr: Emphysema with little airwa	y obstruct-	A	43
EBERT R V: Summary of the conference. Amer Rev Resp Dis 80: 209-12,	1959.	A	44
FEGIZ G: Problemi clinico-pathogenetici e sociali dell'enfisema polmonar 976-92,	and the second s	<b>A</b>	45
FLETCHER C M: Summary of conference. Amer Rev Resp Dis 95: 895-99	, 1967.	A	46
GARDINER J.F: Chronic obstructive emphysema. J Omaha Clin Soc 23:11	33-36,	A	47
GLUCKMAN J C: L'emphyseme pulmonaire. Gaz Hop 141:341-43, 1969.		* <b>A</b>	48
HYDE L and HYDE B: Chronic pulmonary emphysema. Postgrad Med 20: 6	09-14, 1956.	A	49
JOHNSTON R F and KAUFFMAN L A: Pulmonary emphysema. Current TI 104-6, 1970.	nerap 1970:	A	50
JULICH Hand RUCKER K: Der einfluss von pulmophyllin auf die korperlic leistungsfähigkeit von kranken mit lungenemphysem. Disch Gesund 49, 1969.	he 24: 346-	A	51
LECKERT JT: Modern trends in the diagnosis and therapy of pulmonary e <u>J Louisiana Med Soc</u> 116: 318-28, 1964.	mphysema.	<b>A</b>	52
LAWS J W and HEARD B E: Emphysema and the chest film: a retrospection and pathological study. Bull J Radiol 35: 750-61, 1962.	ve radiological		53
LYONS H A: Obstructive pulmonary emphysema. Med Times 93: 1386-140	1965.	. A	54
MACKIEM D. Summary of the 13th Aspen emphysems conference. Chest	so (Supplement 5	γ .	

40S-43S, 1971.

McCOY W T and ROBBINS L L: Emphysema: The pulmonary enigma. Texas J Med 49: 15-18, 1963.	<b>A</b> :
MOTLEY H L: Pulmonary emphysema. Missouri Med 57: 701-9, 1960.	<b>A</b> :
NICKLAUS T M, STOWELL D W, CHRISTIANSEN W R and RENZETTI A D, Jr.: The accuracy of the roentgenologic diagnosis of chronic pulmonary emphysema.  Amer Rev Resp Dis 93: 889-99, 1966.	<b>A</b> :
REBORA F: Formas clinicas. Gaceta Med Mexico 756-76, 1965.	Á :
REID, L and MILLARD F J C: Correlation between radiological diagnosis and structural lung changes in emphysema. Clin Radiol 15: 307-11, 1964.	1
ROKSETH R, SCHAANNING J and AAS T W: Cardiopulmonary function tests in fibrosis and emphysema. Acta Tuber Scand 46: 57-64, 1965.	A
SADOUL P, LAXENAIRE M C, LAMBERT H and BERNADAC P: Tableau fonctionnel de l'emphyseme radiologiquement destructif. J Francais Med Chirc Thorax 24: 183-204, 1970.	A
SCADDING J G: Emphysema. Proc Roy Soc Med 62: 1023-24, 1969.	<b>A</b>
SIMON G: Radiology and emphysema. Clin Radiol 15: 293-306, 1964.  SIMPSON T: The emphysema problem. Brit J Dis Chest 62: 188-94, 1968.	<b>A</b>
SUTINEN S, CHRISTOFORIDIS A J, KLUGH G A and PRATT P C: Roentgenologic criteria for the recognition of nonsymptomatic pulmonary emphysema.  Amer Rev Resp Dis 91: 69-76, 1965.	
SWINNY B, SWINNY B, Jr. and HICKS R R: Pulmonary emphysema. Southern Med <u>J</u> 58: 600-5, 1965.	A
THURLBECK W M, BATES D V, FRASER R G and HENDERSON J: Morphologic and functional studies in emphysema. In: Form and Function in the Human Lun Edited by G. Cumming and L B Hunt, E and S Livingstone LTD, Edinburgh and London, 225-48, 1968.	g, A
TOLL G D: Pulmonary emphysema. Rocky Mountain Med J. 25-27, 1964.	· A

#### Additional Bibliographic List No. 4

그런 선생님은 얼마 되는 그는 그는 그는 그들은 이 얼마를 받는다.	
PULMONARY EMPHYSEMA: THERAPY	
	Reprint
	•
BADGER T L and GREEN G M: Advances in the treatment of respiratory disorders.	A 71
Practitioner 199: 455-64, 1967.	A (1
BAKER I M: The use and results of diagnostic and therapeutic bronchoscopies	
in a prison population. J Amer Ostcopath Ass 69: 646-52, 1970.	A 72
BANYAI A L: Pathogenesis and treatment of pulmonary emphysema. J Med Ass Georgia	
50: 469-75, 1961.	A 73
DADACTI A.L. Danier (C. disconnection of the control of the contro	
BARACH A L: Respiración diafragmática y ejercicio con suministro concomitante de oxígeno en el tratamiento del enfisema pulmonar. Prensa Méd Argent	
53: 114-9, 1966.	A 74
BASS H, WHITCOMB J F and FORMAN R: Exercise training: therapy for patients with chronic obstructive pulmonary disease. Chest 57: 116-21, 1970.	A 75
BEDELL G N and SEEBOHM P M: Treatment of pulmonary emphysema. JAMA	A 76
<b>169:</b> 1699-1702, 1959.	A 10
BLAKE J M: Management of pulmonary insufficiency. NY State J. Med 64: 1308-	
12, 1964.	· A 7.7
BOREN HG, BUSET JF, CORPERF, GRZYBOWSKIS, NEWMAN MM, ORGANICK,	
A B and LESTER W: Treatment of drug-resistant tuberculosis. Amer	
Rev Resp Dis 94: 125-7, 1966.	A 78
CARABASI R J: Chronic ventilatory insufficiency and its management. Clin Med	
70: 725-37, 1963.	A 79
COLE M B and HAMMEL J.V: Management of the emphysematous patient. Med Times	•
<b>89</b> : 1026-30, 1961.	· A 80
CODATAL I MORROUGOS - ACUROURAS A SEAS AND ALCOHOLOGICA	
CORAZZA L J, MORROW C.S and CHESNEY T A: Triamcinolone in the therapy of chronic bronchitis and pulmonary emphysema. Amer J Med Sci 242:	
436-42, 1961.	. A. 81
CORPE R F, GRZYBOWSKI S, MACDONALD F M, NEWMAN M M, NIDEN A H,	
ORGANICK A B and LESTER W.: Chronic obstructive lung disease.	-
Amer Rev Resp Dis 92: 513-8, 1965.	A 82
CURTIS J K, LISKA A P, RASMUSSEN H K and CREE E M: IPPB therapy in chronic	•
obstructive pulmonary disease. JAMA 206: 1037-40, 1968.	A 83
DEANE R St. Aspects of respiratory therapy. Med Proceedings 14: 51-6, 1968.	A 84
	•.
DELARUE N C, WOOLF C B and STRASBERG S M: The surgical treatment of pulmonary	A 85

-		page 211
	List No. 4	Reprint
6	EMIRGIL C, SOBOL B J, NORMAN J, MOSKOWITZ E, GOYAL P and WADHWANI  B: A study of the long-term effect of therapy in chronic obstructive pulmonary disease. Amer J Med 47: 367-77, 1969.	A 86
a	FAIN W R: Surgery in pulmonary emphysema. Southern Med J 64: 65-75, 1971.	A 87
.**	FRANKLIN W, MICHELSON A L, LOWELL F C and SCHILLER L W: Bronchodilators and corticosteroils in the treatment of obstructive pulmonary emphysema.  New Engl J Med 258: 774-8, 1958.	A 88
	HEAD J M, HEAD L R, HUDSON T R and HEAD J R: The surgical treatment of emphysematous blebs and localized vesicular and bullous emphysema.  J Thorac Cardiovasc Surg 40: 443-60, 1960.	A 89
	HEPPER N G G and HYATT R E: Office management of chronic bronchitis and emphysema. Med Clin N Amer 48: 857-65, 1964.	A 90
	HEPPER N.G., BRANSCOMB B.V., CHERNIACK R., PIERCE A.K. and MITCHELL R.S:  Home and clinic care of patients with chronic obstructive lung disease. Aspen  Emphysema Conf. 8: 28-34, 1965.	A 91
	JACKSON D and BERNARD L A: Treatment for bullous emphysema. Med Record Annals 61: 113-6, 1968.	A 92
4	JOANNIDES M. Jr.: Chronic obstructive emphysema. The thoracic surgion's position.  JAMA 192: 365-7, 1965.	A 93
O	JOHNSTON R F and HARRIS H W: Management of chronic bronchitis and pulmonary emphysema. Med Clin N Amer 49: 1309-24, 1965.	A 94
	JONES N L, JONES G and EDWARDS R H T: Exercise tolerance in chronic airway obstruction. Am Rev Resp Dis 103: 477-94, 1971.	A 95
	KILBURN K H: Pathogenesis and treatment of chronic bronchitis. Med Times 95: 161-76	<b>A</b> 96
	KILPATRICK G S: Advances in the treatment of respiratory disorders. <u>Practitioner</u> 207: 460-7, 1971.	A 97
	KIRSCHENFELD J J: Office evaluation and management of the patient with chronic pulmonary disease. Southern Med J 56: 193-205, 1963.	A 98
•	LEBOVITZ J J, LEBOVITZ E and SILVERMAN J D: Pulmonary emphysema. <u>Current Therap</u> 113-6, 1969.	A 99
	LEVINE B E, BIGELOW D B, HAMSTRA R D, BECKWITT H J, MITCHELL R S, NETT L STEPHEN T A and PETTY T L: The role of long-term continuous oxygen administration in patients with chronic airway obstruction with hypoxemia.  Ann Intern Med 66: 639-59, 1967.	M, A 100
	LOPEZ-MAJANO V, KIEFFER R F, MARINE D N, GARCIA D A and WAGNER H N, Jr.:  Pulmonary resection in bullous disease. Amer Rev Resp Dis 99: 554-64, 1969.	A 10

	page 212	2 Repr	int
	MAYOCK R L: Obstructive pulmonary empliysema. GP 16: 109-20, 1957.	A 1	02
C	MILLER W F: Rehabilitation of patients with chronic obstructive lung disease.  Med Clin N Amer 51: 349-61, 1967.	A 1	03
• :	MITTMAN C: The management of chronic obstructive lung disease. Chicago Med 70: 177-81, 1967.	A: 1	04
	OSWALD N: Antibiotics and chronic bronchitis. Practitioner 193: 735-9, 1964.	A Ì	05
	OSWALD N: Modern day treatment of chronic bronchitis. Geriatrics 22: 125-32, 1967.	A 1	06
	PASQUAZZI M: La terapia dell'angina pectoris. Policlinico 75: 313-25, 1968.	A 1	07
	PETTY T L: Can anything be done for emphysema? J Louisiana State M Soc 122: 216-8, 1970.	A 1	08
	PETTY T L and NETT L M: Patient education and emphysema care. Med Times 97: 117-30, 1969.	A 1	09
	REEVES J.E: Pulmonary emphysema in general practice. Med Times 90: 512-7, 1962.	A 1	10
•	RICCITELLI M. L: Modern concepts in the management of emphysema in the aged and infirm. J. Amer Geriatrics Soc. 14: 497-504, 1966.	A 1	11
	SCADDING J G: Treatment of chronic bronchitis. Brit Med J 1: 1281-3, 1966.	A 1	12
_	SCHWABER J R: The years beyond smoking. Gariatrics 23: 106-12, 1968.	A 1	113
0	SNIDER, G L: Ambulatory management of chronic obstructive lung disease. Wisconsin Med J 67: 229-32, 1968.	A I	114
. •	SPEAR H C, DAUGHTRY DC, CHESNEY J G and MARKS A: The surgical management of large pulmonary blobs and bullae. Amer: Rev. Resp Dis 84: 186-96, 1961.	A 1	! 15
	STORY SH, Jr.: The management of pulmonary emphysema. <u>J Med Assæ Georgia</u> 53: 24-5	<b>,A</b> : 1	116
	SZUCS M M, Jr. and LORD G P: Outpatient treatment of chronic obstructive pulmonary disease (COPD). J Maine Med Assoc 58: 209-10, 1967.	<b>A</b> 1	117
	ULMER W T: Chronic obstructive airway disease. Recent aspects of diagnosis and treatment. <u>S Afr Med J 46</u> : 146-51, 1972.	." <b>A</b> 1	118
	WATERMAN DH, DOMM S E, ROGERS WK and BORRELL J L: The effective use of bronchoscopy in chronic bronchitis. Ann Otol Rhinol Laryngol 78: 449-58, 1969.	<b>A</b> :	119
	YANDA R L: Hyperbaric research in chronic pulmonary disease. Amer Rev Resp Dis 92: 564-78, 1965.	<b>A</b> :	120
	YANDAR L and MOTLEY H L: Changes in lung volume and spirometric measurements in emphysema patients who have been exposed to hyperbaric pressure levels. <u>Dis Chest</u> 47: 360-66, 1965.	A	121

Source offes //www.

LEBRUN-BRAIS L, BLANCHETTE G and PROVOST G: Emphysome pulmonaire par

1920-34, 1971.

déficience en alpha - antitrypsine: une étude familiale. Union Med Canad 100:

A 137

and the second s	A STATE OF THE STA	reprint ———
	page 214  LIEBERMAN J: Heterozygous and homozygous alpha; antitrypsin deficiency in patients with pulmonary emphysema. New Engl J Med 281: 279-84, 1969.	A 120
	LIEBERMAN J. MITTMAN C and SCHNEIDER AS: Screening for homozygous and	A 138
(	heterozygous al-antitrypsin deficiency. JAMA 210: 2055-60, 1969.	A 139
	MILLER F and KUSCHNER M: Alphan-antitrypsin deficiency, emphysema, necrotizing angiitis and glomerulonephritis. Amer J Med 46: 615-23, 1969.	A 140
	MITTMAN C: Summary of symposium on pulmonary emphysema and proteolysis.  Amer Rev Resp Dis 105: 430-48, 1972.	A 141
The state of the s	NOMIYAMA K, NOMIYAMA H and MATSUI H: Prevalence of a p-antitrypsin deficiency in Japan. Bull WHO 45: 253-55, 1971.	- A 142
	RESNICK H, LAPP, NL and MORGAN WKC: Serum trypsin inhibitor concentrations in coal miners with respiratory symptoms. JAMA 215: 1101-5, 1971.	A 143
	SANO T, YOKOYAMA M and RHOADS GG: Clinical significance of a <sub>1</sub> -antitrypsin level.  A preliminary evaluation among Japanese men Hawaii Med J 31: 27-32, 1972.	A 144
	SCHLEUSENER A, TALAMO RC, PARE JAP and THURLBECK WM: Familial emphysema.  Amer Rev Resp Dis 98: 692-6, 1968.	A 145
	SMITH JP, FALK GA and SISKIND GW: Serum immunoglobulins and alpha; -antitrypsin:	
	Variation with clinical type of chronic obstructive pulmonary disease. Chest 59 (Suppl 5) 17S-19S, 1971.	A 146
	STEVEN P M, Hnilica V, Johnson P C and BELL RC: Pathophysiology of hereditary emphysema. Chest 59: (Suppl) 19S, 1971.	A 147
0	SZCZEKLIK A, TERESIAK T and KOLACZKOWSKA B: Zachowanie sie antytrypsyny alfa w surowicy krwi u ludzi zdrowych i chorych na rozedme pluc. Pol Tyg Lekar 26: 1569-71, 1971	A 148
	TALAMO RC: The a1-antitrypsin in man. J Allerg Clin Immunol 48: 240-50, 1971.	A 149
	TALAMOR C, AUSTENK F and ALLEN J D: Familial emphysema and alphalantitrypsin deficiency. Public Hith Service No. 1787 491-95, 1972.	A 150
•	TALAMO RC. BLENNERHASSET JB and AUSTEN K F: Familial emphysema and alpha; antitrypsin deficiency. New Engl J Med 275: 1301-5, 1966.	A 151
	TARKOFF M P, KUEPPERS F and Miller W F: Pulmonary emphysema and alphalantitrypsin deficiency. Amer J Med 45: 220-28, 1968.	A 152
•	TOWNLEY R.G., RYNING F. LYNCH H and BRODY A.W.: Obstructive lung disease in hereditary ag-antitrypsin deficiency. JAMA 214: 325-31, 1970.	A 153
	TURINO GM: Proteolytic mechanisms and pulmonary emphysema. Bull Physiol Pathol Resp 7: 719-27, 1971.	A 154
	VARPELA E and SARIS NE: Hereditary alpha 1-antitrypsin deficiency. Ann Clin Res 3: 46-53, 1971.	A 155
<b>E</b>	VIDAL J, CAZAL P, ROBINET-LEVY M and MICHEL F B: Bronchopneumopathies chroniques familiales et déficit homozygote en a-l antitrypsine. Poumon Coeui 27: 79-94, 1971.	A 156
	WELCH M.H. REINECKE M. E. HAMMARSTEN JF and GUENTER C.A. Anticrypsin deficiency in pulmonary disease: the significance of intermediate levels.  Ann Intern Med. 71: 533-42, 1969.	A 157

#### Additional Bibliographic List No. 6

## CHRONIC BRONCHITIS

•.	rioside di interitoriale di la companya della companya della companya della companya della Rio della companya d La companya di la companya della companya della companya della companya della companya della companya della co	ep:	rint
	ASHLEY DJB: Environmental factors in the aetiology of lung cancer and bronchitis. Brit J Prev Soc Med 23: 258-62, 1969.	Ą	158
	EDWARDS G: Acute bronchitis actiology, diagnosis, and management. Brit Med J 1: 963-66, April 16, 1966.	A	159
	FLETCHER C M: Chronic bronchitis. Its prevalence, nature, and pathogenesis. Amer Rev Resp Dis 80: 483-94, 1959.	A	160
-	FLETCHER C M: Définition, classification et étiologie de la bronchite chronique. <u>Le Poumon I</u> <u>Coeur</u> 21: 1239-48, 1965.		161
	FLETCHER C M: Some recent advances in the prevention and treatment of chronic bronchitis and related disorders. With special reference to the effects of cigarette smoking.  Proc Roy Soc Med 58: 918-28, 1965.	A.	162
	FLETCHER C M: Recent clinical and epidemiological studies of chronic bronchitis. Scand J  Resp Dis 48: 285-93, 1967.	A.	163
	FLETCHER C M: Bronchial infection and reactivity in chronic bronchitis. J Roy Coll Phys Ion 2: 183-90; 1968.		164
١.	GEISLER L: Zum problem der chronisch-obstruktiven bronchitis. <u>Deutsches Medizinisches</u> Journal 17: 184-7, 1966.	A.	165
 >/	HAIN E: Soziale Bedeutung und Gestaltungsfaktoren des, Bronchitischen Syndroms. Internist 10: 144-54, 1969.	A	166
•	HARTUNG W and MEYER-CARLSTADT D: Reid's index in the diagnosis of chronic bronchitis.  Beitraege zur Pathologischen Anatomie und Allgemeinen Pathologie 137: 85-98, 1968.	A.	167
	KAATZSCH H: Der Einflub des Rauchens auf die Ventilationsstörung des Bronchitikers.  Med Klin 64: 2196-9, 1969.	A	168
	LICKINT F: Neuere Erkenntnisse auf dem Gebiete der Beziehungen zwischen Raucherbronchitis und Bronchialkrebs. Verli Deut Ges Inn Med 62: 121-25, 1956.		169
	LOOSLICG: Synergism between respiratory viruses and bacteria. Yale J Biol Med 40(5-6): 522-40, 1968.	A	170
• :	LOWE CR: Chronic bronchitis and occupation. Proc Roy Soc Med. 61: 98-102, 1968.	A	17,1
	MACKLEM PT: Airway obstruction and collateral ventilation. Physiological Rev 51(2): 368-4.  April 1971.		: 172
	MICZOCH F: Zivilisationsschäden in der Entstehung und Behandlung der chronischen Bronchiti Wiener Klinische Wschr 77: 506-8, 1965.	s. A	173
,	MORGAN W K C: Chronic bronchitis. The "English disease." Postgrad Med 41: 190-94, 1967		174
			175
•	PEMBERTON J: Occupational factors in chronic bronchitis. Proc Roy Soc Med 61: 95-98, 196		176
	PHILLIPS A M. PHILLIPS R W. and THOMPSON'I IV. Chronic coughs analysis of ctiologic.		- 10

REID L: Chronic bronchitis and hypersecretion of mucus. Lect Sci Basis Med 8: 235-55,	vehti	
1959.	A l	78
REIMANN H A: Acute respiratory tract infections in the aged. Geriatrics 22: 160-67,		
1967.	A 1	70
ROSE B and PHILLS J A: The immune reaction in pulmonary disease. Arch Environ		÷
Health 14: 97-110, 1967.	A 1	80
ROSS C A C, McMICHAEL S, EADIE M B, LEES A W, MURRAY E A, and PINKERTON I: Infective agents and chronic bronchitis. Thorax 21: 461-64, 1966.	A 1	8-1
SCHMIDT O-P: Chronische Tabakrauch-Inhalation und Erkrankungen der Atemwege unter Berücksichtigung des Rauchens von Jugendlichen. Munch Med Wochenschr 104:		
1832-36, 1962.	A I	82
SCHMIDT O P: Prophylaxe des chronischen bronchitischen Syndroms. Med Welt 22:		
1416-7, 1968.	A 1	83
STUART-HARRIS C: Infection, the environment and chronic bronchitis. J Roy Coll		
Phys Lond 5: 351-61, 1971.	. A 1	84
TRINQUET G and MEYER A: Role de l'oxyde de carbone dans le transfert de l'oxygène		
au cours des bronchopathies obstructives. Le Poumon et le Coeur 27: 695-704,		: ·
1971.	A 1	85

Reprint

#### Additional Bibliographic List No. 7

#### BRONCHIAL CARCINOMA

AMERICAN CANCER SOCIETY, INC and NATIONAL CANCER INSTITUTE U.S. PUBLIC HEALTH SERVICE: Discussion. In: Proceedings of the Fourth National Cancer Conference. J B Lippincott Co., Philadelphia 323-33, 1960. A 186 ANONYMOUS: Malignant cells in sputum. A 187 BEDROSSIAN C W M: Coexistent emphysema and bronchogenic carcinoma Cancer Bull 24: 102-7, 1972. A 188 BERNIER J L: Tumors of the lips. In: Tumors of the Oral Regions. Edited by HBG Robinson, WB Saunders Co., Philadelphia, 19-28, 1957. A 189 BLUMBERG B S: Common viruses and their relation to cancer etiology. Cancer À 190 2. 1969. Bull 21: 83-5, 1969. BLUNDI E: A Batalha Do Cancer Do Pulmao. Brasil Med 83: 198-205, 1969. A 191 BONSI R: Dati recenti sull'attivita cancerigena del tabacco. A 192 BRYSON G and BISCHOFF F: The limitations of safety testing. In: Progress in Experimental Tumor Research, Edited by F Homburger, S. Karger, New York, 100-33, 1969. A 193 COSIO VILLEGAS I: La prevencion del cancer pulmonar. Neumol Cir Torax 27: A 194 A 195 CLOUTIER J.P.: La lutte contre le cancer au Quebec. Laval Med 38: 421-23, 1967 A 196 DELARUE N C: Cigarette smoking and lung cancer. Appl Ther 4: 919-25, 1962. FOITZIK E. STAEMMLER M and POSZICH G: Experimental pulmonary carcinomas in albino mice after intranasal infection with influenza virus A, Asia. Z Krebsforschung 77: 77-82, 1972. A 197 GSELL OR: Epidemiology of bronchial carcinoma in a mountain region. Oncology 25: A 198 410-14, 1971. HAMMOND E C: Epidemiologic features of cancer of the lung. In: Cancer Management, sponsored by American Cancer Society, Inc. J B Lippincott Co, Philadelphia, 453-58. A 199 HAMMOND E C: Evolution of concepts concerning etiology and pathogenesis of cancer. Industr Med Surg 37: 616-20, 1968. A 200 HAMMOND E.C: Cancer prevention and competitive risks. Arch Environ Hith 19: 395-402, 1969. A 201 HEALY T M: Carcinoma of the lung: 352 cases. Irish J Med Sci 140: 564-71, 1971. A 202 HYATT C L: Physicians and smoking. J Arkansas Med Soc A 203 KOURILSKY R., HAPPERT J. C. and FRED. D. Donnees Experimentales sur L'Action

Cancérigene Bronchopulmonaire Du Tabac. Bronches 16: 370-80, 1966.

	List No. /	Kep	rint
	KREYBERG L: Actiology of Cancer. Universitetsforlaget, Oslo, 1-90, 1969.	A	205
	LANGSTON, H T: Lung cancerfuture projection. J Thorac Cardiovasc Surg 63: 412-15, 1972.	A	206
<b>S</b>	LEUCHTENBERGER C and LEUCHTENBERGER R: Effects of chronic inhalation of whole fresh cigarette smoke and of its gas phase on pulmonary tumorigenesis in Snell's mice. In: Morphology of Experimental Respiratory Carcinogenesis, Sponsored by National Cancer Institute and U. S. Atomic Energy Commission, 329-46, 1970.		207
	MEINSMA L: Roken en Longcarcinoom. Ned Tijd Genees 111: 233-34, 1967.		208
ं <b>क</b> ्र	MENDEZ F L, Jr., MAURER E R P and FRECKMAN H A: Effect of chemical agents on bronchogenic cancer. Arch Surg 87: 147-53, 1963.		209
	Stonenogene career, Men darg of 147-33, 1783.	, A.	209
	MORGAN J M: Hepatic copper, manganese, and chromium content in bronchogenic carcinoma. Cancer 29: 710-13, 1972.	A	210
	OVERHOLT RH: The challenge of pulmonary carcinoma. Delaware Med J 33: 63-67, 1961.		211
	OVERHOLT R H: Primary carcinoma of the lung. CA 17: 262-67, 1967.	A	212
	OVERHOLT R H, OLIYNYK P N and CADY B: The current status of primary carcinoma of the lung. In: Progress in Clinical Cancer, Edited by I M Ariel, Grune and Stratton, New York IV: 211-22, 19	. А	213
	RASMUSSEN R A, BASINGER C E, HARRISON R W and MEADE R H: Choice of operation in the treatment of bronchogenic carcinoma. Dis Chest 46: 190-7, 1964.	A	214
)	RIGDON R H and KIRCHOFF H: Cancer of the lung 1930 to 1960. A review. Texas  Rept. Biol. Med. 19: 465-513, 1961.	A	215
	RIGGINS H M: Primary lung cancer: present and future.	·A	216
•	ROBERTSON L S: The National Cancer Association of South Africa. S. African Cancer Bull 14: 119-24, 1970.	Ą	217
	ROSS, C A: Cigarette smoking and cancer of the lung. Geriatrics 23: 179-80, 1968.	A	218
	SPAIN D M: The distribution of tracheobronchial metaplasia (regenerative hyperplasia) (age, sex, cigarette smoking and relation to reid indexpreliminary report).  In: Current Research in Chronic Respiratory Disease, Proceedings 11th Aspen Emphysema Conference, Aspen Colorado, June 12-15, 1968, 183-186, 1963.		219
	STASZEWSKI J: Palenie a Rak Wargi, Jamy Ustnej, Migdalkow I Krtani. Nowotwory 20: 121-32, 1960.	A	220
•:	TOKUHATA, GK: Cancer of the lung: host and environmental interaction. Penna Dept. of Hltll. 1-48, 1972.	À	221
	VAN DER WAL AM, HUIZINGA E, ORIE N G M, SLUITER H J and DE VRIES K: Cancer and chronic non-specific lung disease (CNSLD). Scand J. Resp. Dis 47: 161-172, 1966.	<b>A</b>	222
5	WYNDER E L and HOFFMANN D: The epidermis and the respiratory tract as bioassay system in tobacco carcinogenesis.  Brit J Cancer 24: 574-87, 1970.		223
٠.			

Accitional Bibliographic List No. 8	
Activitati Protographic Eist Not 8	page 219
AIR POLLUTANTS AND LUNG DISEASE	
	Reprint
ADAMS A I: The stresses of urban living. Health 19: 12-15, 1969.	A 224
ANDERSON D O: The effects of air contamination on health. Canad Med Ass J 97: 528-36, 1967.	A 225
ANDERSON DO: The effects of air contamination on health. Part II Canad Med Ass <u>J</u> 97: 585-93, 1967.	A 226
ANDERSON D O and FERRIS B.G., Jr.: Community studies of the health effects of air pollution a critique. Canad J Pub Health 57: 209-220, 1966.	A 227
AYRES S M and BUEHLER M E: The effects of urban air pollution on health. Clin Pharmacol Ther 11: 337-71, 1970.	A 228
BALCHUM O J: Environment in relation to respiratory disease. Arch Environ Hith 4: 561-74, 1962.	A 229
BATES D V: Air pollutants and the human lung. Amer Rev Resp Dis 105: 1-13, 1972.	A 230
BEERENS J: Pollution de l'air luchtverontreiniging. Acat Tuber Pneumol 53: 429-48, 1967.	A 231
BRODINE V: Point of damage. Environment 14: 2-15, 1972.	A 232
CAREY G C R, PHAIR J J, SHEPHARD R J and THOMSON: The effects of air pollution on human health. Amer industr Hyg Ass J 19: 363-70, 1958.	n .A 233
CASARETT L J: Toxicology: The respiratory tract. Ann Rev Pharm 11: 425-46, 197	1. A 234
DE LOURDES FONSECA SANTOS M: Aspectos clínicos das Doencas Broncopulmonare de Causa inalatioria. Jornal do Medico 77: 143-49, 1971.	A 235
DEVILLIERS, A: The effects of air pollution on health. Occup. Hith Rev 20: 25-44, 1968 1969.	A 236
ECKARDT R E: Air pollution, lung cancer, and chronic lung disease. J Occup Med 6: 184-8, 1964.	A 237
FERRIS B G: Eipidemiological studies on air pollution and health. Arch Environ Hith 16: 541-55, 1968.	A 238
FERRIS B G and WHITTENBERGER J L: Environmental hazards. Effects of communair pollution on prevalence of respiratory disease. New Engl J Med 275: 1413-1966.	ity 19 A 239
GOLDSMITH J R: The new airborne disease. Calif Med 113: 13-20, 1970.	A 240
GOLDSMITH J R, GREENBURG L, ALTSHULLER, A P, SPICER W S, CASSELL E J, and LANDSBERG H E: Amer Rev Resp Dis 93: 302-12, 1966.	A 241
GROSS P: The pulmonary reaction to man's environment. 106-11,	<b>▲ 242</b>
HAMMOND E C: Epidemiological evidence on the effects of air pollution. 60th Annual Meeting Air Pollution Control Association, Cleveland, Ohio, June 14, 1967.	A 243
HAMMOND E C: Testimony. Before the Subcommittee on Air and Water Pollution of Senate Committee on Public Works, July 30, 1968.	the A 244

List No. 8	Reprint
HEIMANN H: Effects of air pollution on human health. WHO Monogr Ser. No. 46: 159-220, 1961.	A 245
HEIMANN H: Status of air pollution helath research, 1966. Arch Environ Hith 14: 488-503, 1967.	A 246
HIGGINS I T T: Air pollution and chronic respiratory disease. Ashrae J 8: 37-45, 1966.	A 247
INGRAHAM H S: Respiratory disease and air pollution. 4-17.	A 248
KOTIN P: Opportunities in environmental and occupational pathology. <u>Lab Invest</u> 13: 619-23, 1964.	A 249
LINDALL A W: Effect of air pollution from fossil fuel combustion on human health.  Minnesota Med 321-26, 1970.	<b>A</b> 250
MOTLEY H L and PHELPS H W: Pulmonary function impairment produced by atmospheric pollution. Dis Chest 45: 154-62, 1964.	A 251
OCHSNER A: The hazards of air pollution fact or fiction? . Med Tribune, March 11 and 14, 1967.	A 252
PRINCI, F: Air pollution facts and fable. J Occup. Med 461-67, 1963.	A 253
REICHEL G: Umwelteinflusse and obstruktive atemerkrankungen. Schweiz Med Wschr 102: 102-9, 1972.	A 254
SALATA JR: Air pollution. Rocky Mount Med J 63: 47-51, 1966.	A 255
SPICER WS, Jr.: Relation of air pollution to disease. Arch Environ Hlth 9: 600- 5, 1964.	A 256
SPOTNITZ M: Air pollution respiratory syndromes: A military medical problem.  Milit Med 131: 1499-1503, 1966.	A 257
STRESEMANN E: The medical and biological importance of exhaust gases from motor vehicles. Staub-Reinhalt Luft 29: 11-16, 1969.	A 258
WALLER RE: Air pollution and community health. J Roy Coll Physicians Lond 5: 362-8, 1971.	A 259
WINKELSTEIN W, Jr. and FRENCH F E: The role of ecology in the design of a health care system. Calif Med 113: 7-12, 1970.	A 260
health care system. Calif Med 113: 7-12, 1970.	

#### Additional Bibliographic List No. 9

## PULMONARY EMPHYSEMA-BRONCHITIS COMPLEX: ETIOLOGY & PATHOLOGY

	Reprin
ANDERSON DO: Chronic nontuberculous respiratory diseases. In: Preventive Medicine, edited by Duncan W Clark and Brian MacMahon, Little, Brown and Company, Boston, 491-507,	A 261
	· · ·
ANONYMOUS: The nature of chronic bronchitis and emphysema. NTA Bulletin :7-12, 1967.	A 262
BARCLAY WR: Axioms on chronic pulmonary disease. Hosp Med 5: 70-1, 75-9, 1969.	A 263
BATES DV: Chronic bronchitis and emphysema. New Engl J Med 278: 546-51, 1968.	A 264
BATES D V: Chronic bronchitis and emphysema (concluded). New Engl J Med 278: 600-5, 1968.	A 265
BURROWS B, FLETCHER C M, HEARD B E, JONES N L and WOOTLIFF J S: The emphysematous and bronchial types of chronic airways obstruction. A clinico-pathological study of patients in London and Chicago. Lancet 1: 830-5, 1966.	A 266
BURROWS B and KETTEL L J: Important considerations in emphysema-chronic bronchitis syndrome. Geriatrics 24(11): 72-82, 1969.	A 267
COHEN B M: When to suspect that masquerading malady: obstructive lung disease.  Consultant 7: 32-6, 1967.	A 268
DAVID W: Chronic respiratory diseases — the new look in the public health service.  Am J Public Health 57: 1357-62, 1967.	A 269
EBERT R V: Chronic bronchitis and pulmonary emphysema. Postgrad Med 29: 341-5, 1961.	A 270
FLETCHER C M: Discussion on bronchitis and emphysema. Practitioner 188: 403-5, 1962.	A 271
FLETCHER C.M: Bronchitis and emphysema. Proc Roy Soc Med 55: 451-3, 1962.	A 272
FLETCHER C.M: Bronchitis-emphysema syndrome. Disease-a-Month December: 1-48, 1963.	A 273
FRUHMANN GExogene und endogene Faktoren in der Atiologie chronisch-unspezisischer Atemwegerkrankungen. Med Klinik 63: 1288-95, 1968.	A 274
HARTUNG M: Die Klinik der chronischen unspezifischen Atemwegsyndrome. Zugleich ein Beitrag zur klinischen Diagnostik und Klassifizierung. Med Klinik 63: 1296-1302, 1968.	A 275
HARTUNG W: Pathologische Anatomie und Pathogenese der Bronchitis und des Emphysen Internist 10(4): 121-32, 1969.	ns. A 276
HOLLAND W W: Chronische bronchitis und lungenemphysen. In: Nikotin, edited by Helm Schievelbein, Georg Thieme Verlag, Stuttgart, 230-41, 1968.	aut A 277

HUNT JR W B: Criteria for diagnosis of asthma, chronic bronchitis and emphysema. With

	page 224
List No. 9 KINSELLA T J: The indeterminate lung lesion. J Lancet 82: 27-34, 1962.	Reprint A 279
LARSEN K A: Patologisk anatomi og patogenetiske faktoren. Lakartidningen 67: 375-83, 1970:	A 280
MARX HH: Die emphysembronchitis. Disch Med Wochenschr 96: 833-8, 1971.	A 281
MENEFEE JR E E: Bronchitis and emphysema. Virginia Med Monthly 84: 53-6, 19	57. A 282
MITCHELL RS: The fight against chronic non-specific respiratory diseases. In: Bronchitis, edited by NG M Orie and H J Sluiter, Royal Vangorcum, Assen, Netherlands and Charles C Thomas, 343-52, 1964.	A 283
MITCHELL R S, SILVERS G W, DART G A, PETTY T L, VINCENT T N, RYAN S F FILLEY G F: Clinical and morphologic correlations in chronic airway obstru- Am Rev Resp Dis 97: 54-61, 1968.	
MORGAN W K: Irreversible obstructive airway disease: definition, pathology and etiology. West Va Mcd J 64: 288-92, 1968.	<b>A</b> 285
MULLER H: Chronic bronchitis and emphysema. S Afr Med J 33: 805-7, 1959.	A 286
OSWALD N C: The clinical importance of recent advances in bronchitis and emphysic Postgrad Med 35: 233-6, 1964.  PRATT P C and THONG-YAI K: The relative importance of bronchiolitis and extension centrilobular emphysema in pulmonary ventilatory interference. Public Healt Service Publication No. 1787: 339-55, 1968.	A 287
PUGSLEY H E: The underwriting of chronic pulmonary diseases. Proc ALC Med Se : 106-16,	
RAPPAPORT (and MAYER E: Changing forms and causes of pulmonary insufficience present era of medicine. <u>J Am Med Ass</u> 165: 431-6, 1957.	y in
TAUB S J: Emphysema, chronic bronchitis and asthma. Eye Ear Nose Throat Mor 45: 110-2, 1966.	A 291
THURLBECK W M: Chronic obstructive lung disease. : 367-98,	<b>A 292</b>
THURLBECK W M and ANGUS G E: The relationship between emphysema and chron bronchitis as assessed morphologically. Am Rev Resp Dis 87: 815-9, 1963.	ic A 293
THURLBECK W M, HENDERSON J A, FRASER R G and BATES D V: Chronic obstruments of disease. A comparison between clinical, roentgenologic, functional and morphologic criteria in chronic bronchitis, emphysema, asthma and bronchie Medicine 49: 81-145, 1970.	• .
TURNER W A: Chronic obstructive lung disease. Nova Scotia Med Bull 48: 3-11, 19	69. A 295
VALENTIN Hand WOITOWITZ H-J: Chronische Bronchitis und Lungenemphysem arbeitsmedizinisches Problem. Internist 8: 165-72, 1967.	
VOISIN C: Les facteurs etiologiques dans la bronchite chronique et l'emphysème pul. <u>La Revue de Medecine</u> 11: 2175-9, 1970.	monaire. A 297

WESSEL-AAS T: Obstruktive Lungesykdommer. Tidsskr Nor Laegeforen 87: 566-8, 1967. A 298

#### Additional Bibliographic List No. 10

#### PULMONARY EMPHYSEMA-BRONCHITIS COMPLEX: CLINICAL STUDIES

ĊĮ.	ASTIN T W: Reversibility of airways obstruction in emphysema, Lancet 1: 1128, 1972.	Reprint A 299
41.11	The second secon	•
	BARACH A L, CHUSID E L and WOOD L: Ventilatory effect of decreasing functional residual capacity in pulmonary emphysema. Ann Allergy 25: 211-23, 1967.	A 300
	BROCARD H: Poumon et travail; exposé de synthèse. CAMIP 35-36: 88-94, 1969.	A 301
۽ تن	BROWN E A: The pulmonary cripple. Mississippi Valley Med J 80: 80-5, 1958.	A 302
:. <u>,</u> :	BURROWS B: Chronic obstructive lung disease. (Bronchitis-emphysema syndrome)  Diagnosis and physiologic effects. Postgrad Med 39: 105-12, 1966.	A 303
	BURROWS B: The course of patients with chronic obstructive lung disease. In:  Current Research in Chronic Respiratory Diseases. Proceedings of the  lith Aspen Emphysema Conference, Aspen, Colo. U.S. Department of  Health, Education and Welfare, Public Health Service Publication No.	
	1879: 253-8, 1969.	A 304
) .	CARABASI R J: Asthma? Bronchitis? Emphysema? A diagnostic dilemma.	
	Postgrad Med 39: 553-5, 1966.	A 305
•	DALY W J: Single-breath gas distribution in chronic obstructive lung disease.  Arch Intern Med 127: 763-6, 1971.	A 306
٠.	DINES DE: Monitoring certain aspects of respiratory failure. Dis Chest 54: 421-6,	
	1968.	A 307
	EARLE R H and BURROWS B: Prognosis in chronic obstructive lung disease. Public Health Service Publication No. 1787: 453-62, 1972.	A 308
	FARCON E M and MORALES PA: The association of chronic obstructive pulmonary disease (COPD) and lower urinary tract symptoms. J Urology 108/4: 619-22, 1972.	A 309
	FILLEY G.F: Emphysema and chronic bronchitis: clinical manifestations and their physiologic significance. Med Clin N Amer 51: 283-92, 1967.	A 310
•	FLETCHER C M, HUGH-JONES P, McNICOL M W and PRIDE N B: The diagnosis of pulmonary emphysema in the presence of chronic bronchitis. Quart J Med 32:	
	<b>33-47,</b> 1963.	A 311
	GONZALEZ E, WEILL H, ZISKIND M M and GEORGE R B: The value of the single breath diffusing capacity in separating chronic bronchitis from pulmonary emphysem	13
•	Dis Chest 53: 229-36, 1968.	A 312
	HARRIS H.W.: Chronic bronchitis, emphysema, and asthma. Amer J Pub Health 53 suppl. PHI: 7-15, 1963.	A 313
	HEPPER NG, BLACK LF, DINES DE and MILLER RD: The concept of an emphysema clinic. Mayo Clin Proc 43: 306-10, 1968.	A 314

HERZOG H: Chronisch-obstruktive Lungenerkrankungen. Pathophysiologie, Klinik und

	P	a ge 22
	List No. 10	Reprint
	HOGG J C, MACKLEM P T and THURLBECK W M: The resistance of small airways in normal and diseased human lungs. Public Health Service Publication No. 1787: 433-41, 1972.	A 316
*	HUTTNER I and HUTTNER II: Die Motivation des Gesundheitsverhaltens. Teil I.  Zschrärztl Fortbild 64: 471-7,	A 317
	KRUMHOLZ RA: A comprehensive respiratory disease program. Oliio St Med J 65: 147-9, 1969.	A 318
	LORD G P, GAZIOGLU K and KALTREIDER N: The maximum expiratory flow-volume in the evaluation of patients with lung disease. Am J Med 46:72-9, 1969.	A 319
	MARX H H: Gesichtspunkte zur Begutachtung der primär oder sckundar unspezifischen Lungenkrankheiten. Beitr Klin Tuberk 136: 172-8, 1966.	A 3.20
	MOLHO M, RAZ I and KREISLER B: A basis for differentiating between primary emphysema and pulmonary obstruction disease due to chronic bronchitis. :29-35,	A 321
	MORTON J W and TURNBULL K W: The reversibility of chronic bronchitis, asthma and emphysema. Dis Chest 53: 126-32, 1968.	A 322
77.	PENMAN R W and ASTIN T W: Airway obstruction due to hypoxemia in patients with chronic lung disease. Public Health Service Publication no. 1787: 145-59, 1968.	A 323
<b>O</b>	PENMAN R W. O'NEILL R P and BEGLEY L: The progress of chronic airway obstruction in relation to measurements of airway resistance and lung elastic recoil. Am Rev Resp Dis 101: 536-44, 1970.  PIERCE J A: Office management of chronic obstructive pulmonary disease. Clin Notes	A 324
	Resp Dis 7: 3-11, 1968.	A 325
<u>.</u>	PIRCHER F J: Lung scanning in the assessment of obstructive lung disease. Public Health Service Publication No. 1787: 429-31, 1972.	A 326
elv Line	PISTAWKA H J: Chronic airway obstruction. Current Therap : 99-104, 1969.	A 327
•	SIERING H: Bronchitis, Peribronchitis und Emphysem. Zschrärztl Fortbild 60: 602-8,	A 328
	SUFRIN M: To screen or not to screen for COPD. <u>Bull Nat Tuberc Resp Dis Ass</u> 56: 11-13, 1970.	A 329
	SUKUMALCHANTRA YOSVI and WILLIAMS JR MH: Serial studies of pulmonary function in patients with chronic obstructive pulmonary disease. Am J Med 39: 941-5, 1965.	A 330
•	TECULESCU D and RACOVEANU C: A clinical roentgenologic and biologic classification of patients with chronic obstructive lung disease: Rev Roum Med Int. 9: 39-50, 1972.	A 331
	THURLBECK W: The pathological anatomy of chronic obstructive lung disease. J Clin Pathol 24: 482, 1971.	A 332
T	TYSINGER JR DS: Pulmonary function testing for the general hospital and physician. <u>J Med Ass Alab</u> 39: 834-40, 1970.	A 333
1	VARGHA G: Eine neue - auf die Bronchiale Schleimdrüsenhyperplasie Gegründete - Entstehungstheorie für das Obstruktive Emphysem. Acta Med Acad Scient Hung 26. 73.8 1069	A 334

## Additional Bibliographic List No. 11

# MORBIDITY STUDIES

	Reprint
ANON: The ethnic distribution of disease in the United States. J Chron Dis 20: 115-18, 1967.	A 335
ANON: Zur bedeutung chronisch-inhalativer noxen am arbeitsplatx fur chronische bronchitis und lungenemphysem (The significance of chronic inhalable noxae at the place of work for chronic bronchitis and pulmonary emphysema). Munch Med Wochenschr. 113: 1302-7, 1971.	A 336
ASHFORD J R, BROWN S, MORGAN D. C and RAE S: The pulmonary ventilatory function of coal miners in the United Kingdom. Am Rev Resp Dis 97: 810-21, 1968.	A 337
BAKER TR. OSCHERWITZ M, CORLIN R, JARBOE T, TEISCH J and NICHAMAN MZ: Screening and treatment program for mild chronic obstructive pulmonary disease. JAMA 214: 1448-1455, 1970.  BERNDT V H: Bronchialkarzinom und ulcus pepticum. Med Klin 57: 1397-1401, 1962.	A 338
BLACK LF and HYATT RE: Maximal static respiratory pressures in generalized neuromuscular disease. Am Rev Resp Dis 103: 641-50, 1971.	A 340
BROWN K E and CAMPBELL A H: Tobacco, alcohal and tuberculosis Br J Dis Chest 55: 150-8, 1961.	<u>t</u> A 341
BURNS MW and MAY JR: Haemophilus influenzae precipitins in the serum of patients with chronic bronchial disorders. Lancet 1: 354-58, 1967.	A 342
CALIFORNIA MEDICAL ASSOCIATION: The smoking study. Calif Med 109: 339-44, 1968.	A 343
CHERNIACK N S, DOWLING H F, CARTON R W, and McBRYDE V E: The role of acute lower respiratory infection in causing pulmonary insufficiency in bronchiectasis. Ann Intern Med 66: 489-97, 1967.	A 344
COHEN A C and JENNEY F S: The frequency of peptic ulcer in patients with chronic pulmonary emphysema. Am Rev Resp Dis 85: 130-32, 1962.	A 345
CONALLY N T: Early detection of disease in young men. Med Ann DC 38: 672-76, 1969:	A 346
CUNNINGHAM D.A., MONTOYE H.J., HIGGINS M.W. and KELLER J.B.: Smoking habits, chronic bronchitits and shortness of breath and physical fitness. Med Sci Sports 4: 138-45, 1972.	A 347
D'ANELLI A, PROSCIA N, SORISTO F and CARATTI C A: L'enfisema polmonare cronico ostruttivo nel malato ulceroso (Chronic obstructive emphysema in the ulcer patient). Arch Sc Med 17: 191-94, 1965.	A 348
DeMARRAIS G A: The meteorology associated with New Orleans asthma.  Arch Environ Health 11: 787-93, 1965.	A 349

Reprint

A 350

A 351

bronchitis to prednisone. Ann Intern Med 71: 711-78, 1969.

Emphysema. Dis Ches. 41: 514-17, 1962.

KUPKA E and GOLDSMITH J.R: Crossing of the curves II: Tuberculosis and

1311-13, 1969;

List No. 11

DOYLE J T: Cigarette smoking. Minn Med

Arch Environ Health 14: 189-200, 1967.

ENTERLINE P E: The effects of occupation on chronic respiratory disease.

FERRIS B G Jr, ANDERSON D O and BURGESS W A: Prevalence of respiratory

A 368

<b>₩</b>	page 22
List No. 11	Reprint.
LEMLE A, VIEIRA LOBD, MILWARD GAF and MIRANDA JL: Lung function studies in pulmonary South American blastomycosis. Am J Med 48: 434-42, 1970.	A 369
MATHOV E: Complicaciones del asma bronquial. El Dia Medico (4 Dec 1969), pp 2806-10.	A 370
McKEE W D: Environmental problems and the allergist. Ann Allergy 29: 510-24, 1971.	A 371
MEDICAL RESEARCH COUNCIL: Value of chemoprophylaxis and chemotherapy in early chronic bronchitis. Bull Med J 1: 1317-22, 1966.	A 372
MERRILL M H: Public health responsibilities and program possibilities in chronic respiratory diseases. Am J Public Health 53: 25-33, 1963.	A 373
MILLER G J and SERJEANT G R: An assessment of lung volumes and gas transfer in sickle-cell anemia. Thorax 26: 309-15, 1971.	A 374
MILLS M and BAISCH BF: Spontaneous pneumothorax. Ann Thorac Surg 1: 286-97, 1965.	A 375
MOFFAT MAJ and SUTHERLAND JAW: Persistence of viral antibodies in patients with chronic bronchitis. Br Med J 1: 601-603, 1967.	A 376
MONSON R R: Duodenal ulcer as a second disease. Gastroenterology 59: 712-16, 1970.	A 377
OTTO H and HINUBER G V: Zur haufigkeit des emphysema, der tuberkulose und des bronchuskarzinoms bei staublungenerkrankungen. (The incidence of emphysema and bronchogenic carcinoma in silicosis). Prax Pneumol 26: 145-58, 1672	A 378
<b>26:</b> 145-58, 1972.	
PALMER KNV and DIAMENT ML: A comparison of pulmonary function in bronchial asthma and chronic obstructive bronchitis. Thorax 25: 101-4, 1970.	A 379
PRINAL PROCESSOR CONTRACTOR CONTR	
PERALES G M G: Simposio sobre problemas medicos originados por el tabaquismo.  Alteraciones morfologicas en diferentes organos y sistemas.  Neumol Cir  Thorax Mex 32: 27-29, 1971.	A 380
PHILLIPS A M, PHILLIPS R W and THOMPSON J. L: Chronic cough: analysis of etiologic factors in a survey of 1,274 men. Ann Intern Med 45: 216-231, 1956.	A 381
PHILLIPS J R: Gastroduodenal ulceration in chronic obstructive lung diseases. <u>J Abdominal Surg</u> 8: 306-7, 1966.	A 382
SCHUMAN L M: Epidemiologic approaches and problems in the assessment of causal factors in chronic respiratory disease. J Kentucky Mcd Ass 64: 311-16, 1966.	A 383
SCHUNEMANN V G: Die betreuung der patienten mit einem chronisch bronchitischen syndrom in der kreisstelle für tuberkulose und lungenkrankhelten rostock. Z. Ges Ann Med. 25: 32-55, 1970.	A 384
SILEN W, BROWN W H and EISEMAN B: Peptic ulcer and pulmonary emphysema.  Arch Surg 78: 897-903, 1959.	A 385
	A. A
STEBBINGS J H Jr: A survey of respiratory disease among New York City postal and transit workers. III. Anthropometric, smoking, occupational, and ethnic	4 201

iables affecting the FEV among white males. Environ Res 5: 451-66.

요리를 잃어내려면 사람이 되었다.

# INDUSTRIAL HAZARDS

	Reprint
AHEARN V.P., SANDER O.A., VORWALD A.J., WRICHT G.W. and HARTMAN W.C.:  Emphysema pneumoconioses and compensation.  Arch Environ Health 2: 307- 26, 1961.	A 395
AMDUR M O: Respiratory absorption data and SO <sub>2</sub> dose-response curves. Arch Environ Health 12 (6: 729-32, 1966.	A 396
AYER H E and LYNCH J R: Association of disability and selected occupational hazards.  Arch Environ Health 17: 225-31, 1968.	A 397
BALCHUM O J, DYBICKI J and MENEELY G R: The dynamics of sulfur dioxide inhalation Arch Indust Health 21: 564-9, 1960.	A 398
BALLENGER J. J. McFARLAND C.R., HARDING H.B., KOLL M and HALSTEAD D: The effect of air pollutants on pulmonary clearance. <u>Laryngoscope</u> 78: 1387-97, 1968.	A 399
BOUHUYS A and PETERS J.M.: Control of environmental lung disease. New Engl J Med 283: 573-82, 1970.	A 400
CARNOW B, LEPPER MH, SHEKELLE R B and STAMLER J: Chicago air pollution study. Arch Environ Health 18: 768-76, 1969.	A 401
CHESTER E H, GILLESPIE D G and KRAUSE F D: The prevalence of chronic obstructive pulmonary disease in chlorine gas workers. Am Rev Resp Dis 99: 365-73, 1969.	A 402
DEICHMANN W B, KITZMILLER K V and WITHERUP S: Phenol studies VII. Chronic phenol poisoning with special reference to the effects upon experimental animals of the inhalation of phenol vapor. Am J Clin Pathol 14: 273-7, 1944.	A 403
DEICHMANN W B, WITHERUP S and DIERKER M: Phenol studies XII. The percutaneous and alimentary absorption of phenol by rabbits with recommendations for the removal of phenol from the alimentary tract or skin of persons suffering exposure. J Pharmacol Exp Ther 105: 265-72, 1952.	<b>A</b> 404
DISCHER DP, MASSEY F J and WALLACE RR: Screening by chest photofluorography in Los Angeles. The minifilm circa 1970. Arch Environ Health 22: 92-105, 1971.	A 405
ENTERLINE P E: Epidemiology of coal workers' pneumoconiosis. Indust Med 39: 20-2, 1970.	· A 406
FORBES J D: Cutting and grinding fluids in chronic pulmonary airway disease. J Occup Med 9/8: 421-3, 1967.	A 407
FURLANETO JA, ANDERSON A E and FORAKER A G: Soot emphysema in a locomotive engineer. Arch Environ Health 18: 1008-13, 1969.	A 408
GOFF A M and GAENSLER E A: Asbestosis following brief exposure in cigarette filter manufacture. Respiration 29/1: 83-93, 1972.	A 409
GREEN H.L.: Respiratory protection against particulates problems solved and unsolved. Am Indust Hyg Assoc 26: 203-11, 1965.	A 410
GRIMLEY K W, ADAMS C Neand WALTON R A: Screening for chronic pulmonary disease	 A 411

	List No. 12	Reprint
	GROSS P: Some aspects of pneumoconiosis. Pathol Annu 6: 61-79, 1971.	A 412
	HACKETT R L and SUNDERMAN F W: Pulmonary alveolar reaction to nickel carbonyl.  Arch Environ Health 16: 349-61; 1968.	A 413
	HALLETT W Y: Effect of ozone and cigarette smoke on lung function. Arch Environ Health 10: 295-302, 1965.	i A 414
	HEINRICHS E H and HINS B F: Emphysema rehabilitation program in South Dakota.  S Dakota J Med 21: 31-3, 1968.	A 415
	HEPPER N.G., HYATT R E and FOWLER W.S. Detection of chronic obstructive lung disease. Arch Environ Health 19: 806-13, 1969.	A 416
	HEPPLESTON A G: Emphysema in relation to dust exposure. In: Pneumoconiosis.  Proceedings of the international Conference, Johannesburg 1969, edited by H A  Shapiro, Cape Town Oxford University Press, London, 312-4, 1970.	A 417
:	HORTON A W. TYE R and STEMMER K L: Experimental carcinogenesis of the lung.  Inhalation of gaseous formaldehyde or an aerosol of coal tar by C <sub>3</sub> H Mice.  J Nat Cancer Inst 30: 31-43, 1963.	A 418
	LEWIS C E and KERBY G R: An epidemic of polymer-fume fever. J Am Med Ass 191: 375-8, 1965.	A 419
	LOB M: Pathologie respiratoire professionnelle en dehors de la silicose. Therapeutische Umschau/ Revue thérapeutique 27: 10-6, 1970.	A 420
	MORGAN W K: Coal workers' pneumoconiosis. Am Indust Hyg Ass J 32: 29-34, 1971.	A 421
	MURPHY R L, FERRIS B G, BURGESS W A, WORCESTER J and GAENSLER E A:  Effects of low concentrations of asbestos. Clinical, environmental, radiologic  and epidemiologic observations in shippard pipe coverers and controls. New Engl  J Med 285: 1271-8, 1971.	A 422
	MURPHY S D, LENG J K, ULRICH C E and DAVIS H V: Effects on animals of exposure to auto exhaust. Arch Environ Health 7: 60-70, 1963.	A 423
,	NAEYE R L and DELLINGER W S: Coal workers' pneumoconiosis. Correlation of roentgenographic and postmortem findings. J Am Med Assoc 220(2): 223-7, 1972.	A 424
	NIDEN: A H: Effects of ammonia inhalation on the terminal airways. In: Current Research in Chronic Respiratory Diseases. Proceedings of the 11th Aspen Emphysema Conference, Aspen, Colo. U.S. Department of Health, Education, and Welfare, Public Health Service Publication 1879: 41-4, 1969.	A 425
	PATTLE R E and BURGESS F: Toxic effects of mixtures of sulphur dioxide and smoke with air. J Path Bact 73: 411-9, 1957.	A 426
	PENMAN R W: A summary of the conclusions from an international conference on coal workers' pneumoconiosis. Am. Rev Resp Dis 102: 243-7, 1970.	A 427
	SARIC M and GOLDSMITH J R: Methods for investigating the relationship of chronic bronchitis and emphysema to occupational exposures. In: XIVth International Congress of Occupational Health, Volume II, edited by P Sangro, G Akoun, and H L Beale, Excerpta Medica Foundation, Amsterdam, 741-51, 1964.	A 428
	STOKINGER H E, WAGNER W D and DOBROGORSKI O J: Ozone toxicity studies. III.  Chronic injury to lungs of animals following exposure at a low level. Arch indust	

VIIKERI M, JÄÄSKELAINEN J and TÄHTI E: Ultrasonic examination of pleural thickenings and calcifications in occupational asbestosis. Dis Chest 54: 17-20, 1968.

A 430

#### PULMONARY FUNCTIONAL TESTS

		wet	orint
•	BEECKMANS J M: The deposition of aerosols in the respiratory tract. II. Deposition in cigarette smoking. Canad J Physiol Pharmacol 43: 707-14, 1965.	A	431
	BOUHUYS A, GEORG J, JONSSON R, LUNDIN G and LINDELL S-E: The influence of histamine inhalation on the pulmonary diffusing capacity in man. J Physiol 152: 176-81, 1960.	· A	432
	BOUHUYS A, JONSSON R, LICHTNECKERT S, LINDELL S-E, LUNDGREN C, LUNDIN  G and RINGQUIST T R: Effects of histamine on pulmonary ventilation in man.  Clin Sci 19: 79-94, 1960.		433
	BRISCOE W A and DUBOIS A B: The relationship between airway resistance, airway conductance and lung volume in subjects of different age and body size.		,
·	J Clin Invest 37: 1279-85, 1958.  BURGER JR E J and MACKLEM P: Airway closure: demonstration by breathing 100%	A	434
•	O <sub>2</sub> at low lung volumes and by N <sub>2</sub> washout. <u>J Appl Physiol</u> 25: 139-48, 1968.  CALDWELL E J and FRY D L: Pulmonary mechanics in the rabbit. <u>In:</u> Current Research	Α	435
•	in Chronic Respiratory Diseases. Proceedings of the 11th Aspen Emphysema Conference, Aspen, Colo. U.S. Department of Health, Education, and Welfare, Public Health Service Publication 1879: 307-20, 1969.	A	436
<b>)</b>	CASARETT L J: Role of radioactive substances in effects of smoking. In: Toward a Less Harmful Cigarette. National Cancer Institute Monograph 28, edited by E L Wynder and D Hoffmann, U.S. Department of Health, Education, and Welfare, Public Health Service, National Cancer Institute, Bethesda, Maryland, 199-209, 1968.	A	437
	DALHAMN T, EDFORS M-L and RYLANDER R: Retention of cigarette smoke components in human lungs. Arch Environ Health 17: 746-8, 1968.	A	438
	DAVIES CN: The handling of particles by the human lungs. Brit Med Bull 19: 49-52, 1963.	A	439
	DuBOIS A B, BOTELHO S Y and COMROE JR J H: A new method for measuring airway resistance in man using a body plethysmograph: values in normal subjects and in patients with respiratory disease. J Clin Invest 35: 327-35, 1956.	A	440
	FERIN J: Pulmonary clearance after deposition of aerosols. Lekarske Prace 5: 5-67, 1966.	. А	44 i
	GAVRILESCU N, TECULESCU D, STANESCU D and CONSTANTIN I: Pulmonary mechanics in young healthy men normal values for some less frequently used parameters.  Int Z Angew Physiol 24: 194-207, 1967.		442
•	HALE F C; OLSEN C R and MICKEY JR M R: The measurement of bronchial wall componed Am Rev Resp Dis 98: 978-87, 1968.		443
	HERZOG H, KELLER R, MAURER M, BAUMANN H R and NADJAFI A: The distribution bronchial resistance in obstructive pulmonary disease and in dogs with artificially induced tracheal collapse. Public Health Service Publication No. 1787: 55-91, 1968.		444
	HYATT R E and WILCOX R E: Extrathoracic airway resistance in man. J Appl Physiol 16: 326-30, 1961.	A	445
	LaBELLE C W, LONG J E and CHRISTOFANO E E: Synergistic effects of aerosols.  Arch Indust Health II: 297-304, 1955.	, <b>A</b>	446

List No. 13		1	Reprint
LANGER G and FISHER M A: Concentration  Arch Indust Health 13: 372-8, 1956.	on and particle size of cigarett	e-smoke particles.	A 447
LOUDON R G: Determinants of expiratory a Respiratory Diseases. Proceedings Aspen, Colo. U.S. Department of Health Service Publication 1879: 275-	of the 11th Aspen Emphysema eaith, Education, and Welfare	Conference,	A 448
LOUDON R G and SHAW G B: Mechanics of obstructive respiratory disease. Am	1 Rev Resp Dis 96: 666-77, 19	9 <b>67.</b>	A 449
MACKLEM PT: Obstruction in small airway 52: 721-4, 1972.	ays - a challenge to medicine.	Am J Med	<b>A</b> 450
MAISEL J C, SILVERS G W, MITCHELL R dynamic expiratory collapse. Am Re		al atrophy and	A 451
MELLORS R C: Microscopic localization of tracts of animals exposed to cigarette 2: 325, 1958.			A 452
MITCHELL R I: Controlled measurement o		the respiratory	A 453
NADEL JA, SALEM H, TAMPLIN B and during inhalation of sulfur dioxide.			A 454
SAFFIOTTI U, BORG SA, GROTE MI and carcinogens in the lungs. Studies in duction. The Quarterly: 10-7, 1	an experimental model for lun		A 455
SIMONSSON BG, JOCOBS FM and NADE the cough reflex in the increased responsitive airway disease. J Clin Is	ponsiveness of airways in pati		A 456
WOLFF W A, PURDOM E G and ISENHOW tracers in cigarette smoke. N Caro	ER J A: The use of radioiso lina Med J 15: 159-63, 1954.	topes as	A 457
			•

some observations on free charges in signrette smoke. Acta Med Scand: 170 (Suppl.

369): 119-20, 1961.

#### Additional Bibliographic List No. 15

#### CARDIOPULMONARY EFFECTS OF CIGARETTE SMOKE

	Reprint
ANON: Cigarettes and atherosclerosis. Practitioner 202(1208): 195-6, 1969.	A 473
BARER G.R. HOWARD P and McCURRIE J.R.: The effect of carbon dioxide and changes in blood pH on pulmonary vascular resistance in cats. Clin Sci 32: 361-76, 1967.	A 474
BOREN H G and KRAHL V E: Ventilatory, neural, and other factors influencing the pulmonary microcirculation of the frog. Bibliotheca Anatomica 7: 86-91, 1965.	A 475
COLLOQUES INTERNATIONAUX DU CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE: Le rôle de la paroi artérielle dans l'athérogénèse. 169: 7-139, 1968.	A 476
CONROY J P: Smoking and the anesthetic risk. Anesth Analg Current Researches 48: 388-400, 1969.	A 477
DENNISON JR A D: Athletics and the heart. J Indiana St Med Ass : 850-5, 1964.	A 478
DOYLE JT: The prevention of coronary heart disease. In: Coronary Heart Disease, edited by Albert N. Brest, F A Davis Company, Philadelphia, 305-16, 1969.	A 479
ECHT R: Estimations of pulmonary surface activity in sedentary and exercised rats.  IN: Current Research in Chronic Fespiratory Diseases. Proceedings of the lith Aspen Emphysema Conference, Aspen, Colo. U.S. Department of Health, Education, and Welfare, Public Health Service Publication 1879; 323-31, 1969.  EMMANUEL G E, ROSENBLUTH A, ANSARI I, VILLANO R and CARDUCCI R:	<b>A 4</b> 80
Evidence of active alveolar closure in the human lung. IN: Current Research in Chronic Respiratory Diseases. Proceedings of the 11th Aspen Emphysema Conference, Aspen, Colo. U.S. Department of Health, Education, and Welfare, Public Health Service Publication 1879: 287-96, 1969.	° <b>A 481</b>
FILLEY G.F. BECKWITT H.J. REEVES J.T and MITCHELL R.S.: Chronic obstructive bronchopulmonary disease. (I. Oxygen transport in two clinical types. Am J. Med 44: 26-38, 1968.	A 482
FOWLER NO: Chronic cor pulmonale. Gentatrics 22/6:156-71, 1967.	A 483
GALY P and PERRIN L-F: Le surfactant alveolaire dans le poumon normal et pathologique. Rev Lyonnaise Med 17: 677-84, 1968.	A 484
GERLACH U: Zur pathogenese der arteriosklerose. Therapiewoche 2: 59-63, 1969.	A 485
GIAMMONA S T, KERNER D and BONDURANT S: Effect of oxygen breathing at atmospheric pressure on pulmonary surfactant. J Appl Physiol 20: 855-8, 1965.	A 486
GROVER R.F., ALEXANDER J.F. and HARTLEY L.H. Myocardial function in corpulmonale. In: Current Research in Chronic Respiratory diseases. Proceedings of the 11th Aspen Emphysema Conference, Aspen, Colo. U.S. Department of Health, Education, and Welfare, Public Health Service Pub-	
lication 1879; 121-7, 1969.	A 487

HURST J W: Prevention of coronary atherosclerosis. In: The Heart. Arteries and

	List No. 15	Reprint
<b>T</b>	KILBURN KH, ASMUNDSSON T. BRITT R C and CARDON R: Effects of breathing 10 per cent carbon dioxide on the pulmonary circulation of human subjects. Circulation 39: 639-53, 1969.	L A 489
	LLOYD JR T C: Hypoxic pulmonary vasoconstriction: role of perivascular tissue. <u>J Appl Physiol</u> 25: 560-5, 1968.	A 490
	MALISZEWSKI TF and BASS DE: 'True' and 'Apparent' thiocyanate in body fluids of smokers and nonsmokers. J Appl Physiol 8: 289-91, 1955.	A 491
	MASSUMI R A and NUTTER DO: Cardiac arrhythmias associated with Cheyne-Stokes respiration: a note on the possible mechanisms. Dis Chest 54: 21-2, 1968.	A 492
	McLAUGHLIN R.F., TYLER W.S and CANADA R.O: A study of the subgross pulmonary anatomy in various mammals. Am J. Anatomy 108: 149-65, 1961.	A 493
	MENDENHALL JT: Evaluation and management of pulmonary insufficiency in surgical patients. Surg Clin N Amer 48: 773-8, 1968.	A 494
	NAEYE R L and LAQUEUR W A: Chronic cor pulmonale. Its pathogenesis in Appalachian bituminous coal workers. Arch Path 90: 487-93, 1970.	A 495
• •	OUTHRED K.G., McKENZIE H. I and WHITE K.H.: Some findings on the post-mortem association of right ventricular wall thickness and chronic obstructive bronchitis, pneumoconiosis, emphysema and tuberculosis. Med J. Aust 2: 950-4, 1970.	A 496
)	PAUTRAT J: Les complications cardiaques des broncho-pneumopathies. G M de France 1: 4993-8, 1967.	A 497
••	PENMAN R W, HOWARD P and STENTIFORD N H: Factors influencing pulmonary gas exchange in patients with acute edematous cor pulmonale due to chronic lung disease  Am J Med 44: 8-15, 1968.	A 498
	PHILLIPS R W: The electrocardiogram in cor pulmonale secondary to pulmonary emphysema: A study of 18 cases proved by autopsy. Am Heart J 56: 352-71, 1958.	A 499
	PLATTS M M, HAMMOND J D and STUART-HARRIS C H: A study of cor pulmonale in patients with chronic bronchitis. Q J Med 29: 559-74, 1960.	A 500
	READ J, LEE J H and PAIN M C: Two groups of subjects with obstructive lung disease, defined by pulmonary vascular reactivity. Public Health Service Publication No. 1787: 229-39, 1968.	<b>A</b> 501
	SAID SI: Role of pulmonary surfactant in health and disease. Med Clin North Am 51: 391-402, 1967.	A 502
•	SANDERS W. L.: Heart disease and pneumoconiosis. Thorax 25: 223-5, 1970.	A 503
	SEKULIC S M, HAMLIN III J T, ELLISON R G and ELLISON L T: Pulmonary surfactant and lung circulation in experimental atelectasis. Am Rev Resp Dis 97: 69-75, 1968	I. A 504
	SHERMAN M: Smoking and heart disease. National Advisory Heart Council : 1-55, 196	2. A 505
•	SHMOCK C L, MITCHELL R S, POMERANTZ B, PRYOR R, and MAISEL J. C: The electrocardiogram in chronic airways obstruction. The role of bronchitis and	. A 506
	emphysema. Chest 60: 335-40, 1971.  STAMLER J: Chronic pulmonary heart disease (Cor pulmonale). In: Lectures on Preventive Cardiology by I Stamler, Grune, and Stratton, Inc., New York,	

STONE R M, GINSBERG R J, COLAPINTO R F and PEARSON F G: Bronchial artery regeneration after radical hilar stripping. Surg Forum 17: 109-10, 1966.

A 508

VUGA D: Risposta del tratto bronchiolo-alveolare al fumo di sigaretta. Riflessi sul surfactant e sulla chiarificazione polmonare. Riforma Medica 82: 203-8, 1968.

A 509

WEST JB, GLAZIER JB, HUGHES JM and MALONEY JE: Effect of gravity on the morphology of pulmonary capillaries and alveoli. In: Current Research in Chronic Respiratory Diseases. Proceedings of the 11th Aspen Emphysema Conference, Aspen, Colo. U.S. Department of Health, Education, and Welfare, Public Health Service Publication 1879: 135-7, 1969.

A 510

ANDERSON 1967 (A 225) 219 ABBOTTO, HOPKINS, VAN FLEIT and ROBINSON 1953 (43) 23, 21 ANDERSON 1967 (A 226) 219 ABELSON 1968 (827) 174, 160 ANDERSON 1968 (307) 71, 69 ABLIN 1971 (A 122) 213 ANDERSON (A 261) 221 ADAMS 1969 (A 224) 219 ANDERSON 1964 (1005) 193 ANDERSON 1962 (230) 54, 52 ADDINGTON 1970 (384) 93, 91 ADELMAN 1967 (A:1): 205 ANDERSON: 1971 (738) 143 ADELSTEIN 1967 (344) S1 ANDERSON 1962 (385) 93, 84 AGUZZI, WOOLF and PATERSON 1966 ANDERSON and FERRIS 1965 (596) 123, 121 ANDERSON and FERRIS 1966 (A 227) 219 (491):111, 104 AHEARN, SANDER, VORWALD, WRIGHT ANDERSON, FERRIS and ZICKMANTEL 1965 (492) and HARTMAN 1961 (A 395) 229 111, 104 ANDERSON, ZICKMANTEL and FERRIS 1963 (386) AIACHE, CHEMINAT, LAKATOS and 93, 84 MOLINA 1972 (345) 81, 79 ALBERT, ALESSANDRO, LIPPMANN ANDERSON, DUNNILL and RYDER 1972 (59) 28, 25 and BERGER 1971 (666) 135, 134 ANDERSON and SHEPHARD 1969 (737) 143, 141 ANDERSON, ANDERSON, HERNANDEZ and ALBERT, LIPPMANN and BRISCOE 1969a (667):135, 131 FORAKER 1964 (A 13) 206 ANDERSON and WILLIAMS: 1969 (739) 143, 141 ALBERT, LIPPMANN, PETERSON, BERGER, SANBORN and BOHNING ANDOSCA 1966 (387) 93, 85 ANON (837) 184 1973 (668) 135, 134 ALBERT, SPIEGELMAN, SHATSKY ANON (840) 184 and LIPPMANN 1969b (669) 135, 134 ANON (839) 184 ALLEN and SABIN 1971 (621) 128, 125 ANON (838) 184 ANON: 1964 (900):181 ALLEN, ANGERMANN and FACKLER 1968 (1) 11, 8 ANON 1967 (A 262) 221 ALLI 1972 (24) 19, 18 ANON 1967 (A 335) 225 ALVAREZ 1964 (58) 28, 25 ANON 1968 (902) 187 AMDUR 1966 (A 396) 229 ANON 1969 (A 473) 235 AMERICAN CANCER SOCIETY 1967 ANON 1970: (901) 187 ANON: 1971 (A 336) 225 AMERICAN CANCER SOCIETY, INC ANON 1972 (A 187) 217 and NATIONAL CANCER INSTITUTE ANTHOMSEN, BASS, ORIOL, PLACE and BATES U S PUBLIC HEALTH SERVICE 1960 1968 (201):48, 47 ANTONIO 1963 (44) 23, 21 (A 186) 217 ARCHER, CARROL, BRINTON and SACCOMANNO AMERICAN CANCER SOCIETY NEWS SERVICE RELEASE 1968 (967) 191 1964 (388) 93, 92 ARCHIVES OF OTOLARYNGOLOGY 1969 (903) 187 AMERICAN COLLEGE OF CHEST PHYSICIANS 1966 (1044) 196 ARING 1969 (1046) 196 AMERICAN DRUGGIST 1970 (1139):201 ASIAN MEDICAL JOURNAL 1964 (968) 191 AMERICAN JOURNAL OF PUBLIC ASHFORD, BROWN, DUFFIELD, SMITH and FAY HEALTH 1970 (1045) 196 1961 (444) 101, 100 ASHFORD, BROWN, MORGAN and RAE 1968 (A 337) AMERICAN MEDICAL ASSOCIATION 1966 (S98) 187 225 ASHFORD, MORGAN, RAE and SOWDEN 1970 (445) AMERICAN REVIEW OF RESPIRATORY DISEASES 1970 (899) 187. 101, 100 ASHLEY 1969 (A 158) 215 ANDERSON, AZCUY, BATCHELDER ASMUNDSSON and KILBURN 1969 (316) 75, 73

ASTIN 1972 (A 299):223

AUERBACH 1964 (904) 187

(770) 147, 145

AUERBACH 1967 (105) 35, 31

AUERBACH 1969 (106):35, 31

128, 126

ATTINGER, GOLDSTEIN and SEGAL 1958 (622).

AUERBACH, HAMMOND and GARKINKEL 1968

and FORAKER 1964 (104) 35, 31

ANDERSON and FORAKER 1967 (A 12)

ANDERSON, FURLANETO and

ANDERSON: 1963 (306) 71, 69

FORAKER 1970 (25) 19, 16

ANDERSON, HERNANDEZ, ECKERT

and FORAKER 1964 (26):19, 15

ANDERSON, HERNANDEZ, HOLMES

and FORAKER 1966 (27) 19, 15

BALINT, BONDURANT and KYRIAKIDES 1971 (788)

BALLENGER, DAWSON, DeRUYTER and HARDING

BALLENGER, McFARLAND, HARDING, KOLL and

152, 150

1965 (A 458) 234

BANYAI 1961 (A 73) 210

BANYAI 1962 (A 36) 208

BANYAI 1966 (A 37) 208

BANYAI 1967 (1049) 196

BARACH 1964 (319) 75, 72

BALLENGER 1960 (670) 135, 131

HALSTEAD 1968 (A 399) 229.

BARACH 1966 (A 74) 210 GARFINKEL 1961 (173) 44, 43 BARACH 1967 (1140) 201 AUERBACH, STOUT, HAMMOND and BARACH 1970 (1141) 201 GARFINKEL 1962a (174) 44, 43 BARACH 1971 (318) 75, 72 BARACH 1967 (A 300) 223 AUERBACH, STOUT, HAMMOND and GARFINKEL 19625 (175) 44, 43 BARACH and SEGAL 1968 (320) 75, 72 AUERBACH, STOUT, HAMMOND and BARCLAY 1966 (969) 191 GARFINKEL 1963 (29) 19, 16 BARCLAY 1969 (A 263) 221 AUERBACH, STOUT, HAMMOND and BARCKLEY (1050) 196 GARFINKEL 1964 (905) 187 BARER, HOWARD and McCURRIE 1967 (A 474) 235 AUERBACH, STOUT, HAMMOND and BARNES and SIMPSON 1968 (493) 111, 109 GARFINKEL 1966 (30) 19, 16 BASS, WHITCOMB and FORMAN 1970 (A 75) 210 BATES 1965 (A 38):208 AUERBACH, STOUT, HAMMOND and GARFINKEL 1968 (176) 44, 43 BATES 1968 (A 264) 221 AVIADO 1972 (623) 128, 127 BATES 1968 (A 265) 221 AVIADO 1969 (110) 35, 32; (624) 128, BATES 1970 (A 39) 208. BATES 1972 (A 230) 219 AVIADO and McKINNEY 1969 (111) BATES, BOUCOT and DORMER 1955 (740) 143, 141 BATES, GORDON, PAUL, PLACE, SNIDAL and AVIADO and PALECEK 1967 (625) WOOLF 1966 (202) 48, 46 BATTISTA and KENSLER 1970a (671) 135, 134 AVIADO, SADAVONGVIVAD and BATTISTA and KENSLER: 1970b (672) 135, 134 BAUMANN 1968 (A 40) 208 CARRILLO: 1970 (112):35, 33; (626) BAUMBERGER 1927 (A 459) 234. AVIADO and SAMANEK 1965 (627) BEAUDRY 1968 (494) 111, 104 BECKERMAN: 1967 (841) 184 BECKLAKE, GOLDMAN, BOSMAN and FREED 1957 AVIADO, SAMANEK and FOLLE 1966 (115):35, 33 BEDELL and SEEBOHM 1959 (A 76) 210 AVIADO, WATANABE and BELEJ 1973 BEDELL and SEEBOHM 1966 (60) 28, 24 AYER and LYNCH 1968 (A 397) 229 BEDROSSIAN 1972 (A 188) 217 AYRES 1968 (1047) 196 BEECKMANS 1965 (A 431) 232 AYRES 1970 (A 228) 219 BEERENS 1967 (A 231) 219 BACHMAN 1970 (1048) 196 BEHNKE: 1970 (780) 149, 148 BADGER and GREEN 1967 (A.71) 210 BELCHER 1966 (1051):196 BAILEY, HABERMAN and SHEINBERG BELL (1142) 201 BELL 1970 (61) 28, 26 BAIR, PORTER, BROWN and WEHNER BELLOMY (1052) 196 BENNETT 1972 (835) 182, 161 BERARD and EMONOT 1972 (346) 81, 79 BAKER: 1970 (A-72):210 BAKER, OSCHERWITZ, CORLIN, BERITIC 1964 (842) 184

BERNDT 1962 (A 339) 225

BERNIER 1957 (A 189) 217.

BEUMER 1963 (1053) 196

BALCHUM, FELTON, JAMISON, GAINES, BIERSTEKER 1968 (495) 111, 106

BEST, WALKER, BAKER, DELAQUIS, McGREGOR

and McKENZIE 1967 (308) 71, 69

BICKERMAN and BARACH 1954 (45) 23, 22

BIEGEL and KRUMHOLZ 1968 (241) 57, 56

AUERBACH, HAMMOND, GARFINKEL

and BENANTE 1972 (28) 19, 18, 17

UERBACH, HAMMOND, KIRMAN and GARFINKEL 1967 (107) 35, 31

AUERBACH, HAMMOND, KIRMAN and

GARFINKEL and STOUT 1967 (254)

GARFINKEL and STOUT 1968 (109):

GARFINKE L. 1970 (108):35, 31

AUERBACH, HAMMOND, KIRMAN,

NUERBACH, HAMMOND, KIRMAN,

AUERBACH, STOUT, HAMMOND and

35, 31

127

35, 32

128, 127

128, 127

128, 127

(628) 128

(787) 152, 151

1969 (317) 75, 74

1969 (113) 35, 32

1970 (A 338) 225

1960 (A 398) 229

BALCHUM 1962 (A 229) 219

JARBOE, TEISCH and NICHAMAN

BALCHUM, BUCKLEY, SHERWIN and

BALCHUM, DYBICKI and MENEELY

GARDNER 1965 (114):35, 33

BIERSTEKER 1969 (496):111, 106 BIERSTEKER 1969 (497) 111, 106 BISHOP 1963 (A 2) 205 BLACK and HYATT 1971 (A: 340) 225 BLACK, HYATT and STUBBS: 1972 (62) 28, 26 BLACKBURN, TAYLOR, PARLIN, KIHLBERG and KEYS 1965 (390) 93, 91BRINKMAN and BLOCK 1966 (394) 93, 90 BLAIR, HENRY and EHRLICH 1969 (116) BRINKMAN and BLOCK 1968 (203) 48, 47 35. 32 BLAKE 1964 (A 77) 210 BLUE 1970 (347) 81, 79 BLUMBERG 1969 (A 190) 217. BLUNDI 1969. (A.191) 217. BOAKE: 1958 (242) 57, 56 BOATMAN and MARTIN: 1965 (117) 35, 34 BRITISH MEDICAL JOURNAL 1971 (906) 187 BOCHE and QUILLIGEN 1959 (673) 135, 134 BOIN 1969 (1006) 193 BOKHOVEN and NIESSEN 1961 (118) 35, 33 BONDURANT 1960 (789) 152, 150 BONNELL 1955 (119) 35, 34 BONNELL, KAZANTZIS and KING 1959 (120) 36, 34 BONOMO and D'ADDABBO 1964 (217) 51, 50 BONSI (A 192) 217 BOREN 1965 (121) 36, 31 BOREN, BUSET, CORPE, GRZYBOWSKI, NEWMAN, ORGANICK, and LESTER 1966 (A 78):210 BOREN, DELAHANT and STEENKEN 1965 (122) 36, 33 BOREN, KORY and SYNER 1966 (597) 123, 120 BOREN and KRAHL 1965 (A 475) 235 BORGATTA and EVANS 1968 (2) 11, 8; (970) 191 BOUCOT 1968 (1007) 193 BOUCOT, COOPER and WEISS 1962 (391) BURNS and MAY 1967 (A 342) 225 93, 87 BOUDIK, GOLDSMITH, TEICHMAN and KAUFMANN 1970 (498) 111, 108 BOUHUYS, BARBERO, SCHILLLING and VAN DE WOESTIJNE 1969a (499) 111, 107 BOUHUYS, GEORG, JÖNSSON, LUNDIN and LINDELL 1960 (A 432) 232 BOUHUYS, JONSSON, LICHTNECKERT, LINDELL, LUNDGREN, LUNDIN and RINGQUIST 1960 (A 433)232 BOUHUYS and PETERS 1970 (A 400) 229 BOUHUYS, SCHILLING and VAN DE WOESTIJNE 19698 (500) 111, 107 BCUHUYS, WOLFSON, HORNER, BRAIN and ZUSKIN: 1969 (392) 93, 90 BOUSHY, KOHEN, BILLIG and HEIMAN

1968 (47) 23, 21

BOUSHY and LEWIS 1964 (46) 23, 21 BOWER: 1961 (393):93, 91. BRILLE 1969 (501) 111, 106 BRILLE 1970 (502) 111, 106 BRILLE, COUDRAY, ISRAEL-ASSELAIN, LE MELLETIER, MICHEL, MIGUERES. PHAM and VOISIN 1970 (503) 111, 106 BRINKMANN and BLOCK 1972 (204) 48, 47 BRINKMAN and BLOCK 1972 (395):93: 90 BRINKMAN and COATES 1962 (396) 93, 90 BRISCOE and DUBOIS 1958 (A 434) 232 BRITISH MEDICAL JOURNAL 1967 (843) 184 BRITISH MEDICAL JOURNAL 1971 (844):184 BROCARD 1969 (A 301) 223 BRODINE 1972 (A 232) 219 BROWN 1958 (A 302) 223 BROWN and CAMPBELL 1961 (A 341) 225 BROWNING 1967 (845):184 CAHAN and KIRMAN 1968 (124) 36,31 BROWNING and THORP 1969 (1054) 196 BROWNLEE (1143) 201 BRUN, BIOT and KOFMAN 1969 (A 123) 213 BRUN, FOFMAN, G UICHARD and PERRIN-FAYOLLE 1967 (A 14) 206 BRYSIEWICZ, BULUK, CESARZ-FRONCZYK, KORDECKA, LESZCZYŃSKI, ŁUKJAN and SADOKIERSKA 1970 (504) 111, 107 BRYSON and BISCHOFF 1969 (A 193) 217 BUCKLEY and LOOSLE 1969 (123) 36, 33 BUECHNER, BROWN and TRETOLA 1964 (63) 28, 27 BUELL, DUNN and BRESLOW 1967 (321) 75, 73 BULLETIN OF THE LOS ANGELES COUNTY MEDICA ASSOCIATION 1968 (846) 184 BURCH 1961 (971) 191 BURGER: 1968 (A 435) 232 BURGESS: (972):191 BURGESS and BURGESS 1966 (48) 23, 21 BURROWS 1962 (64) 28, 25 BURROWS 1966 (A 303) 223 BURROWS 1967 (741) 143 BURROWS 1968 (A 41) 208 BURROWS 1969 (A 304) 223 BURROWS and EARLE 1964 (322) 75, 73 BURROWS and EARLE 1969 (205) 48, 47 BURROWS, FLETCHER, HEARD, JONES and WOOTLIFF 1966 (A 266) 221 BURROWS and KETTEL 1969 (A 267) 221 CAIRD and AKHTAR: 1972 (206):48, 46 CALDWELL and FRY 1969 (A 436) 232 CALIFORNIA GOVERNOR'S ADVISORY COMMITTEE ON CIGARETTE SMOKING AND HEALTH 1964 (1055) 196 CALIFORNIA MEDICAL ASSOCIATION 1967 (1056) 19 CALIFORNIA MEDICAL ASSOCIATION 1968 (A 343) 2 CAMNER and PHILIPSON 1971 (674) 135, 132

COLLEGE HEALTH 1965 (1011) 193

COLLEY and REID 1970 (450) 101, 99 COLLOQUES INTERNATIONAUX DU CENTRE

NATIONAL DE LA RECHERCHE

SCIENTIFIQUE 1968 (A 476) 235 COLP, PARK and WILLIAMS: 1970 (A 43):208

COMSTOCK, BROWNLOW, STONE and

SARTWELL 1970 (399):94, 87

COLLEY and HOLLAND 1967 (449) 101, 98

CECILIONI 1970 (977):191 CEDERLÖF 1966 (505) 112, 105

112, 105

112, 110

(508) 112, 105

(509) 112, 105

CEDERLÖF 1967 (506) 112, 105

CEDERLOF, FRIBERG and HRUBEC 1969 (507)

CEDER LOF, FRIBERG, JONSSON and KAIJ 1965

CEDERLOF, FRIBERG, JONSSON and KAIJ 1966

CELIS, DIAZ, GOROCICA and CANO 1968 (510)

CAMNER and PHILIPSON 1972 (675) 135. 132 CAMNER, PHILIPSON and ARVIDSSON 1971 (676) 135, 132 CAMNER, PHILIPSON and ARVIDSSON 1973 (677):135, 132 CAMNER, PHILIPSON and FRIBERG 1972 (678):135, 132 CAMPBELL 1969 (A 15) 206 CANADIAN DEPARTMENT OF NATIONAL HEALTH AND WELFARE 1966a (309) 71, 69 CANADIAN DEPARTMENT OF NATIONAL CHADWICK 1967 (1144) 201 HEALTH AND WELFARE 1966b (310) CANADIAN DEPARTMENT OF NATIONAL CHANG 1957 (178) 44, 41 HEALTH AND WEL FARE 1966c (311) CHANG: 1958 (179) 44, 41 CANADIAN FAMILY PHYSICIAN 1968 (1057):196 CANADIAN JOURNAL OF PUBLIC HEALTH 1965 (973) 191 CANADIAN MEDICAL ASSOCIATION JOURNAL 1963 (1008):193 CANADIAN MEDICAL ASSOCIATION JOURNAL 1964 (974) 191 CANADIAN MEDICAL ASSOCIATION JOURNAL 1967 (975) 191 CANADIAN MEDICAL ASSOCIATION JOURNAL 1969 (1009):193 CANCER NEWS (976) 191 CANDER and BLUMENTHAL 1963 (65) 28, 25 CANTER and LUCHSINGER 1967 (66) 28. 26 CANTER and LUCHSINGER 1968 (67) 28, 26 CAPLAN, SIMON and REID 1966 (446): 101, 100 CARABASI 1963 (A 79) 210 CARABASI 1966 (A 305):223 CAREY, DAWSON and MERRETT 1967 (598) 123, 122 CAREY, PHAIR, SHEPHARD and THOM50N 1958 (A 233) 219 CARIELLO (A 42) 208 CARILLI, KOTZEN and FISCHER: 1973 (68) 28, 25 CARLSON, DYER and JOHNSON 1968 (125) 36, 34 CARLSON, YOKOYAMA and DICKINSON 1972 (126) 36, 34 CARNOW 1968 (951) 190 CARNOW, LEPPER, SHEKELLE and STAMLER 1969 (A 401) 229 CARSON; GOLDHAMER and CARPENTER 1966 (679):135, 133 CASARETT: 1968: (A. 437): 232:

CASARETT 1971 (A 234):219

CATLETT and KIDERA 1969 (599) 123, 121

CASSUTO 1970 (907) 187

DEHAMEL and O'DONNELL 1972 (518) 112, 109

DEICHMANN, KITZMILLER and WITHERUP

DEICHMANN, WITHERUP and DIERKER 1952

DELARUE, WOOLF and STRASBERG 1965 (A 85)

DE LOURDES: FONSECA SANTOS 1971 (A 235)

1944 (A 403) 229

DELARUE 1962 (A 196) 217

DELARUE 1968 (828): 175, 160

(A 404) 229 DELARUE 1962 (1145) 201

210

219

DAVIS and MCCLEMENT 1969 (327) 75, 73 DOWLING, JACKSON and INOUYE 1957 (245)

57, 55

232

DuBOIS, BOTELHO and COMROE 1956 (A 440)

DURDA and SZAFRAŃSKI 1971 (521) 113, 107

DUNNILL and RUDEN, 1971 (182) 44, 42

DYSINGER and LEMON: 1963 (325) 75, 74

DeMARRAIS 1965 (A 349) 225 DENNISON 1964 (A.478) 235 DENSEN, JONES, BASS and BREUER 1967 (402) 94, 87 DENSEN, JONES, BASS, BREUER and REED 1964 (403) 94, 87 DEODHAR 1972 (849) 184 DEUTSCHER and HIGGINS 1970 (75) 29, 27 DEVILLIERS 1969 (A 236) 219 **DEVINE** 1960 (1031) 195 DIAGNOSIS NEWS (1032) 195 DIAMOND, LIPSCOMB and JOHNSON 1971 (637) 129, 125 DIEHL (850) 184 DIEHL 1969 (4):11, 8 DIEHL 1969 (851) 184 DIEHL 1970: (1065):197: DINES 1968 (A 307) 223 DISCHER, MASSEY and WALLACE 1971 (A 405): 229 DKURIC, RAICEVIC and KONSTANTINOVIC 1962 (A 461) 234 DOBRZYNSKI, KISIELEWICZ and KOCIECKA 1970a (519) 112, 107 DOBRZYŃSKI, KISIELEWICZ and KOCIĘCKA 1970ь (520):113, 107 DOLL 1963 (952) 190 DOLL and HILL 1964a (313) 71, 70 DOLL and HILL 1964b (314) 71, 70 DORMANDY 1965 (743):143 DORN 1958 (286) 67, 65 DORN 1959 (287):67, 65 DORN 1960 (288) 67, 65 DA SILVA and HAMOSH 1973 (636) 129, DORN 1961 (289) 67, 65 DOUGLAS and WALLER 1966 (244) 57, 56 DOYLE 1968 (852) 184 DOYLE 1969 (A 350):226 DOYLE: 1969: (A.479):235

CONALLY 1969 (A 346) 225 CONROY 1969 (A 477) 235-COOK 1966 (1059) 196 COOK and WEBB 1966 (791) 152, 151 COOKE 1964 (908) 187 **COOKE** 1970 (1060):196 CORAZZA, MORROW and CHESNEY 1961 (A 81) 210 CORPE, GRZYBOWSKI, MACDONALD, NEWMAN, NIDEN; ORGANICK and LESTER 1965 (A 82) 210 COR WIN 1970 (1061):196 COSIO VILLEGAS 1966 (A 194) 217 COTES, ROSSITER, HIGGINS and | GILSON 1966 (600) 123, 119 COUDRAY, SERISE and FRÉOUR 1969 - (514) 112, 106 COUNCIL OF SCIENTIFIC RESEARCH ON CHRONIC NONSPECIFIC RESPIRATORY DISEASES IN CRACOW, de TAKATS 1965 (1064) 197 POLAND: 1972 (515) 112, 1:07 COURANT 1967 (243) 57,56 CRALLEY 1969 (847) 184 CRENSHAW 1968 (771) 147, 145 CRESS, SPOCK and HEATHERINGTON 1965 (680) 136 CROFTON: 1965 (323) 75, 73 CROMPTON 1968 (634) 128, 125 CROSS, WALZ, PALMER and WARNER 1961 (181) 44, 42 CROXATTO and OEMEDO 1968 (A 16) 206 DIMOND 1972 (1066) 197 CULLEN, KAEMMERLEN, DAOUD and KATZ 1970 (779) 149, 148 CULLEN, KATZ and KAEMMERLEN 1965 (348) 81, 79 CUNNINGHAM, MONTOYE, HIGGINS and KELLER 1972: (A. 347): 225 CURTIS, ASHLEY, RASMUSSEN and CREE 1966 (635) 123, 125 CURTIS, LISKA, RASMUSSEN and CREE 1968 (A 83) 210 D'ANELLI, PROSCIA, SORISTO and CARATTI 1965 (A 348) 225 D'ANELLI, PROSCIA, SORISTO and CARATTI 1968 (349):81 DANOVICH, KOROVINA, SMIRNOW and ANISIMOVA 1969 (517) 112, 108

DAVID (1063):197

DAVID 1967 (A 269):221

DAVIS 1966 (848) 184

(A 460) 234

(401) 94, 92

DAVIES 1963 (A: 439) 232

DAWSON 1966 (400) 94, 92

DEANE 1968 (A 84) 210

1968 (74) 28, 27

DAVIS, BATTISTA and KENSLER 1967

DEANE, GOLDSMITH and TUMA 1965

DeCENCIO, LESHNER and LESHNER

EARLE and BURROWS 1972 (A 308) 223 EBERLY, TYLER and GILLESPIE 1966 (127) 36, 34 EBERT 1959 (A 44) 208 EBERT 1961 (A 270) 221 EBERT and PIERCE 1963 (A 3) 205 ECHT 1969 (A 480) 235 ECKARDT 1964 (A 237) 219 ECKARDT 1972 (404) 94, 84 EDELMAN, MITTMAN, NORRIS, COHEN and SHOCK 1966 (601): 123, 121 EDITORIAL 1964 (819) 160, 165 EDWARDS, McKEOWN and WHITFIELD **1959** (451) 101, 98 EDWARDS 1966 (A 159) 215 福州EDWARDS 1957 (350) 81, 78 EHRLICH and HENRY 1968 (128) 36, 33 EICH, GILBERT and AUCHINCLOSS 1956 (49) 23, 22 EICH, GILBERT and AUCHINCLOSS 1957 (50) 23, 22 EL DIA MEDICO 1967 (853) 184 ELLIOT and TIGHE 1968 (1067) 197 EL-SEWEFY 1970 (522) 113, 109 EL-SEWEFY and AWAD 1971 (523) 113, 109 ELWOOD, PEMBERTON, MERRETT, CAREY and McAULAY 1965 (452) 101, 100 ELY, PEDLEY, HEARNE and STILLE 1970 (405) 94, 87 EMIRGIL, SOBOL, NORMAN, MOSKOWITZ, EMMANUEL, ROSENBLUTH, ANSARI, VILLANO and CARDUCCI 1969 (A 481) ENTERLINE 1967 (A 351) 226 ENTERLINE 1970 (A 406) 229 ENTERLINE and LAINHART 1967 (406) 94, 89 ERIKSSON 1965 (76) 29, 26, 27 ESPINO 1967 (953) 190 EVANG 1964 (854) 184 **EVANG 1969 (855) 184** EVANS, WILKES and DALRYMPLE-SMITH 1969 (453): 101, 99 EVERT 1965 (696) 136, 132 FAIN 1971 (A 87) 211 FAIRBAIRN and REID 1958 (232) 54, 52 FALK and BRISCOE 1970 (909) 187 FALK and BRISCOE 1970 (910) 187 FALK, SISKIND and SMITH 1971 (A 124) 213 FALK, TREMER and KOTIN 1959 (697) 136, 133 **FALLAT** 1970 (77) 29

FALLAT, POWELL, KUEPPERS, NADEL and MURRAY 1971 (79) 29, 26 FALLAT, POWELL, KUEPPERS, LILKER 1973 (78) 29, 26 FARCON and MORALES 1972 (A 309) 223 FEGIZ (A 45) 208 FERIN 1966 (A 441) 232 FERIN 1967 (698) 136 FERIN, URBANKOVA, VICKOVA and REICHRTOVA 1966 (699) 137, 134 FERRIS 1968 (A 238) 219 FERRIS and ANDERSON 1965 (407) 94, 84 FERRIS, ANDERSON and BURGESS 1962 (A 352) 226 FERRIS, BURGESS and WORCESTER 1967 (408) 94, 85 1 FERRIS, BURGESS and WORCESTER 1968 (409) 94, 86 FERRIS, HIGGINS, HIGGINS, PETERS, VAN GANSE and GOLDMAN 1971 (410) 94, 84 FERRIS, HIGGINS, HIGGINS and PETERS 1973 (411) 94, 84 FERRIS, SPEIZER, WORCESTER and CHEN 1971 (1012) 193 FERRIS and WHITTENBERGER 1966 (A 239) 219 FEUEREISL, KRZHIVANEK, REIL, KUBIK and FELKEL 1972 (524) 113, 108 FIESER 1964 (817) 160, 162 FILLEY 1967 (A 310) 223 FILLEY, BECKWITT, REEVES and MITCHELL 1968 (A 482) 235 GOYAL and WADHWANI 1969 (A 86) 211 FINGERLAND, HUSAK and BENFLOVA 1971 (31) 19, 17 FINKLEA, HASSELBLAD, SANDIFER, HAMMER and LOWRIMORE 1971 (412) 94, 90 FINKLEA, SANDIFER and SMITH 1969 (246) **57**, 56 FINLEY 1972 (792) 152 FINLEY and LANDMAN 1972 (793) 152, 151 FISHBEIM 1964 (822): 160, 168 FISHER 1964 (911) 187 FLETCHER 1959 (A 160) 215 FLETCHER 1962 (A 271) 221 FLETCHER 1962 (A 272) 221 FLETCHER 1963 (454) 101, 97 FLETCHER 1963 (A 273) 221 FLETCHER 1965 (A 161) 215 FLETCHER 1965 (A 162) 215 FLETCHER 1967 (A 46) 208 FLETCHER 1967 (A 163) 215 FLETCHER 1968 (207) 48, 47 FLETCHER 1968 (856) 184; (979) 191 FLETCHER 1968 (A 164) 215 FLETCHER 1969 (857) 185

FLETCHER and HORN 1970 (5) 11,

FLETCHER, HORN, AS and RAMSTROM 1971 (1068) 197 FLETCHER, HUGH-JONES, McNICOL and PRIDE 1963 (A 311) 223 FLICK and PATON 1959 (602) 123, 120 FLINT, MAXWELL and RENZETTI **1971** (255) 61, 59. FODOR, GLASS and WEINER FOGARTY 1967 (824) 160, 171 FOGH, FROST and GEORG 1969 (525) 113, 106 FOITZIK, STAEMMLER and POSZICH 1972 (A 197) 217 FOLEY, AMOE, COON and CULVER 1969 (1070) 197 FONTANA (978) 191 FORBES 1967 (A 407) 229 FORD 1971 (1013) 193 FORSTER, ROUGHTON, CANDER, BRISCOE and KREUZER 1957 (744)143 **FOURNIER** and ZIVY 1970 (526) 113, 106 FOWLER 1967 (A 483) 235 FRANKLIN and LOWELL 1961 (603) 123, 120, 121 FRANKLIN, MICHELSON, LOWELL and SCHILLER 1958 (A 88) 211 FRANCES, ALESSANDRO, LIPPMANN, PROCTOR and ALBERT 1970 (700) **137.** 134 FRASCA, AUERBACH, PARKS and JAMIESON 1968a (256) 61, 60 FRASCA, AUERBACH, PARKS and JAMIESON 19685 (257) 61, 60 FRASCA, AUERBACH, PARKS and JAMIESON 1971 (129) 36, 31 FREDERICKSON 1968 (1071) 197 FREDERICKSON 1970 (1072): 197 FREEMAN 1968 (130) 36, 33 FREEMAN, CRANE, STEPHENS and FURIOSI 1968a (131) 36, 33 FREEMAN, CRANE, STEPHENS and FURIOSI 1968b (132) 36, 33 FREEMAN, CRANE, STEPHENS and FURIOSI 1969 (133) 36, 33 FREEMAN and HAYDON 1964 (134) 36, 33 FREOUR and COUDRAY 1971 (6) 11, 8 FREOUR, COUDRAY, ROUSSEL and SERISE 1966 (527) 113, 106 FRUHMANN: 1968 (A. 274) 221 FRY 1954 (328): 75, 73 FULLMER 1968 (218) 51, 49 FULLMER, SHORT, ALLEN and WALKER 1969 (219) 51, 50 FURLANETO, ANDERSON and FORAKER 1969 (A: 408): 229 **FU**TORYANSKY, DETSIL, SHUFLAT SIMONCHIK and BARANSKY 1971

GALY and PERRIN 1968 (A 484) 235 GANDEVIA 1969 (529) 113, 109 GANDEVIA and MILNE 1965 (530) 113, 109 GARCIA ECHEVERRIA (858):185 GARDINER 1963 (A 47) 208 GARFINKEL 1963 (980) 191 GASINSKA and GBUREK 1970 (532) 113, 108 (1069) 197 GAUTHIER 1970 (531) 113, 104 GAVRILESCU, TECULESCU, STANESCU and CONSTANTIN 1967 (A 442) 232 GEDDA 1967 (70) 28, 27 GEISER and STEINMANN 1969 (351) 81, 78 GEISLER 1966 (A 165) 215 GEISLER, BACHMANN, LAUMEN, NOLTE WENTZEL and ROST 1972 (80) 29, 27 GERLACH 1969 (A 485) 235 GIAMMONA 1967 (794) 152, 150 GIAMMONA, KERNER and BONDURANT 1965 (A 486) 235 GIAMMONA, TOCCI and WEBB 1971 (795) 152, 150 **G**ILGEN 1970 (7) 11, 8 GILLESPIE and TYLER 1967 (136) 36, 34 GILLESPIE, TYLER and EBERLY 1964 (137) 36, 34 GILLESPIE, TYLER and EBERLY 1966 (138) 37, 34 GLAUSER and GLAUSER 1969 (139) 37, 34 GLAUSER 1971 (A 125): 213 GLUCKMAN 1969 (A 48) 208 GODBER (859) 185 GOFF and GAENSLER 1972 (A 409) 229 GOLDRING, GREENBURG and RATNER 1968 (140) 37, 34 GOLDSMITH 1965 (A 353) 226 GOLDSMITH 1966 (954) 190 GOLDSMITH 1969 (955) 190 GOLDSMITH: 1970 (A: 240) 219 GOLDSMITH 1971 (912) 187 GOLDSMITH and DEANE 1965 (A 354) 226 GOLDSMITH, GREENBURG, ALTSHULLER, SPICER, CASSELL and LANDSBERG 1966 (A 241) 219 GOLDSMITH, HECHTER, PERKINS and BOHRANI 1962 (413) 94, 92 GOLLI 1970 (533) 113 GONZALES, WEILL, ZISKIND and GEORGE 1968 (A 312) 223 GOTTSCHING and GOTTSCHING 1966 (A355) 226 GOUGH 1955 (A 17) 206 **1005051561 GOUGH** 1968 (A 4) 205 GRAY 1959 (141) 37, 33 GREEN and HORN 1968 (1073) 197 GREEN 1968 (247) 57, 55 GREEN 1968a (258) 61, 58 GREEN 1968b (248) 57, 55 GREEN 1968b (259) 61, 58 GREEN 1968c (260) 61, 58

GREEN 1969 (261) 61, 58 GREEN and CAROLIN 1967 (262) 61, 58 GREEN 1965 (410) 229 GREENBERG, MILNE and WATT 1970 (A 356) 226 GREGG 1966 (956) 190 GREGG 1967 (A 357) 226 GRIMLEY 1969 (A 411) 229 GROSS (A 242) 219 GROSS 1964 (81): 29, 25 GROSS 1966 (A 18) 206 **G**ROSS 1967 (701): 137, 131 GROSS 1967 (913) 187 GROSS 1971 (A 412) 230 GROSS and DeTREVILLE 1969 (142) 137, 34 GROSS and DeTREVILLE 1969 (914) 187 GROSS, DeTREVILLE, BABYAK, KASCHAK and TOLKER 1968a (143) **37**, 33 GROSS, DeTREVILLE, BABYAK, KASCHAK and TOLKER 1968b (144) GROSS, DeTREVILLE, TOLKER, KASCHAK and BABYAK 1969 (263) GROSS and HATCH 1963 (702) 137, 131 GROVER, ALEXANDER and HARTLEY 1969 (A 487) 235 GSELL 1971 (A 198) 217 GUENTER, WELCH, FERGUSON, HENDERSON and HAMMARSTEN 1971 (A 126) 213 GUENTER, WELCH and HAMMARSTEN 1971 (82) 29, 26 GUENTER, WELCH, RUSSEL, HYDE and HAMMARSTEN 1968 (A 127) 213 GUILLERM, BADRE, SAINDELLE, HEE and FLAVIAN 1968 (1033) 195 GUILLERM, BADRE and VIGNON 1961 (703) 137, 134 GUPTA, PURI, INDIRA and DATTA 1968 (A 358) 226 **GUTHRIE** (860) 185 GUYATT and ALPERS 1968 (638) 129, 125 GUYATT, BERRY, ALPERS, BRAMLEY and FLETCHER 1970 (639) 129, 125 HAAGEN-SMIT, BRUNELLE and HARA 1959 (145) 37, 33 HACKETT and SUNDERMAN 1968 (A413) HAENSZEL and HOUGEN 1972 (534) 113, 107 HAGERUP and LARSEN 1971 (535) 113

HAHN, WESTINGHOUSE, SCHREIBER

BRILLMAN, LOMBARD and

EISENBERG 1970 (1146) 201

HAIN 1969 (A 166) 215 HALE, OLSEN and MICKEY 1968 (A 443) 232 HALLETT 1965 (A 414) 230 HAMMARSTEN, WELCH, RICHARDSON, PATTERSON and GUENTER 1968 (A 128) 213 HAMMOND (A 199) 217 HAMMOND 1959 (290) 67, 66 HAMMOND 1964 (291) 67, 66 HAMMOND 1964 (861) 185 HAMMOND 1964 (981) 191 HAMMOND 1965 (1074) 197 HAMMOND 1966 (292) 67, 66 HAMMOND 1967 (982) 191 HAMMOND 1967 (A 243) 219 HAMMOND 1968 (A 244) 219 HAMMOND 1968 (A 200) 217 HAMMOND 1969 (A 201) 217 HAMMOND 1971 (983) 191 HAMMOND, AUERBACH, KIRMAN and GARFINKEL 1970 (146) 37, 31 HAMMOND, BERGLUND and KASLOW 1965 (293) 67, 66; (862) 185 HAMMOND, FENT, REED and WILSON 1964 (294) 67, 66 HAMMOND and GERBER 1965 (295) 67, 66 HAMMOND, GRIETHUYSEN, DIBELER SNEDDON and HALLIGAN 1965 (300) HAMMOND and HORN 1958 (296) 67, 66 HAMMOND and LETTON 1965 (297) 67, 66 HAMMOND and STOCKS 1964 (298) 67, 66 HAMMOND and STREET 1964 (299) 67, 66 HAMMOND and WILSON 1965 (301) 67, 66 HANY, BURCKHARDT and BUHLMANN 1967 (536) 113, 106 HARKEN 1968 (829) 160, 176 HARRIS 1963 (A 313) 223 HARRIS, MENEELY, RENXETTI, STEELE and WYATT 1962 (32) 19, 14 HARRISON and REEVES 1968 (A 359) 226 HARTUNG 1968 (A 275) 221 HARTUNG and MEYER-CARLSTADT 1968 (A 167) 215 HARTUNG 1969 (A 276) 221 HARVEY 1969 (1075) 197 HAUSKNECHT 1962 (A 360) 226 HAY 1967 (8) 11, 8 HAYDON, DAVIDSON, LILLINGTON and WASSERMAN 1967 (147) 37 HAYDON, FREEMAN and FURIOSIS 1965 (148) 37, 33HAYES 1970 (84) 29, 26 HAYES and SUMMERELL 1969 (83) 29, 26 HAYNES, KRSTULOVIC and BELL 1966 · 大学的一个一个一种的政治等的关系 (249) 57, 56 HAWAII MEDICAL ASSOCIATION 1968 (1076) - 197 The state of the s HEAD, HEAD, HUDSON and HEAD 1960

HEALY 1971 (A 202) 217 HEATH 1958 (85) 29, 27 HEDRICK 1969 (9) 11, 8 HEDRICK 1971 (1077) 197 HEIMANN 1967 (A 246) 220 **HEIMANN 1961 (A 245) 220 HEIMBACH 1971 (233) 54, 52** HEINRICHS and HINS 1968 (A 415) 230 HEISE and MYRVIK 1967 (264) 61, 58 HELMHOLZ 1968 (984) 191 HENION 1971 (A 462) 234 HENRY, EHRLICH and BLAIR 1969 (149) 37, 33 HENRY, FINDLAY, SPANGLER and EHRLICH 1970 (150) 37, 33 HEPPER 1969 (1014) 193 HEPPER, BLACK, DINES and MILLER .... **19**68 (A 314) 223 HEPPER, BRANSCOMB, CHERNIACK PIERCE and MITCHELL 1965(A 91) 211 HEPPER, CARR, ANDERSEN, FONTANA, ROSENOW and HANSON 1970 (1078) 197 HEPPER and HYATT 1964 (A 90) 211 HEPPER, HYATT and FOWLER 1969 (A 416) 230 HEPPLESTON 1970 (A 417) 230 **HERBERG** 1966 (86) 29, 25 HERNANDEZ, ANDERSON, HOLMES and FORAKER 1966 (151): 37, 31 HERNANDEZ, ANDERSON, HOLMES and FORAKER 1966 (A 19): 206 **HERXHEIMER and HOLMSTEDT 1959** (A 361) 226 HERZOG 1969 (A 315) 223 HERZOG, KELLER, MAURER, BAUMANN, and NADJAFI 1968 (A 444) 232 HICKEN, HEATH, BREWER and WHITAKER 1965 (772) 147, 145 HICKEY, SCHOFF and CLELLAND 1967 **(32**9) 75, 73 **HIGGINS** (985) 191 HIGGINS 1957 (455) 101, 97 HIGGINS 1959 (456) 101, 98 HIGGINS 1966 (A 247) 220 HIGGINS and COCHRAN 1958 (457) 101, 98 HIGGINS, COCHRANE, GILSON and WOOD 1959 (458) 101, 100 HIGGINS, GILSON, FERRIS, WATERS, CAMPBELL and HIGGINS 1968 (459) 102, 100 HIGGINS, GILSON, FERRIS, WATERS, CAMPBELL and HIGGINS 1969 (460): 102, 100 HIGGINS, HIGGINS, LOCKSHIN and CANALE 1968 (414) 94, 88 HIGGINS and OLDHAM 1962 (461) 102, 100 HIGGINS, OLDHAM, COCHRANE and GILSON 1956 (462) 102, 97

HIGGINS and KJELSBERG 1967 (415) 95, 90 HILDING 1956a (704): 137, 134 HILDING 1956b (705) 137, 134 HILDING 1967 (706) 137 HILDING and FILIPI 1966 (707) 137, 134 HILL 1928 (708) 137, 134 HISLOP 1964(1079) 197 HOFFMAN 1967 (863) 185 HOGG, MACKLEM and THURLBECK 1972 (A 316): 224 HOLCOMB and MEIGS 1972 (416), 95, 86 HOLDAWAY and TURK 1967 (A 362) 226 HOLE and WASSERMAN 1965 (A 129) 213 HOLLAND, MILIC-EMILI, MACKLEM and BATES 1968 (745) 143 HOLLAND 1963 (A 363) 226 HOLLAND 1968 (A 277) 221 HOLLAND and ELLIOTT 1968 (463) 102, 99 HOLLAND, HALIL, BENNETT and ELLIOTT 1969a (464) 102, 99 HOLLAND, HALIL, BENNETT and ELLIOTT 1969ь (465) 102, 99 HOLLAND, HALIL and ELLIOTT 1969c (466) 102, 99 HOLLAND, KOZLOWSKI and BOOKER 1963 (265) 61, 58 HOLLAND and REID 1965 (467) 102, 98 HOLLAND, REID, SELTSER and STONE 1965 (468) 102, 97 HOLLAND and STONE 1965 (417) 95, 87 HOLLAND, WILSON, MORRIS, McCALL and LANZ 1958 (266) 61, 58 HOLMA 1969 (709) 137, 132 HONG, GANDEVIA and LOVELL 1967 (537) 114, 109 HORAK 1969 (10) 11, 8 HORN 1966 (1015) 193 HORN and WAINGROW 1966 (A 364) 226 HOROWITZ 1969 (352) 81, 79 HORTON, TYE and STEMMER 1963 (A 418) 230 HORTON 1965 (1016) 193 HOWARD 1970 (469) 102, 99 HUHTI 1965 (538) 114, 105 HUHTI 1966 (539) 114, 105 HUHTI 1967 (604) 123, 120 HUHTI, RYHANEN, VUOPALA and TAKKUNEN 1970 (540) 114, 105 HUKILL 1963 (1147) 201 HUNT 1968 (A 278) 221 HURST 1959 (A 130) 213 (A 488) 235 HURST HUTCHISON, COOK, BARTER, HARRIS and HUGH-JONES 1971 (88) 29, 26 HUTCHISON, BARTER, COOK, LAWS MARTELLI and HUGH-JONES, 1972 (87) 29, 26 

HUTCHISON 1968 (830) 160, 177

HUTTEMANN, OSWALD, LODE and HUCKAUF 1971 (640) 129, 125 HUTTEMANN, OSWALD, LODE and HUCKAUF 1972 (641) 129, 125 HUTTNER and HUTTNER (A 317) 224 HYATT (986) 192 HYATT (A 203) 217 HYATT, KISTIN and MAHAN 1964 (418) 95, 88 HYATT and WILCOX 1961 (A 445) 232 HYDE and HYDE 1956 (A 49) 208 IDE, SUNTZEEF and COWDRY 1959 (183) 44, 42 **ING**RAHAM (A 248) 220 INOH and AVIADO 1971 (152) 37, 32 INTERNATIONAL SYMPOSIUM IN SAN JUAN 1971 (541) 114, 105 IRNELL and KIVILOOG 1968 (542) 114, 105 ISHIKAWA, BOWDEN, FISHER and · WYATT 1969 (33) 19, 17 ISHIKAWA, FATTAL, ZYLAK, CHERNIACK and WYATT 1968 (A 5) 205 ITO and AVIADO 1968a (153) 37, 33; (642) 129, 127 ITO and AVIADO 1968b (154) 37, 32 IZARD 1967 (A 463) 234 JACKSON and BERNARD 1968 (A 92) 211 **JACOBS** 1970 (834) 161, 181 JAMES 1963 (1080) 197 JAMES 1965 (A 365) 226 JAMES and ROSENTHAL 1961 (302) 68 JAMES 1970 (643) 129, 126 JANCIK 1970 (543) 114, 106 JANCIK and JANCIK-MAK 1972 (544) 114, 108 JARVINEN 1960 (1081) 197 JARVINEN, PATIALA and THOMANDER **19**66 (545) 114, 105 JELEA, ILIE and GAVRILA 1964 (546) 114, 108 **JERRY** and RITCHIE 1964 (A 20) 206 JIMINEZ-DIAZ and CUENZE 1935 (353) 81, 79 **JO**ANNIDES 1965 (A 93) 211 JOHANNSEN and PREIBISZ 1966 (644) 129, 126 JOHNSTON and HARRIS 1965 (A 94) 211 JOHNSTON and KAUFFMAN 1970 (A 50) 208 **JONES** 1970 (1082) 198 JONES and THOMAS 1971 (89) 29, 26 JONES, BURROWS and FLETCHER 1967 (208) 48, 47 JONES, JONES and EDWARDS 1971 (A 95) 211 JONES, BAETJER and REID 1971 (267) 61, 59 The control of the control

ONES, BOLDUC and REID 1972 (268)

JONGKEES 1965 (987) 192 JORGENSEN and SVENSSON 1970 (547) 114, 105 JOULES 1954 (1017) 193 JOURNAL OF THE AIR POLLUTION CONTROL **19**66 (915) 188 JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION 1965 (916) 188 JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION 1965 (957) 190 JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION 1966 (917) 188 JOURNAL OF THE AMERICAN MEDICAL **ASSOCIATION** 1967 (918) 188 JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION 1969 (919) 188 JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION 1969 (988) 192 **JOURNAL OF THE AMERICAN MEDICAL** ASSOCIATION 1972 (1083) 198 JOURNAL OF IOWA MEDICAL SOCIETY **1968** (864) 185 JOURNAL OF IOWA MEDICAL SOCIETY 1969 (1018) 193 JOURNAL OF THE MISSISSIPPI STATE MEDICAL ASSOCIATION (1084) 198 JOURNAL OF THE NATIONAL MEDICAL ASSOCIATION: 1967 (1019): 193 JULICH and RUCKER 1969 (A 51) 208 KAATZSCH 1969 (A 168) 215 KAHN 1966 (303) 68, 65 KAHANA, ARONOVITCH and PLACE 1963 **(51)** 23, 21 KALACIC 1970 (548) 114, 108 KAMINSKI, FANCHER and CALANDRA 1968 **(710)** 137, 133 KAMPMEIER 1964 (818) 160, 164 KARPICK, PRATT, ASMUNDSSON and KILBURN 1970 (184) 44, 43 KATZ and KUNOFSKY 1964 (354) 81, 78 KELSEY, MOOD and ACHESON 1968 (419) 95, 86 KENDIG 1967 (A 366) 226 KENNEDY and ELLIOTT 1970 (712) 137, 134 KENSLER 1967 (711) 137, 133 KENSLER and BATTISTA 1963 (713) 137, 133 KENSLER and BATTISTA 1966 (714) 137, 133 KHOSLA, MEHROTRA and ANKLESARIA 1969 **(781)** 149, 148 KIBELSTIS 1972 (420) 95, 88 KILBURN 1967 (A 96) 211 KILBURN 1967 (715) 137, 131 KILBURN 1968 (716) 137, 131 KILBURN, ASMUNDSSON, BRITT and CARDON 1969 (A 489) 236 KILPATRICK 1971 (A 97) 211 KINSELLA 1962 (A 279) 222 KIRSCHENFELD 1963 (A 98) ZI KISS 1966 (605) 123, 121

KLEIN 1970 (A 6) 205

KLEIN 1970 (989) 192 KLEIN, SALVAGGIO and KUNDUR 1969 (A 367) 226 KLEINFELD, MESSITE, SWENCICKI and SHAPIRO 1968 (421) 95, 87 KLER 1945 (250) 57, 55 KLIDE and AVIADO 1968 (645) 129, 127 KNYVETT 1967 (355) 81, 80 KOLLER 1968 (865) 185 KORDICK, BULBRING and BURN 1952 **(717)** 138, 137 KOTIN 1964 (A 249) 220 KOTIN 1967 (1148) 201 KOTIN 1970 (1149) 201 KOURILSKY, BRILLE and HATTE 1966 (549) 114, 106 KOURILSKY, HAPPERT and FREDY 1966 (A 204) 217 KOWALYSHYN and SATALINE 1969 (A 131) 213 KRAHL and BULMASH 1969 (718) 138, 134 KREYBERG 1969 (A 205) 218 KROEKER and LEON 1962 (356) 81, 80 KRUEGER and SMITH 1958 (719) 138, 132 KRUMHOLZ 1969 (A 318) 224 KRUMHOLZ, CHEVALIER and ROSS 1964 (746) 143, 141 KRUMHOLZ, CHEVALIER and ROSS 1965 (747) 143, 141 KRUMHOLZ, CHEVALIER and ROSS 1965 (773) 147, 145 KRUMHOLZ and HENDRICK 1973 (422) 95, 91 KUCEWICZ 1969a (550) 114, 107 KUCEWICZ 1969b (551) 114, 107 KUEPPERS 1969 (A 132) 213 KUEPPERS 1971 (A 133) 213 KUEPPERS 1971 (A 134) 213 KUEPPERS, FALLAT and LARSON 1969 (A 135) 213 KUPKA and GOLDSMITH 1962 (A 368) KYLE and RIESEN 1970 (269) 62, 59

ndusfrydocuments.ucsf.edu/docs/pfmk0000

LABELLE, BEVILACQUA and Buleger LEWIS and NICHOLLS 1972 (797) 152, 151 1966 (720) 138, 132 LEWIS and KERBY 1965 (A 419) 230 LABELLE, LONG and CHRISTOFANO LEWIS, LAPP and BURRELL 1970 (251) 57, 56 1955 (A 446) 232 LEWIS, LYLE and MILLER 1969 (796) 152, 151 LAENNEC 1819 (34) 19, 14 LICKINT 1956 (A 169) 215 LAKE 1969 (552) 115, 109 LIEBERMAN 1969 (A 138) 214 LAMB and REID 1969 (270) 62, 59 LIEBERMAN, MITTMAN and SCHNEIDER 1969 LAMBERT and REID 1970 (470) 102, 99 (A 139) 214 LANCASTER and TOMASHEFSKI 1963 LIEBOW 1959 (775) 147, 145 (357) 81, 78 LINDALL 1970 (A 250) 220 LANDAU and MORTON 1961 (990) 192 LINDALL, MEDINA and GREIMER 1967 (92) LANGER and FISHER 1956 (A 447) 233 29, 24 LANGSTON: 1972 (A 206) 218 LISTER 1971 (866) 185 LAPP, HANKINSON, BURGESS and LLOYD 1968 (A 490) 236 O'BRIEN 1972 (423) 95, 89 LOB 1970 (A 420) 230 LONG 1964 (1085) 198 LA PRESSE MEDICALE: 1970 (958) 190 LOOSLI 1968 (A 170):215 LARSEN 1970 (A 280) 222 LOPEZ-MAJANO, KIEFFER, MARINE, GARCIA LARSON 1963 (606) 123, 121 and WAGNER 1969 (A 101) 211 LARSON 1965 (35) 19, 14 LORD, GAZIOGLU and KALTREIDER 1969 (A 319) LARSON 1966 (920) 188 224 LARSON and BARMAN 1965 (A 136) 213 LOUDON 1969 (A 448) 233 LARSON, BARMAN, KUEPPERS and LOUDON and SHAW 1967 (A 449) 233 FUDENBERG 1970 (90) 29, 27 LOURENCE, KLIMER and BOROWSKI 1971 LARSON, FUKUDA and MURRAY 1964 (721) 138, 132 (A 464) 234; 1965 (774) 147, 145 LORENZONI 1967 (1020) 193 LAURELL 1971 (921) 188 LOWE 1956 (360) 82, 78 LAURENZI, GUARNERI, ENDRIGA and LOWE 1968 (A 171) 215 CAREY 1963 (271) 62, 60 LOWE, CAMPBELL and KHOSLA 1970 (472) LAWS and HEARD: 1962 (A 53) 208 102, 100 LAWTHER: 1966 (358) 81, 78 LOWE and KHOSLA 1972 (473):102, 100 LEBOVITZ, LEBOVITZ and SILVERMAN LOWE, PELMEAR, CAMPBELL, HITCHENS. 1969 (A 99) 211 KHOSLA and KING 1968 (474) 102, 100 LEBRUN-BRAIS, BLANCHETTE and LOWELL 1959 (361) 82, 79 PROVOST 1971 (A 137) 213 LOWELL, FRANKLIN, MICHELSON and LECKERT 1964 (A 52) 208 SCHILLER 1956 (362) 82, 80 LEFCOE and WONNACOTT 1970 (553) LOWRY and SCHUMAN 1956 (155) 38, 33 115, 104 LUBAN-PLOZZA 1967 (1086) 198 LEGRAND, BOTTO and LESOBRE 1971 LUDWIG and PYLE 1969 (1087) 198 (359) 81, 79 LUNDAR 1965 (867) 185 LEIBESCHUETZ 1959 (471) 102, 97 LUNDMAN 1966 (554) 115, 105 LEMLE, VIEIRA, MILWARD and LUPU and VELICAN 1962 (798):152, 151 MIRANDA 1970 (A 369) 227 LUPU, VELICAN and RACOVEANU 1966 (363) LEMON and WALDEN 1966 (330) 75. 74 82, 79 LEPINE and MYRE 1969 (52) 23, 21 L'UNION MEDICALE DU CANADA 1968 (991) LEUCHTENBERGER and LEUCHTENBERGER 1965 (272) 62, 60 LYNCH (1088) 198 LEUCHTENBERGER and LYONS 1965 (A 54) 208 LEUCHTENBERGER 1970 (A 207) 218 LYONS, RYDER, CAMPBELL and GOUGH 1972 - LEUCHTENBERGER, LEUCHTENBERGER (475) 102, 100 and DOOLIN 1958 (273) 62, 60 LEUCHTENBERGER, LEUCHTENBERGER ZEBRUN and SHAFFER 1960 (274) 62, 60 LEVIN (1150):201 LEVINE, BIGELOW, HAMSTRA, BECKWITT, MITCHELL, NETT.

STEPHEN and PETTY 1967 (A 100) EVINE, TALAMO, SHANNON and KAZEMI Shano dahira / Komining

MACKLEM 1971 (A 55) 208 MACKLEM 1971 (A 172) 215 MACKLEM 1972 (A 450) 233 MAISEL, SILVERS, MITCHELL and PETTY 1968 (A 451) 233 MALISZEWSKI and BASS 1955 (A 491) 236 MALLAN (1151) 201 MANNHART 1962 (555) 115, 107 MARCATILI, BONO and ZUCCHETTO 1967 (185):44, 41 MARCO, MASS, MERANZE, WEINBAUM and KIMBEL 1971 (156) 38, 32 MARKUSH 1969 (331) 75, 73 MARTIN and BOATMAN 1965 (77.6) 147, 145 MARTT 1962 (748) 143, 141 MARX 1966 (A 320) 224 MARX 1971 (A 281) 222 MASIN and MASIN 1966 (220) 51, 49 MASON 1970 (1021) 193 MASS, IKEDA, MERANZE, WEINBAUM and KE MBEL 1972 (157) 38, 32 MASSARO and BOTTOMS: 1967 (209):48, 46 MASSUMI and NUTTER 1968 (A 492) 236 MATHOV 1969 (A 370) 227 MATHUR and MISRA 1972 (556) 115, 109 MATOUSEK 1957 (1152): 201 MAYOCK 1957 (A 102): 212 McCALLUM 1972 (476) 102, 100 McCARROLL, CASSELL, WOLTER, MOUNTAIN, DIAMOND and MOUNTAIN 1967 (234) 54, 52 McCOY and ROBBINS 1963 (A 56) 209 McDERMOTT and COLLINS 1965 (646) 129, 126 McDONALD, BECKLAKE, FOURNIER-MASSEY and ROSSITER 1972 (557) 115, 104 McFADDEN and LINDEN 1972 (647) 129, 125 McFARLAND, GIMBEL, DONALD and FOLKENBERG 1964 (1089) 198 McGRATH and THOMSON 1959 (749) 143. 141 McILREATH and COHEN: 1966 (607) 123, 121 McKEE 1963 (992) 192 McKEE 1971 (A 371) 227 McKENZIE, GLICK and OUTHRED: 1969 (558) 115, 109 McKUSICK and MUTALIK 1968 (93) 30, 27 McLAUGHLIN and TUELLER: 1971 (36) 19, 17 McLAUGHLIN, TYLER and CANADA 1961 (A 493): 236 McLAUGHLIN, TYLER, EDWARDS, CRENSHAW, CANADA, FOWLER, PARKER and REIFENSTEIN 1965 (158):38. 34 McMILLEN (868) 185 McMILLEN (869) 185 MEDICAL ARTS AND SCIENCES 1965

(1090) 118

MEDICAL BULLETIN ON TOBACCO 1967 (994) 192 MEDICAL BULLETIN 1968 (993) 192 MEDICAL NEWS 1968 (1022) 193 MEDICAL NEWS 1969 (870) 185 MEDICAL RESEARCH COUNCIL 1966 (A: 372) 227 MEDICAL TIMES 1970 (959) 190 MEDICAL TRIBUNE (873) 185 MEDICAL TRIBUNE (871):185 MEDICAL TRIBUNE 1972 (872): 185 MEDICAL TRIBUNE (926) 188 MEDICAL TRIBUNE (927) 188 MEDICAL TRIBUNE 1963 (924) 188 MEDICAL TRIBUNE 1970 (925) 188 MEDICAL TRIBUNE 1970 (923) 188 MEDICAL TRIBUNE 1966 (922) 188 MEDICAL TRIBUNE (996) 192 MEDICAL TRIBUNE (995) 192 MEDICAL TRIBUNE 1968 (1034) 195 MEDICAL TRIBUNE (1094) 198 MEDICAL TRIBUNE 1969 (1092) 198 MEDICAL TRIBUNE 1969 (1091) 198 MEDICAL TRIBUNE 1969 (1093): 198 MEDICAL WORLD NEWS 1967 (875) 185 MEDICAL WORLD NEWS 1965 (928) 188 MEDICAL WORLD NEWS 1967 (929) 188 MEDICAL WORLD NEWS 1967 (997) 192 MEDICAL WORLD NEWS 1969 (874): 185 MEDICAL WORLD NEWS 1969 (1023) 193 MEDICAL WORLD NEWS 1970 (930) 188 MEDICINA 1968 (876) 185 MEGAHED 1966 (210) 48, 46 MEGAHED, SENNA, EISSA, SALEH and EISSA 1967 (211) 48, 47 MEINSMA 1967 (A 208) 218 MELLORS 1958 (A 452) 233 MENDEZ, MAURER and FRECKMAN 1963 (A 209) 218 MENDENHALL 1968 (A 494) 236 MENDENHALL and SHREEVE 1937 (722) 138, 134 MENEFEE 1957 (A 282) 222 MERCHANT, KILBURN, O'FALLON, HAMILTON and LUMSDEN: 1972 (424) 95, 90 MERRILL 1963 (A 373) 227 MERRILL 1965 (877) 185 MESERVE (1095): 198 (878) 186 MEURMAN MICZOCH 1965 (A 173) 215 MILLER, BONDURNAT, BRATTON and McILROY 1960 (800) 152, 150 MILLER and BONDURANT 1962 (799) 152, 150 MILLER: 1963 (221) 51, 50 MILLER and KUSCHNER 1969 (A 140) 214 MILLER, ASHCROFT, BEADNELL, WAGNER and PEPYS 1971 (364) 82, 79 MILLER, BREADNELL and ASHCROFT 1968 (365) 82. 79 MILLER and SERJEANT 1971 (A 374) 227

MILLER and SPROULE 1966 (648) 129 MILLER 1964 (1096) 198 MILLER 1967 (A 103) 212 MILLS 1950 (304) 68 MILLS and BAISCH 1965 (A 375) 227 MILNE and WILLIAMSON 1972A (477) 102, 99 MILNE and WILLIAMSON 1972B (478) 102, 99 MITCHELL 1962 (A 453) 233 MITCHELL 1962 (53) 23, 21 MITCHELL 1964 (A 283) 222 g MITCHELL 1966 (960) 190 MITCHELL 1969 (998) 192 MITCHELL, MAISEL, DART and SILVERS 1971 (1024) 193 MITCHELL, RYAN, PETTY and FILLEY 1966 (186) 44, 41 MITCHELL, SILVERS, DART, PETTY, VINCENT, RYAN and FILLEY 1968 (A 284) 222 MITCHELL, TOLL and FILLEY 1962 (A 7) 205 MITCHELL, VINCENT and FILLEY 1964 (54) 23, 21 MITCHELL, WALKER and MAISEL 1968 (782) 149, 148 MITCHELL, WALKER, SILVERS, DART and MAISEL 1969 (A 21):206 MITCHELL, WEBB and FILLEY 1964 (332):75, 73 MITTMAN 1967 (A 104) 212 MITTMAN 1972 (159) 38, 31 MITTMAN 1972 (A 141) 214 MITTMAN, LIEBERMAN, MARASSO and MIRANDA 1971A (94) 30, 26 MITTMAN, LIEBERMAN, MARASSO and MIRANDA 1971B (95) 30, 26 MOERSCH 1968 (1097) 198 MOFFAT and SUTHERLAND 1967 (A 376) MOHANTY and GUPTA 1968: (608): 123, 121 MOLHO, RAZ and KREISLER (A 321) 224 MOLYNEUK and TOMBLESON 1970 (479) 103, 100 MONSON 1970 (A 377) 227 MOOLTEN 1968 (723) 138 MOORE and VINCENT (1098) 198 MORGAN 1971 (801) 152, 151 MORGAN 1972 (A. 210): 218 MORGAN 1968 (1099) 198 MORGAN 1967 (A 174) 215 MORGAN 1968 (A 285) 222 MORGAN 1971 (A 421) 230 MORIYAMA 1963 (1025) 194 MORIYAMA, DAWBER and KANNEL 1966 (1026): 194 MORK 1962 (366): 82, 78

MORTON 1969 (750) 143, 142

MORTON 1944 (751) 143, 142

MORTON and TURNBULL 1968 (A 322) 224 MOTLEY 1960 (A 57) 209 MOTLEY 1971 (235) 54, 52 MOTLEY and KUZMAN 1958 (752) 143, 141 MOTLEY and PHELPS 1964 (A 251) 220 MOTLEY and YANDA 1966 (236) 54, 53 MUELLER 1969 (1100) 198: MUELLER, KEBLE, PLUMMER and WALKER 1971A (425):95, 91 MUELLER, WALKER and PLUMMER 1971B (426):95, 91 MULDOON and WARWICK 1972 (480) 103, 100 MULLER 1959 (A 286) 222 MURPHY, KATZ, MASSARO and LUCHSINGER 1962 (96) 30, 25 MURPHY, FERRIS, BURGESS, WORCESTER and GAENSLER 1971 (A 422) 230 MURPHY, LENG, ULRICH and DAVIS 1963 (A 423) 230 MURRAY 1967 (879) 186 MYDLIL and TOMANEK 1964 (1101) 198 MYRVIK and EVANS 1967 (275) 62, 58 NADEL and COMROE 1961 (649) 129, 126 NADEL, SALEM, TAMPLIN and TOKIWA 1965 (A 454) 233 NAEYE 1972 (427) 95 NAEYE and DELLINGER 1972 (A 424) 230 NAEYE and LAQUEUR 1970 (A 495) 236-NAEYE, MAHON and DELLINGER 1971 (428) 95, 88 NAKASHIMA 1938 (724):138, 133 NAKAJIMA 1966 (367):82, 78 NANDI, HERSHEL, SLONE, SHAPIRO and LEWIS 1969 (160) 38, 34 NASIELL 1968: (222) 51, 49 NEBRASKA STATE MEDICAL JOURNAL 1968 (931) 188 NEGREIROS 1968 (A 175) 215 NERI, MANDEL, O'CONNOR and LARSON 1970 (559) 115, 104 NEW ENGLAND MEDICAL JOURNAL 1963 (1153) 201 NEWMAN, VATTER and REISS 1967 (A 22) 206 NICOLAS 1964 (223):51, 50 NICKLAUS, STOWELL, CHRISTIANSEN and RENZETTI 1966 (A 58) 209 NIDEN 1969 (A 425) 230 NIESENBAUM 1968 (97) 30, 25 NIEWOEHNER, KLEINERMAN and KNOKE 1972 (212) 48, 47 NISHIMOTO, BURROWS, MIYANISHI, KATSUTA, SHIGENOBU and KETTEL 1970 (560) 115, 109 NOMIYAMA, NOMIYAMA and MATSUI 1971 (A 142) 214

NORMAN-TAYLOR and DICKINSON 1972

(1035) 195

(481) 103, 99

NYLAN 1968 (831) 160, 178

NTRDA BULLETIN

**OCHSNER** (1103) 199 OCHSNER (880) 186 **OCHSNER** (1104) 199 OCHSNER 1967 (A 252) 220 OCHSNER 1968 (832) 160, 179 OCHSNER 1968 (1102) 199. OCHSNER 1971 (961) 190 OCHSNER and DAMRAU 1970 (A98) 30, OLSEN, FONTANA, PEABODY and ORDSTRAND 1968 (1036) 195 OLSEN and GILSON: 1960 (482): 103, · 97 ORR 1969 (1105) 199 OSWALD 1964 (A 105) 212 OSWALD 1964 (A 287) 222 OSWALD 1967 (A 106) 212 OSWALD 1968 (932) 188 OSWALD, HAROLD and MARTIN 1953 (483) 103, 97 OSWALD and MEDVEI 1955 (484) 103, 97 OTTO 1969 (A. 23): 206 OTTO and HINUBER 1972 (A 378) 227 OTTO, ZEILHOFER and REISSINGER 1970 (37) 19, 17 OUTHRED and FLYNN 1960 (561) 115, 109 OUTHRED, McKENZIE and WHITE 1970 (A 496) 236 OVERHOLT 1961 (A 211):218 OVERHOLT 1964 (1154) 201 OVERHOLT 1967 (A 212) 218 OVERHOLT, OLIYNYK and CADY (A213) 218 OWSINSKI, JANUS and MYSIK 1971 (562) 115, 107 OYANGUREN, DONOSO, PRIETO, VARGAS and PAZ 1972 115, 110 PAGE 1967 (1106) 199 PAIN, GLAZIER, SIMON and WEST 1967 (753) 144, 142 PALECEK and AVIADO 1967 (650) 129, PALECEK, OSKOUI and AVIADO 1967 (651) 130, 127 PALECEK, PALECEKOVA and AVIADO 1967 (161) 38, 32 PALMER (1107) 199 PALMER and DIAMENT 1970 (A 379) 227 PANZER 1972 (38) 19 PANZER 1972 (933) 188: PARADE 1969 (783) 149, 148 PARAS, MERCADO and QUINTERO 1970 (564) 115, 110 PARAS, MERCADO and QUINTERO 1971 (565) 115, 110 PARNELL, ANDERSON and KINNIS 1966 (252) 57, 56 PASQUAZZIM 1968 (A 107) 212 PATTLE 1958 (276) 62, 53 PATTLE 1965 (277) 62, 58 PATTLE and BURGESS 1957 (A: 426)

PAUL 1968 (934) 188 PAUTRAT 1967 (A 497) 236 PAVIA, SHORT and THOMSON 1970 (725) 138, 131 PAVIA, THOMSON and POCOCK 1971 (726) 138, 131 PAVLIK and CERMAKOVA 1964 (368) 82, 79 PAYNE and KJELSBERG 1964 (429) 95, 90 PEABODY 1962 (1108) 199 PELZER and THOMSON 1966 (652) 130, 125 PEMBERTON 1966 (1027) 194 PEMBERTON 1968 (A 176) 215 PEMBERTON and GOLDBERG 1954 (333) 75, 73 PENMAN 1969 (754) 144 PENMAN 1970 (A 427): 230 PENMAN and ASTIN 1968 (A 323) 224 PENMAN and HOWARD 1966 (755) 144, 142 PENMAN, HOWARD and STENTIFORD 1968 (A 498) 236 PENMAN, O'NEILL and BEGLEY 1970 (A 324) 224 PENNSYLVANIA HEALTH 1965 (1109) 199 PENNSYLVANIA MEDICINE 1969 (1110) 199 PERALES 1971 (A 380) 227 PETERS and MILLERS 1960 (609) 123, 122 PETERS and PRICKMAN 1957 (935) 188 PETERS and FERRIS 1967 (430) 95, 85 PETERS and FERRIS 1967 (431) 95, 85 PETERSON, LONGERGAN and HARDINGE 1968, (610) 123, 122 PETROVA 1956 (566) 115, 108 (1111) 199 PETTY PETTY 1970 (A 108) 212 PETTY, BRINK, MILLER and CORSELLO 1970 (611) 123, 122 PETTY and NETT 1969 (A 109) 212 PETTY, RYAN and MITCHELL 1967 (39) 19, 17 PHELPS and KOIKE 1962 (237) 54, 53 PHILLIPS 1964 (881) 186 PHILLIPS, PHILLIPS and THOMPSON 1956 (A 177) 215 PHILLIPS 1966 (A: 382) 227 PHILLIPS: 1958 (A 499) 236 PHYSICAL THERAPY 1967 (936) 188 PHYSICAL THERAPY 1968 (937) 189 PICK 1960 (1112) 199 PIERCE 1968 (A 325) 224 PIPER 1958 (756) 144 PIRCHER 1972 (A 326) 224 PISTAWKA 1969 (A 327) 224 PLATT and BOINES 1965 (1113) 199 PLATTS, HAMMOND and STUART-HARRIS 1960 (A 500) 236 POCHMON, SKARABELLOVA and LEGOSTOV

1968 (567) 115, 108

POPPIUS 1969 (757) 144, 142 POPPIUS 1970 (882) 186 POSSNER 1970 (568):115, 106 PRATT 1969 (A 24) 206 PRATT, JUTABHA and KLUGH 1963 (A 26) 206 PRATT and KILBURN 1970 (A 25) 206 PRATT and THONG-YAI 1968 (A 288) PRATT, FINLEY, SMITH and LADMAN 1969 (802): 152, 151 PRATT, SMITH, LADMAN and FINLEY 1971 (224):51, 49 PREGOWSHI, BRYSIEWICZ, CESARZ-FRONCZYK, KORDECKA, LESZC-BINSKI, LUKJAN and SADOKIERSKA 1970 (569) 116, 107 PREISER, DONNER and VAN METRE 1969 (369) 82, 79 PRINCI 1963 (A: 253) 220: PRINSLOO and LAUBSCHER: 1970: (570) 116, 108 PROETZ 1939 (278) 62, 58 PUBLIC HEALTH 1971 (883) 186 PUBLIC HEALTH REPORT 1962 (938) 189 PUBLIC HEALTH REPORT 1967 (1114) 199 PUBLIC HEALTH SERVICE PUBLICATION: NO. 1103 1963 (999) 192 PUDELSKI, OKLEK and DELOFF 1971 (571) 116, 107 PUGSLEY (A 289) 222 PUTZ 1969 (1115) 199. QUINLAN, SALMAN SWIFT, WAGNER and PROCTOR 1969 (727) 138, 132 RADELFINGER (1116) 199 RAKIETEN, RAKIETEN, FELDMAN and BOYKIN 1912 (728) 138, 131 RAMIREZ 1969 (1037) 195 RANDHAW 1963 (884) 186 RANKIN (1038) 195 RANKIN, GEE and CHOSY 1965 (758) 144, 141 RANKIN, HALE, WILKINSON, O'DAY, SANTAMARIA and BABARCZY 1969 76. 74 RAO, COHN, ELDRIDGE and HANCOCK 1969 (784) 149, 148 RAPPORT and MAYER 1957 (A 290) 222 RASMUSSEN, BASINGER, HARRISON and MEADER 1964 (A 214) 218 RASMUSSEN 1971 (432):95, 89 RASMUSSEN and NELSON 1971 (433) RASTELLI 1969 (885) 186 RAVENHOLT 1964 (820) 160, 166 RAVENHOLT 1965 (1000): 192 RAY 1965 (55) 23, 21 RAY 1965 (99) 30, 25 READ, LEE and PAIN: 1968 (A 501) 236 READ and SELBY 1961 (612) 124, 119 REBORA 1965 (A 59) 209

REEVES 1962 (A 110) 212 REICHEL 1972 (A 254) 220 REID: 1956 (335) 76 REID, ANDERSON, FERRIS and FLETCHER 1964 (486) 103, 97 REID, CORNFIELD, MARKUSH, SPEIGEL, PEDERSEN and HAENSZEL 1966 (487) 103, 97 REID and FLETCHER: 1971 (315) 71, 69 REID 1959 (A 178) 216 REID 1960 (189) 45, 41 REID 1967 (225) 51, 50 REID and MILLARD 1964 (A 60) 209 REID 1954 (188) 44, 41 REIF, RHODES and COHEN 1970 (162) 38, 31 REGAN, TAGG, WALFORD and THOMSON 1971 (485) 103, 100 REIMANN 1967 (A 179) 216 REINTJES, SEIERENGA and BOGAARD 1972 (653) 130, 126 RENZETTI 1967 (336) 76, 73 RENZETTI, McCLEMENT and LITT 1966 (337) 76, 73 RENZETTI, McCLEMENT and LITT 1966 (1001) 192 RESNICK, LAPP and MORGAN 1971 (A 143) 214 REVOTSKIE, KANNELL, GOLDSMITH and DAWBER 1962 (434) 96, 85 RHODE ISLAND MEDICAL JOURNAL 1966 (939) 189 RICCITELLI 1966 (A 111) 212 RIGDON: 1960 (A 465) 234: RIGDON 1961 (A 215) 218 RIGGINS (A 216) 218 RIMINGTON 1968 (226) 51, 49 RIMINGTON 1969 (488) 103, 99 RIVERO 1966: (A. 27) 206 ROBERTS 1970 (940) 189 ROBERTSON 1969 (654) 130 ROBERTSON 1970 (A 217) 218 ROBBINS, EPSTEIN, DIAMOND, PLISHNER and FINKELSTEIN 1969 (435) 96, 87 ROBBINS 1966 (227) 51, 49 ROCKEY and SPEER: 1966 (163) 38, 32 RODMAN and STERLING 1969 (A 8) 205 RODRIGUEZ, ACHON, RODRIGUEZ. DE LA PAZ, PITA DE LA VEGA. DURAND and SOLLET 1969, (572) 116, 110 ROKSETH, SCHAANNING and AAS 1965 (A 61) 209. ROLLINS 1969 (1155) 201 RONKA 1960 (1117) 199. ROQUE and PICKREN 1968 (228) 51, 49 ROSE and PHILLS 1967 (A 180) 216 ROSEN 1946 (370) 82 ROSEN and LEVY 1950 (371) 82, 79 ROSENBERG 1962 (1118) 199

ROSENBLATT 1969 (941) 189

SANGHVI and KOTIA 1963 (785) 149, 148

SARIC and GOLDSMITH 1964 (A 428) 230

SANO, YODOYAMA and RHOADS 1972

SCHLEUSENER, TALAMO, PARE and

SCHLOTZHAUER and SCHMELTZ 1967

THURLBECK 1968 (A 145) 214

(100):30, 27

(A 468) 234

SCHMIDT 1962 (A 182):216

SAUNDERS 1958 (A 467) 234

SCADDING 1966 (A 112) 212 SCADDING 1969 (A 63) 209

SCHARBORT 1967 (729) 138, 133

(A 328) 224

SIMONNSON, JOCOBS and NADEL 1967

SILEN, BROWN and EISEMAN 1959 (A 385)

SIERING

227

SIMON 1964 (A 64) 209

SIMON 1966 (1157) 202

SIMONSSON 1962 (660) 130, 126

SIMONSSON 1965 (661) 130, 126

SIMONSSON 1965 (214) 48, 46

ROSENBLATT 1970 (338) 76, 72 ROSENBLATT 1972 (339) 76, 72 ROSENBLUM 1964 (836) 186 ROSENKRANTZ, ESBER and SPRAGUE 1969 (803) 153 ROSENKRANTZ and SPRAGUE 1969 (A 466) 234 ROSS 1968 (A: 218) 218: ROSS, McMICHAEL, EADIE, LEES, MURRAY and PINKERTON 1966 (A 181) 216 ROSS, LEY, KRUMHOLZ and RAHBARI 1967 (759) 144, 141 ROWE and ROWE 1965 (372) 82, 79 ROWLANDS, CADENA and GROSS 1967 (804) 153, 151 ROYAL COLLEGE OF PHYSICIANS 1962 (11) 11, 8 ROYAL COLLEGE OF PHYSICIANS: 1971 (12) 11, 8 ROYAL SOCIETY OF HEALTH JOURNAL 1971 (887) 186 RUFF, SALEM, BUSY, de VARNEJOUL, EVEN and BROUET 1972 (655) 130, 125 RUIKKA, SOURANDER and KASANEN 1966 (573) 116, 105 RULLIERE and SAIGOT 1971 (13) 11 RYAN, McDONALD and DEVINE 1955 (190) 45, 43 RYAN: 1968 (1119): 199 RYDER 1966 (1002) 192 RYDER, LYONS, CAMPBELL and GOUGH 1970 (489) 103, 100 RYDER, DUNNILL and ANDERSON 1971 (191) 45, 42 RYDER, THURLBECK and GOUGH: 1969 (A 28) 207 RYLANDER 1968 (279) 62, 59 RYLANDER 1969 (253) 57, 55 RYLANDER 1969 (280) 62, 59. RYLANDER 1970 (14) 11, 8 RYLANDER 1971 (281) 62, 59 SACHS 1972 (805): 153, 151 SACHS and FINLEY 1972 (1039) 195 SACHS 1970 (942):189. **SADOUL 1968 (A. 29): 207** SADOUL, LAZENAIRE, LAMBERT and BERNADA:C 1970 (A 62): 209 SAFFIOTTI, BORG, GROTE and KARP 1964 (A 455) 233 SAHA and JAIN 1970 (56) 23, 21 SAID 1967 (A 502) 236 SAINDELLE, RUFF, GUILLERM and PARROT 1968 (806) 153 SALATA 1966 (A 255): 220 SAMANEK and AVIADO 1965 (656) 130, 127 SAMANEK and AVIADO 1966 (777): 147, 145 SAMANEK, AVIADO and PESKIN 1965 (657) 130, 127 **SANDERS 1970 (A 503) 236** SANDERUD 1958 (192) 45, 42

SCHMIDT 1968 (A 183) 216 SCHOETTLIN 1962 (238) 54, 52 SCHUMAN 1966 (A 383) 227

STONE, GINSBERG, COLLAPINTO and PEARSON 1966 (A 508) 237 STORY 1964 (A 116) 212 STRAWBRIDGE 1960A (167) 38, 31 STRAWBRIDGE 1960B (168) 38, 34 STRAWBRIDGE 1960C (169) 38, 34 (1126) 200 STRESEMANN 1969 (A 258) 220 STRIEDER and KAZEMI 1967 (761) 144, 142 STRIEDER, MURPHY and KAZEMI 1968 (762): 144, 142 STROBEL and GSELL 1965 (579) 116, 107 STRONG, RICHARDS, MCGILL, EGGEN and MCMURRY 1969 (786) 149, 148 STUART-HARRIS 1968 (763) 144 STUART-HARRIS 1971 (A 184):216 STURGIS 1967 (962) 190 STURGIS 1971 (963) 190 STUBER 1967 (1127) 200 SUFRIN 1970 (A 329) 224 SUKUMALCHANGRA, DINAKARA and WILLIAMS 1966 (340) 76, 73 SUKUMALCHANTRA, and WILLIAMS 1965 (A 330) 1965 SUSSMANN 1967 (376) 82, 78 SUTINEN, CHRISTOFORIDIS, KLUGH and PRATT 1965 (A 66) 209 SWARTZ 1971 (15) 11, 8 SWEET, WYATT, FRITSCH and KINSELLA 1961 (40) 20, 14 SWINNY, SWINNY and HICKS 1965 (A 67) 209 SYNDER, LEMON and MANTEL 1965 (A 393) 228 SYZGANOV, GOLOVIN and TROITSKAYA 1971 (170) 38, 32 SZADKOWSKI, SCHULTZE, SCHALLER and LEHNERT 1969 (808) 153, 151 SZCZEKLIK, TERESIAK and KOLACZKOWSKA 1971 (A 148) 214 SZUCS and LORD 1967 (A 117) 212 SZYMCZYKIEWICZ, KUNSKI and GIELEC 1970 (580):116, 107 TAKENOUGHI 1968 (581) 116, 109 TALAMO 1971 (A 149) 214 TALAMO, ALLEN, KAHAN and AUSTEN 1968 (102): 30, 26 TALAMO, AUSTEN and ALLEN 1972 (A 150):214 AMO, BLENNERHASSET and AUSTEN 1966 (A 151) 214 TANNER 1967 (341) 76, 73 TARKOFF, KUEPPERS and MILLER 1968 (A 152): 214. TAUB 1966 (A 291) 222 TAUBERGER 1964 (A 470) 1964 TAYLOR 1963 (964) 190 TAYLOR 1969 (1041):195 TECULESCU: and RACOVEANU 1972 (A 331) 224

TERRELL and SCHMELTZ 1968 (A 471)

234

SPAIN, BRADESS, TARTER and MATERO SPEAR, DAUGHTRY, CHESNEY and MARKS SPICER 1964 (A 256) 220 SPIEGEL and COHEN 1970 (1124) 199 SPODINK, CHUSHMAN, KERR, BLIDE and SPICER 1966 (239):54, 52 **SPOTNITZ** 1966 (A 257) 220 SPURGASH, EHRLICH and PETZOLD (A: 507) 236 STANEK, FODOR, HEJL, WIDIMSKY, CHARVAT, SANTRUCEK, ZAJIC and VAVRIK 1966 (577) 116, 108 STANESCU, GAVRILESCU and TECULESCU **STA**NESCU, PILAT, GAVRILESCU, TECULESCU and CRISTESCU STANESCU, TECULESCU, PAGURARU and GAVRILESCU 1968 (760) 144, 141 STASZEWSKI 1960 (A 220) 218: STEBBINGS 197/1A (438) 96, 89 STEBBINGS 1971B (439) 96, 89 STEBBINGS 197/10 (440) 96, 89 STEBBINGS 197.1D (4-11) 96, 89 STEBBINGS 1972 (A 386) 227 STEBBINGS 1973 (A 387) 228 STEDMAN and MILLER: 1967 (A: 469) 234 STEINFELD 1972 (1158) 202 STEPHENS, FREEMAN and EVANS 1971 STER LING 1967 (662) 130, 126 STEVENS, HNILICA, JOHNSON, BELL

SIMPSON 1968 (215) 48, 46 SIMPSON 1968 (A 65) 209

**19676 (576) 116, 108** 

1967a (575) 116, 108

SMITH, FALK and SISKIND 1971

SMITH, SMITH and McCALL 1960

SNIDER, DOCTOR, DEMAS and SHAW

SOBONYA and KLEINERMAN 1972 (193)

SIRTORI

116, 108

(A 146) 214

(165) 38, 34 **SNIDER** 1968 (A 114) 212

秀日 | 1971 (375) 82, 78

SPAIN 1963 (A 219) 218

SPAIN 1969 (194) 45, 42

1970 (195) 45, 42

1961 (A 115) 212

1968 (282) 62, 60

1968 (615) 124, 120

1967 (578) 116, 108

(166) 38, 33

1971 (101) 30, 26

STEVEN 1971 (A. 147): 214 STEWART 1970 (A 388) 228

STEWART 1967 (1125) 200

1957 (A 429) 230

STOKINGER, WAGNER and DOBROGORSKI

**STAMLER** 

45, 42

(1123): 199

SLUIS-CREMER and SICHEL 1968 (574)

SLUIS-CREMER, WALTERS and SICHEL

SLUIS-CREMER, WALTERS and SICHEL

TERRIS 1968 (1128) 200 TERRY (1129) 200 TERRY 1968 (889) 186 TERRY (16) 11, 8 (1130) 200 THE CHART THE JOURNAL OF SCHOOL HEALTH (1003) 192 THE JOURNAL OF SCHOOL HEALTH (1131) 200: THE LANCET 1969 (945) 189 THE LANCET 1970 (946) 189 THE LANCET 1972 (1132):200 THE MEDICAL JOURNAL OF AUSTRALIA 1964 (947) 189 THE MEDICAL JOURNAL OF AUSTRALIA 1970 (1133) 200 THE MEDICAL JOURNAL OF AUSTRALIA 1971 (890): 186 THE MEDICAL OFFICER 1969 (965) 190 THOMSON and PAVIA 1973 (730) 138, 131 THOMSON 1970 (A 9) 205 THURLBECK (A 292) 222 THURLBECK 1963 (171):38, 31 THURLBECK 1963 (41) 20, 17 THURLBECK 1966 (A 30) 207 THUR LBECK 1970 (A 31) 207 THURLBECK 1970 (A 32) 207 THURLBECK 1971 (A 332) 224 THURLBECK and ANGUS 1963 (A 293) 222 THURLBECK and ANGUS 1964 (196) 45, 42 THURLBECK and ANGUS 1967 (197) THURLBECK, ANGUS and PARE 1963 (199) 45, 42 THURLBECK, BATES, FRASER and HENDERSON 1968 (A 68) 209. THURLBECK, FRASER and BATES 1965 (198) 45, 41 THURLBECK, HENDERSON, FRASER and BATES 1970 (A 294) 222 **TOBACCO REPORTER 1967 (1043) 195** TOBACCO REPORTER 1967 (1042) 195 TOKUHATA, DESSAUER, PENDERGRASS, HARTMAN, DIGON and MILLER 1970, (442) 96, 88 TOKUHATA 1972 (A 221) 218 TOLL 1964 (A 69): 209 TOMASHEFSKI and PRATT 1967 (A 10) 205 TOWNLEY, RYNING, LYNCH and BRODY 1970 (A 153) 214 TRINQUET and MEYER: 1971 (A. 185) 216 TSUMETOSHI, SHIMIZU, TAKAHASHI, ICHINOSAWA, UEDA, NAKAYAMA, YAMAGATA and Olishino 1971 (582) 116, 109 TUBERCULOSIS ASSOCIATION OF LIARYLAND 1968 (948) 189: TURINO 1971 (A 154) 214

page 256 TURNER 1969 (A 295) 222 TYSINGER 1970 (A 333) 224 ULMER 1970 (583) 116, 106 ULMER 1972 (A 118) 212 ULMER and REIF 1966 (586) 117, 106 ULMER, REICHEL and WERNER 1968 (585) 117, 106 ULMER and REICHEL 1970 (584) 117, 106 U.S. DEPARTMENT OF HEALTH, EDUCATION AND WELFARE 1964 (17) 11, 8 U.S. DEPARTMENT OF HEALTH, **EDUCATION AND WELFARE 1965** (823) 160, 169 U.S. DEPARTMENT OF HEALTH, **EDUCATION AND WELFARE 1967** (18) 11, 8 U.S. DEPARTMENT OF HEALTH, **EDUCATION AND WELFARE 1967** (891) 186 U.S. DEPARTMENT OF HEALTH, **EDUCATION AND WELFARE 1968** U.S. DEPARTMENT OF HEALTH **EDUCATION AND WELFARE 1969** (20) 12, 8 U.S. DEPARTMENT OF HEALTH **EDUCATION AND WELFARE 1971** (21) 12, 8 U.S. DEPARTMENT OF HE AL TH **EDUCATION AND WELFARE 1973** (22) 12, 8 U.S. MEDICINE 1960 (1134) 200 U.S. MEDICINE 1968 (966) 190 U.S. MEDICINE 1968 (1028) 194 U.S. MEDICINE 1969 (893) 186 VALIC and ZUSKIN 1972 (587) 117, 108 VALENTIN and WOITOWITZ 1967 (A 296). 222 VAN DER WAL, HUIZINGA, ORIE, SLUITER and DE VRIES 1966 (A 222) 218 VAN GANSE, DEPOORTER, MEHEUS, EYLENBOSCH, VANHOORNEEN, VUYLSTEEK, DE VREEZE and RAVEEL 1972 (764) 144, 141 VARGHA 1969 (A 334) 224 VARPELA and SARIS 1971 (A 155) 214 VASSAR, CULLING and SAUNDERS 1960 (229) 51, 49 IDAL, CAZAL, ROBINET-LEVY and MICHEL 1971 (A 156) 214 VIGY 1970 (588) 117, 106 VIIKERI, JAASKELAINEN and TAHTI 1968 (A 430) 231 VINCENT 1967 (892) 186 VISWANATHAN 1965 (778) 147, 145 VISWANATHAN, MODY, PRASAD and SINHA 1965 (377) 83, 78 VOISIN 1970 (A 297) 222 VON: OETTINGEN 1941 (17/2) 39,: 33 VOORST VADER 1965 (894) 186

VOTCHAT, LOZINSKY, GRIBOVA, TAGAI and SHEFER 1970 (663) 130

YUGA: 1968 (A 509): 237-VYSKOCIL 1968 (589) 117, 108 WADDELL 1966 (1004) 192 WAKERLIN 1964 (821) 160, 167 WALKER, ARCHIBALD and ATTFIELD 1971 (490): 103, 100 WALKER 1917 (A 389): 228 WALKER and KIEFER 1966 (731) 138, 134 WALLNER 1963 (23) 12, 8 WALLER 1971 (A 259) 220 WALSHE and HAYES 1967 (590):117, 110 WATERMAN, DOMM, ROGERS and BORRELL 1969 (A 119) 212 WARE 1964 (895) 186 WARING 1969 (1135) 200 WARTHEN 1964 (1159) 202 WEBB, COCK, LANIUS and SHAW 1967 (809) 153, 150 WEBSTER, KETTEL, MORAN, BUGAIESKI and CUGELL 1968 (103) 30, 24 WEHRLE, BRENT, DOYLE, FARR, FAGAN, FINBERG, NAHMIAS, PICKERING and YAMAZAKI 1969 (A 390) 228 WEIR and DUNN: 1970 (305):68, 65 WEISS 1967 (378) 83, 79 WEISS 1968 (896) 186 WEISS 1968 (732) 138, 134 WEISS 1969 (379) 83, 79 WEISS 1971 (380) 83, 79 WEISS, BOUCOT, COOPER and CARNAHAN 1963 (616) 124, 121 WEISS and DULFANO 1968 (765) 144, 142 WEISSBECKER, CREAMER and CARPENTER 1971 (733) 138, 133 WEISSBECKER, CARPENTER, LUCHSINGER and OSDENE 1969 (283) 62, 58 WELCH, REINECKE, HAMMARSTEN and GUENTER 1969 (A 157) 214 WELLER 1955 (200) 45, 42 WENDOHL and FACKLER 1968 (949) 189 WENDEL 1966 (591) 117, 106 WENDEL 1968 (216) 48, 46 WESCHESTER MEDICAL BULLETIN 1969 (1136) 200. WESSEL-ASS 1967 (A 298) 222 WEST, GLAZIER, HUGHES and MALONEY 1969 (A 510) 237. WEST, BURNS, DANIEL and JACKSON 1959 (A 391) 228 WESTERMARK 1961 (A: 472) 234 WHATLEY 1967 (826): 160, 173 WIGHTMAN 1968 (766) 144, 142 WIKLANDER and NORLIN 1957 (767) 144, 142 WILHELMSEN 1967 (664) 130, 125 WILHELMSEN, ORHA and TIBBLIN 1969 (592):117, 105 WILHELMSEN and TIBBLIN 1966 (593) 117, 105 WILKELSTEIN, KANTOR, DAVIS, MANERI and MOSHER 1967 (342):76, 73 WILLIAMS and ANDERSON 1968a (768) 144,

page: 257 WILLIAMS and ANDERSON 1968b (769) 144, 141 WILSON, MEADOR, JAY and HIGGINS 1960 (617) 124, 122 WINKELSTEIN and KANTOR 1969 (240):54, 52 WINKELSTEIN and FRENCH 1970 (A: 260) 220 WINKENWERDER, ALLAN and FISHER 1962 (381) 83, 79 WOOLCOCK, BLACKBURN, FREEMAN, ZYLSTRA and SPRING 1970 (594) 117, 109 WOOLCOCK, VINCENT and MACKLEM 1969 (665) 130, 125 WOOLF and SUERO 1971 (618) 124, 121 WOLFF, PURDOM and ISENHOWER 1954 (A 457) 223 WORLD HEALTH ORGANIZATION 1970 (1137) 200 WORTH, MUYSERS and SMIDT 1968 (A 33) 207 WRIGHT 1965: (A. 70) 209. WRIGHT and KLEINERMAN: 1963 (A 11) 205 WRIGHT 1968 (A 34):207 WYATT and FISCHER 1964 (A 35) 207 WYNDER, GOODMAN and HOFFMANN 165 (7.34) 138, 134, 133 WYNDER, GOODMAN and HOFFMANN 1965 (735) 139, 134 WYNDER and HOFFMANN 1970 (A 223) 218 WYNDER: and HOFFMANN: 1972 (897) WYNDER, KAISER, GOODMAN and HOFFMANN 1963 (736) 139, 134 WYNDER, KAUFMAN and LESSER 1967 (619) 124, 122 WYNDER, LEMON and MANTEL 1965 (A 392) 228 WYNDER, TAGUCHI, BADEN and HOFFMANN 1968 (284) 62, 60 YAMANAKA 1970 (42) 20, 17 YANDA 1965 (A 120) 212 YANDA and MOTLEY 1965 (A 121) 212 YEAGER 1969 (285) 63, 58 ZAGONA 1967. (1138): 200 ZASLY, BAUM and RUMBALL 1960 (A 394) 228 ZEIDBERG, HORTON and LANDAU 1967 (343) 76, 73 ZEIDBERG, PRINDLE and LANDAU 1964 (443) 96, 90 ZIVY 1970 (382) 83, 79 ZIVY and FOURNIER 1970 (383) 83, 79 ZWI, GOLDMAN and LEVIN 1964 (620) 124. 121 ZWI. SLUIS-CREMER and DU PREEZ 1967

(595) 117, 108

3